Name

## CAMBRIDGE INTERNATIONAL EXAMINATIONS International General Certificate of Secondary Education

0620/03 **CHEMISTRY** 

Paper 3

May/June 2003

1 hour 15 minutes

Candidates answer on the Question Paper. No Additional Materials required.

## **READ THESE INSTRUCTIONS FIRST**

Write your name, Centre number and candidate number in the spaces provided at the top of this page. Write in dark blue or black pen in the spaces provided on the Question Paper. You may use a pencil for any diagrams, graphs or rough working.

Do not use staples, paper clips, highlighters, glue or correction fluid.

Answer all questions.

The number of marks is given in brackets [ ] at the end of each question or part question.

A copy of the Periodic Table is provided on page 12.

If you have been given a label, look at the details. If any details are incorrect or missing, please fill in your correct details in the space given at the top of this page.

Stick your personal label here, if provided.

For Exam	iner's Use
1	
2	
3	
4	
5	
TOTAL	

This document consists of 12 printed pages.




- No one knows where iron was first isolated. It appeared in China, the Middle East and in Africa. It was obtained by reducing iron ore with charcoal.
  - (a) Complete the following equation.

(c)

[2]

**(b)** In 1705 Abraham Darby showed that iron ore could be reduced using coke in a blast furnace.



(1)	exothermic reaction that causes this high temperature.
(ii)	In the furnace, the ore is reduced by carbon monoxide. Explain how this is formed.
	[3]
	formation of slag removes an impurity in the ore. Write a word equation for the nation of the slag.

(d)	Stai carb	nless steel is an alloy of iron. It contains iron, other metals and about 0.5% of oon.
	(i)	State a use of stainless steel.
	(ii)	Name a metal, other than iron, in stainless steel.
	(iii)	The iron from the blast furnace is impure. It contains about 5% of carbon and other impurities, such as silicon and phosphorus. Describe how the percentage of carbon is reduced and the other impurities are removed.
		[6]
(e)		of the methods used to prevent iron or steel from rusting is to electroplate it with ther metal, such as tin. Complete the following.
	The	anode is made of
	The	cathode is made of
	The	electrolyte is a solution of

0620/03/M/J/03 [Turn over

2 Calcium and other minerals are essential for healthy teeth and bones. Tablets can be taken to provide these minerals.

## **Healthy Bones**

## Each tablet contains

calcium

magnesium

zinc

copper

boron

(a)	Bor	on is a non-metal with a macromolecular structure.
	(i)	What is the valency of boron?
	(ii)	Predict <b>two</b> physical properties of boron.
	(iii)	Name another element and a compound that have macromolecular structures.
		element
		compound
	(iv)	Sketch the structure of one of the above macromolecular substances.

(b)		cribe the reactions, if any, of zinc and copper(II) ions with an excess of aqueous um hydroxide.
	(i)	zinc ions
		addition of aqueous sodium hydroxide
		excess sodium hydroxide
	(ii)	copper(II) ions
		addition of aqueous sodium hydroxide
		excess sodium hydroxide
		[4]
(c)	Eac read	n tablet contains the same number of moles of $CaCO_3$ and $MgCO_3$ . One tablet ted with excess hydrochloric acid to produce 0.24 dm <sup>3</sup> of carbon dioxide at r.t.p.
		$\begin{array}{l} CaCO_3 \ + \ 2HCl \ \to \ CaCl_2 \ + \ CO_2 \ + \ H_2O \\ MgCO_3 \ + \ 2HCl \ \to \ MgCl_2 \ + \ CO_2 \ + \ H_2O \end{array}$
	(i)	Calculate how many moles of CaCO <sub>3</sub> there are in one tablet.
		number of moles CO <sub>2</sub> =
		number of moles of CaCO <sub>3</sub> and MgCO <sub>3</sub> =
		number of moles of $CaCO_3$ =[3]
	(ii)	Calculate the volume of hydrochloric acid, $1.0\mathrm{mol}/\mathrm{dm^3}$ , needed to react with one tablet.
		number of moles of $CaCO_3$ and $MgCO_3$ in one tablet =
		number of moles of HCl needed to react with one tablet =
		volume of hydrochloric acid, 1.0 mol/dm³, needed to react with one tablet =

[4]

3	Alkenes are unsaturated	l hydrocarbons.	They undergo addition	on reactions.

- (a) Two of the methods of making alkenes are cracking and the thermal decomposition of chloroalkanes.
  - (i) Complete an equation for the cracking of the alkane, decane.

 $C_{10}H_{22} \rightarrow \dots + \dots$  decane

(ii) Propene can be made by the thermal decomposition of chloropropane. Describe how chloropropane can be made from propane.

reagents propane and ......

conditions

**(b)** The following alkenes are isomers.

$$\begin{array}{ccc} \mathrm{CH_3-CH_2-CH=CH_2} & & \mathrm{CH_3-C=CH_2} \\ & & \mathrm{CH_3} \end{array}$$

(i) Explain why they are isomers.

(ii) Give the name and structural formula of another hydrocarbon that is isomeric with the above.

name .....


structural formula

(c) Give the name of the product when but-1-ene reacts with each of the following.

steam .....

hydrogen .....

- (d) Alkenes can polymerise.
  - (i) Deduce the name and structural formula of the monomer from the structure of the polymer.



name of monomer .....

structural formula

(ii) Draw the structure of the polymer formed from the following monomer.

(iii) Describe the pollution problems caused by the disposal of polymers in landfill sites and by burning.

landfill sites	 	 	
	 	 	[2]

burning .....

\_\_\_\_\_[1]


- 4 Nitrogen dioxide, NO<sub>2</sub>, is a dark brown gas.
  - (a) Most metal nitrates decompose when heated to form the metal oxide, nitrogen dioxide and oxygen.
    - (i) Write a symbol equation for the decomposition of lead(II) nitrate.

$$Pb(NO_3)_2 \rightarrow \dots + \dots + \dots + \dots$$
 [2]

(ii) Potassium nitrate does not form nitrogen dioxide on heating. Write the word equation for its decomposition.

.....[1]

**(b)** When nitrogen dioxide is cooled, it forms a yellow liquid and then pale yellow crystals. These crystals are heated and the temperature is measured every minute. The following graph can be drawn.



(i) Describe the arrangement and movement of the molecules in the region A-B.

	(ii)	Name the change that occurs in the region <b>B</b> – <b>C</b>
		[4]
(c)	Nitro	ogen dioxide and other oxides of nitrogen are formed in car engines.
	(i)	Explain how these oxides are formed.
	(ii)	How are they removed from the exhaust gases?
		[4]
d)	Des	ogen dioxide, oxygen and water react to form dilute nitric acid. Scribe how lead(II) nitrate crystals could be prepared from dilute nitric acid and $d(II)$ oxide.
		[3]

5

		t three elements in Period and lanthanum.	6 of the Periodic Tal	ble of the Elements are	caesium,
(a	tha	w many <b>more</b> protons, electric in one atom of caesium. It is you.			
	nur	nber of protons			
	nur	nber of electrons			
	nur	nber of neutrons			[3]
(b	•	three metals can be obtaine he aqueous halides does no	-	of a molten halide. The ele	ectrolysis
	(i)	Complete the equation felectrode (cathode).	or the reduction of	lanthanum ions at the	negative
		La <sup>3+</sup> +	→		
	(ii)	Name the <b>three</b> products f		sis of aqueous caesium l	
(с	) All	three metals react with cold	water. Complete the	word equation for these re	eactions.
		metal + water $\rightarrow$	+		[2]
(d	cor aro	rium chloride is an ionic cor npound, the charges on the und the negative ion. e electron distribution of a ba	ions and gives the arr	angement of the valency	
		e x to represent an electron			

Use o to represent an electron from a chlorine atom.

(e)		cribe, by means of a simple diagram, the lattice structure of an ionic compount as caesium chloride.	∩d,
			[2]
(f)	The	reactions of these metals with oxygen are exothermic.	
		$2Ba(s) + O_2(g) \rightarrow 2BaO(s)$	
	(i)	Give an example of bond forming in this reaction.	
	(ii)	Explain using the idea of bond breaking and forming why this reaction exothermic.	is
			.[3]

DATA SHEET

				I	_			,		_
		0	4 <b>He</b> Helium	20 <b>Ne</b> Neon 10	40 <b>Ar</b> Argon	84 <b>Kr</b> Krypton 36	131 <b>Xe</b> Xenon Xenon 54	Rn Radon 86		ļ
		<b>=</b>		19 Fluorine	35.5 <b>C1</b> Chlorine	80 <b>Br</b> Bromine 35	127 <b>I</b> lodine 53	At Astatine 85		,
		5		16 Oxygen	32 <b>S</b> Sulphur 16	79 Selenium 34	128 <b>Te</b> Tellurium 52	<b>Po</b> Polonium 84		-
		>		14 Nitrogen	31 Phosphorus 5	75 <b>AS</b> Arsenic	122 <b>Sb</b> Antimony	209 <b>Bi</b> Bismuth 83		
		≥		12 Carbon 6	28 <b>Si</b> Silicon	73 <b>Ge</b> Germanium 32	<b>Sn</b>	207 <b>Pb</b> Lead		
		≡		11 Boron 5	27 A1 Aluminium	70 <b>Ga</b> Gallium 31	115 <b>In</b> Indium 49	204 <b>T t</b> Thallium		
ts						65 <b>Zn</b> Zinc 30	112 <b>Cd</b> Cadmium 48	201 <b>Hg</b> Mercury 80		
The Periodic Table of the Elements						64 Copper 29	108 <b>Ag</b> Silver	197 <b>Au</b> Gold 79		ļ
e of the	Group					59 <b>X</b> Nickel	106 <b>Pd</b> Palladium 46	195 <b>Pt</b> Platinum 78		,
dic Tabl	Gre					59 <b>Co</b> Cobatt	103 <b>Rh</b> Rhodium 45	192 <b>Ir</b> Iridium 77		
he Perio			1 <b>X</b> Hydrogen			56 <b>Fe</b> Iron	Ruthenium 44	190 <b>OS</b> Osmium 76		
F						55 Wn Manganese 25	Tc Technetium 13	186 <b>Re</b> Rhenium 75		-
						Cr Chromium 24	96 <b>Mo</b> Molybdenur 42	184 <b>W</b> Tungsten 74		-
						51 V Vanadium 23	93 Nobium 41	181 <b>Ta</b> Tantalum 73		-
						48 <b>Ti</b> Titanium	91 <b>Zr</b> Zirconium 10	178 <b>Hf</b> Hafnium 72		
						Scandium 21	89 <b>×</b>	139 <b>La</b> Lanthanum 57 * 7	227 <b>Ac</b> Actinium 89	
		=		9 Beryllium	24 Mg Magnesium	40 <b>Ca</b> Calcium	St Strontium	137 <b>Ba</b> Barium 56	226 <b>Ra</b> Radium R8	
		_		7 <b>L</b> :thium	Na Sodium	39 K	Rubidium 37	133 <b>Cs</b> Caesium 55	<b>Fr</b> Francium 87	
							3/M/J/03			

מסומסי ליוסים	140	141	144		150	152	157		162		167	169	173	
iola senes	ပီ	Ŗ	PZ	Pm	Sm	Ē	Вg	Тр	۵	우	ш	Ε	ΛÞ	Ľ
Ju selles	Cerium	Praseodymium	Neodymium	Promethium	Samarium	Europium	Gadolinium		Dysprosium		Erbium	Thulium	Ytterbium	
Г	58	29	09	61	62	63	64	65	99	67	89	69	70	
a = relative atomic mass	232		238											
X = atomic symbol	Т	Ра	<b>-</b>	Ν	Pu	Am	S	路	₽	Es	Fm	Md	8	בֿ
b = proton (atomic) number	Thorium	Protactinium 0-1	Uranium	Neptunium	Plutonium 94	Americium os	Curium	Berkelium o7	Californium	Einsteinium	Fermium	Mendelevium	Nobelium	Lawrencium
7	8	5	36	200	100	6	00	91	000	00	3	2	102	2

в **X** 

Key

\*58-71 Lanthanoid series †90-103 Actinoid series

The volume of one mole of any gas is 24 dm<sup>3</sup> at room temperature and pressure (r.t.p.).