

Centre Number

Candidate Number

Name

UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS  
International General Certificate of Secondary Education

**CHEMISTRY****0620/03**

Paper 3

May/June 2004

**1 hour 15 minutes**

Candidates answer on the Question Paper.  
No Additional Materials required.

**READ THESE INSTRUCTIONS FIRST**

Write your Centre number, candidate number and name on all the work you hand in.  
Write in dark blue or black pen in the spaces provided on the Question Paper.  
You may use a pencil for any diagrams, graphs or rough working.  
Do not use staples, paper clips, highlighters, glue or correction fluid.  
You may use a calculator.

Answer **all** questions.

The number of marks is given in brackets [ ] at the end of each question or part question.  
A copy of the Periodic Table is printed on page 12.

**For Examiner's Use**

1

2

3

4

5

6

7

**Total**

If you have been given a label, look at the details. If any details are incorrect or missing, please fill in your correct details in the space given at the top of this page.

Stick your personal label here, if provided.

This document consists of **12** printed pages.



1 It was reported from America that a turbine engine, the size of a button, might replace batteries. The engine would be built from silicon which has suitable properties for this purpose.

(a) (i) Why are batteries a convenient source of energy?

..... [1]

(ii) The engine will run on a small pack of jet fuel. What other chemical is needed to burn this fuel?

..... [1]

(b) Silicon has the same type of macromolecular structure as diamond.

(i) Explain why one atom of either element can form four covalent bonds.

.....  
..... [2]

(ii) Predict **two** physical properties of silicon.


.....  
..... [2]

(iii) Name a different element that has a similar structure and properties to silicon.

..... [1]

(c) Silicon is made by the carbon reduction of the macromolecular compound, silicon(IV) oxide.

(i) Balance the equation for the reduction of silicon(IV) oxide.



(ii) Explain why the silicon(IV) oxide is said to be reduced.


..... [1]

(iii) Describe the structure of silicon(IV) oxide. You may use a diagram.

..... [2]

2 Sulphur is used to make sulphuric acid. In the UK, the annual production of the acid is about 2.5 million tonnes.

(a) The reactions in the manufacture of sulphuric acid by the Contact Process are shown below.



(i) Give a large scale source of the element sulphur.

..... [1]

(ii) State another use of sulphur dioxide.

..... [1]

(iii) How is sulphur changed into sulphur dioxide?

..... [1]

(iv) Name the catalyst used in reaction 2.

..... [1]

(v) Reaction 2 is exothermic. Why is a catalyst, rather than a higher temperature, used to increase the rate of this reversible reaction?

..... [2]

(vi) Write a word equation for reaction 3.

..... [1]

(vii) Write a symbol equation for reaction 4.

..... [1]

(b) About one third of this production of acid is used to make nitrogen and phosphorus-containing fertilisers.

(i) Name the third element that is essential for plant growth and is present in most fertilisers.

..... [1]


(ii) Name a nitrogen-containing fertiliser that is manufactured from sulphuric acid.

..... [1]

(iii) Rock phosphate (calcium phosphate) is obtained by mining. It reacts with concentrated sulphuric acid to form the fertiliser, superphosphate. Predict the formula of each of these phosphates.

fertiliser	ions	formula
calcium phosphate	$\text{Ca}^{2+}$ and $\text{PO}_4^{3-}$	.....
calcium superphosphate	$\text{Ca}^{2+}$ and $\text{H}_2\text{PO}_4^-$	..... [2]

(iv) The ionic equation for the reaction between the phosphate ion and sulphuric acid is shown below.



Explain why the phosphate ion is described as acting as a base in this reaction.

..... [2]

3 An organic compound decomposes to form nitrogen.



(a) Explain the state symbols.

aq .....


l .....

g ..... [2]


(b) Draw a diagram to show the arrangement of the valency electrons in **one** molecule of nitrogen.

[2]

(c) The rate of this reaction can be measured using the following apparatus.



The results of this experiment are shown on the graph below.



(i) How does the rate of this reaction vary with time?

.....  
 ..... [1]

(ii) Why does the rate vary?

.....  
 ..... [2]


(iii) The reaction is catalysed by copper powder. Sketch the graph for the catalysed reaction on the same grid. [2]

(iv) Why is copper powder more effective as a catalyst than a single piece of copper?

..... [1]

4 (a) Insoluble compounds are made by precipitation.

(i) Complete the word equation for the preparation of zinc carbonate.




(ii) Complete the following symbol equation.



(iii) Write an ionic equation for the precipitation of the insoluble salt, silver(I) chloride.



(b)  $2.0 \text{ cm}^3$  portions of aqueous sodium hydroxide were added to  $4.0 \text{ cm}^3$  of aqueous iron(III) chloride. Both solutions had a concentration of  $1.0 \text{ mol/dm}^3$ . After each addition, the mixture was stirred, centrifuged and the height of the precipitate of iron(III) hydroxide was measured. The results are shown on the following graph.



(i) Complete the ionic equation for the reaction.



(ii) On the same grid, sketch the graph that would have been obtained if iron(II) chloride had been used instead of iron(III) chloride? [2]

- (iii) If aluminium chloride had been used instead of iron(III) chloride, the shape of the graph would be different. How are the shapes of these two graphs different and why?

difference in shape .....

.....

reason for difference .....

..... [2]

- 5 (a) Copper has the structure of a typical metal. It has a lattice of positive ions and a "sea" of mobile electrons. The lattice can accommodate ions of a different metal.

Give a **different** use of copper that depends on each of the following.

- (i) the ability of the ions in the lattice to move past each other

..... [1]


- (ii) the presence of mobile electrons

..... [1]

- (iii) the ability to accommodate ions of a different metal in the lattice

..... [1]

- (b) Aqueous copper(II) sulphate solution can be electrolysed using carbon electrodes. The ions present in the solution are as follows.



- (i) Write an ionic equation for the reaction at the negative electrode (cathode).

..... [1]

- (ii) A colourless gas was given off at the positive electrode (anode) and the solution changes from blue to colourless.

Explain these observations.

.....

..... [2]

(c) Aqueous copper(II) sulphate can be electrolysed using copper electrodes. The reaction at the negative electrode is the same but the positive electrode becomes smaller and the solution remains blue.

(i) Write a word equation for the reaction at the positive electrode.

..... [1]

(ii) Explain why the colour of the solution does not change.


.....  
..... [2]

(iii) What is the large scale use of this electrolysis?

..... [1]




- 6 In 2002, Swedish scientists found high levels of acrylamide in starchy foods that had been cooked above 120 °C. Acrylamide, which is thought to be a risk to human health, has the following structure.



- (a) (i) It readily polymerises to polyacrylamide. Draw the structure of this polymer.

[2]

- (ii) Starch is formed by polymerisation. It has a structure of the type shown below. Name the monomer.



[1]

- (iii) What are the differences between these two polymerisation reactions, one forming polyacrylamide and the other starch?

[2]

- (b) Acrylamide hydrolyses to form acrylic acid and ammonium ions.

- (i) Describe the test for the ammonium ion.

test

.....  
 .....


result

..... [2]


- (ii) Given an aqueous solution, concentration 0.1 mol / dm<sup>3</sup>, how could you show that acrylic acid is a weak acid.

.....  
 ..... [2]

- (c) The structural formula of acrylic acid is shown below. It forms compounds called acrylates.



- (i) Acrylic acid reacts with ethanol to form the following compound.



Deduce the name of this compound. What type of organic compound is it?

name .....

type of compound ..... [2]

- (ii) Acrylic acid is an unsaturated compound. It will react with bromine. Describe the colour change and draw the structural formula of the product of this addition reaction.

colour change .....

structural formula of product


[2]

- 7 Chemists use the concept of the mole to calculate the amounts of chemicals involved in a reaction.

(a) Define *mole*.

..... [1]

(b) 3.0 g of magnesium was added to 12.0 g of ethanoic acid.



The mass of one mole of Mg is 24 g.

The mass of one mole of CH<sub>3</sub>COOH is 60 g.

(i) Which one, magnesium or ethanoic acid, is in excess? You must show your reasoning.

..... [3]


(ii) How many moles of hydrogen were formed?

..... [1]

(iii) Calculate the volume of hydrogen formed, measured at r.t.p.

..... [2]

(c) In an experiment, 25.0 cm<sup>3</sup> of aqueous sodium hydroxide, 0.4 mol/dm<sup>3</sup>, was neutralised by 20.0 cm<sup>3</sup> of aqueous oxalic acid, H<sub>2</sub>C<sub>2</sub>O<sub>4</sub>.



Calculate the concentration of the oxalic acid in mol/dm<sup>3</sup>.

(i) Calculate the number of moles of NaOH in 25.0 cm<sup>3</sup> of 0.4 mol/dm<sup>3</sup> solution.

..... [1]

(ii) Use your answer to (i) and the mole ratio in the equation to find out the number of moles of H<sub>2</sub>C<sub>2</sub>O<sub>4</sub> in 20 cm<sup>3</sup> of solution.

..... [1]

(iii) Calculate the concentration, mol/dm<sup>3</sup>, of the aqueous oxalic acid.

..... [2]

**DATA SHEET**  
**The Periodic Table of the Elements**

Group																																																																																																						
I	II	III	IV	V	VI	VII	0																																																																																															
7 <b>Li</b> Lithium 3	9 <b>Be</b> Beryllium 4	1 <b>H</b> Hydrogen 1	12 <b>C</b> Carbon 6	14 <b>N</b> Nitrogen 7	16 <b>O</b> Oxygen 8	19 <b>F</b> Fluorine 9	20 <b>Ne</b> Neon 10	11 <b>B</b> Boron 5	13 <b>Al</b> Aluminium 13	14 <b>Si</b> Silicon 14	15 <b>P</b> Phosphorus 15	16 <b>S</b> Sulphur 16	17 <b>Cl</b> Chlorine 17	18 <b>Ar</b> Argon 18	27 <b>Al</b> Aluminium 13	28 <b>Si</b> Silicon 14	31 <b>P</b> Phosphorus 15	32 <b>Ge</b> Germanium 32	33 <b>As</b> Arsenic 33	34 <b>Se</b> Selenium 34	35 <b>Br</b> Bromine 35	36 <b>Kr</b> Krypton 36	49 <b>In</b> Indium 49	50 <b>Tl</b> Thallium 81	51 <b>Sb</b> Antimony 51	52 <b>Te</b> Tellurium 52	53 <b>I</b> Iodine 53	54 <b>Xe</b> Xenon 54	81 <b>Fr</b> Francium 87	82 <b>Ra</b> Radium 88	83 <b>Bi</b> Bismuth 83	84 <b>Po</b> Polonium 84	85 <b>At</b> Astatine 85	86 <b>Rn</b> Radon 86	87 <b>Fr</b> Francium 87	88 <b>Ra</b> Radium 88	89 <b>Ac</b> Actinium 89	90 <b>Th</b> Thorium 90	91 <b>Pa</b> Protactinium 91	92 <b>U</b> Uranium 92	93 <b>Np</b> Neptunium 93	94 <b>Pu</b> Plutonium 94	95 <b>Am</b> Americium 95	96 <b>Cm</b> Curium 96	97 <b>Bk</b> Berkelium 97	98 <b>Cf</b> Californium 98	99 <b>Es</b> Einsteinium 99	100 <b>Fm</b> Fermium 100	101 <b>Md</b> Mendelevium 101	102 <b>No</b> Nobelium 102	103 <b>Lr</b> Lawrencium 103	104 <b>Rf</b> Rutherfordium 104	105 <b>Db</b> Dubnium 105	106 <b>Sg</b> Seaborgium 106	107 <b>Bh</b> Bohrium 107	108 <b>Hs</b> Hassium 108	109 <b>Mt</b> Meitnerium 109	110 <b>Ds</b> Darmstadtium 110	111 <b>Rg</b> Roentgenium 111	112 <b>Cn</b> Copernicium 112	113 <b>Nh</b> Nihonium 113	114 <b>Fl</b> Flerovium 114	115 <b>Mc</b> Moscovium 115	116 <b>Lv</b> Livermorium 116	117 <b>Ts</b> Tennessine 117	118 <b>Og</b> Oganesson 118	140 <b>Ce</b> Cerium 58	141 <b>Pr</b> Praseodymium 59	142 <b>Nd</b> Neodymium 60	143 <b>Pm</b> Promethium 61	144 <b>Sm</b> Samarium 62	145 <b>Eu</b> Europium 63	146 <b>Gd</b> Gadolinium 64	147 <b>Tb</b> Terbium 65	148 <b>Dy</b> Dysprosium 66	149 <b>Ho</b> Holmium 67	150 <b>Er</b> Erbium 68	151 <b>Tm</b> Thulium 69	152 <b>Yb</b> Ytterbium 70	153 <b>Lu</b> Lutetium 71	154 <b>Yt</b> Ytterbium 70	155 <b>Lu</b> Lutetium 71	156 <b>Yb</b> Ytterbium 70	157 <b>Lu</b> Lutetium 71	158 <b>Yt</b> Ytterbium 70	159 <b>Lu</b> Lutetium 71	160 <b>Yb</b> Ytterbium 70	161 <b>Lu</b> Lutetium 71	162 <b>Yb</b> Ytterbium 70	163 <b>Lu</b> Lutetium 71	164 <b>Yt</b> Ytterbium 70	165 <b>Lu</b> Lutetium 71	166 <b>Yb</b> Ytterbium 70	167 <b>Lu</b> Lutetium 71	168 <b>Yt</b> Ytterbium 70	169 <b>Lu</b> Lutetium 71	170 <b>Yb</b> Ytterbium 70	171 <b>Lu</b> Lutetium 71	172 <b>Yt</b> Ytterbium 70	173 <b>Lu</b> Lutetium 71	174 <b>Yb</b> Ytterbium 70	175 <b>Lu</b> Lutetium 71

\*58-71 Lanthanoid series  
90-103 Actinoid series

**Key**  

a	<b>X</b>
b	

 a = relative atomic mass  
 X = atomic symbol  
 b = proton (atomic) number

The volume of one mole of any gas is 24 dm<sup>3</sup> at room temperature and pressure (r.t.p.).