

# UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS International General Certificate of Secondary Education

CANDIDATE NAME					
CENTRE NUMBER		CAND NUMB			

CHEMISTRY 0620/63

Paper 6 Alternative to Practical

October/November 2013

1 hour

Candidates answer on the Question Paper.

No Additional Materials are required.

#### **READ THESE INSTRUCTIONS FIRST**

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use a pencil for any diagrams, graphs or rough working.

Do not use staples, paper clips, highlighters, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

Answer all questions.

Electronic calculators may be used.


You may lose marks if you do not show your working or if you do not use appropriate units.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [ ] at the end of each question or part question.

For Examiner's Use

1 A student tried to separate a mixture of salt and sand. Salt, sodium chloride, is soluble in water. Sand, silicon(IV) oxide, is insoluble in water. He added the mixture to water. Three sets of apparatus are shown.




[Total: 7]

For Examiner's nd Use

2 Sulfur dioxide is a poisonous gas which is denser than air and soluble in water. Sulfur dioxide can be prepared by adding dilute hydrochloric acid to sodium sulfite and warming the mixture.

Study the diagram of the apparatus used.



(a) Identify and explain three mistakes in the diagram.

	1	mistake	
		reason	[2]
	2	mistake	
		reason	[2]
	3	mistake	
		reason	[2]
(b)	Sta	ate <b>one</b> precaution that should be taken when carrying out this experiment.	
,			[1]

[Total: 7]

3 A scientist measured the boiling point of water at different pressures.

For Examiner's Use

(a) Use the thermometer diagrams in the table to complete the boiling point temperatures.

pressure /atmospheres	thermometer diagram	boiling point /°C
1	105	
2	125 120 13 115	
3	135	
4	135	
5	155 150 145	
6	155 150	
7	170 1-165 160	

[3]

**(b)** Plot the points on the grid and draw a smooth line graph. 170 160 150 boiling point/°C 140 130 120 110 100 0 2 3 5 6 7 8 pressure/atmospheres [4] (c) State which point is inaccurate. Why have you chosen this point? (d) Use your graph to find the boiling point of water at a pressure of 7.5 atmospheres. Show clearly on the grid how you obtained your answer. **(e)** State a chemical test for water. [Total: 14]

For Examiner's Use **4** A student investigated the reaction between aqueous potassium manganate(VII), which is purple, and two different colourless acidic solutions, **D** and **E**.

For Examiner's Use


Three experiments were carried out.

## (a) Experiment 1

A burette was filled with the solution of potassium manganate(VII) to the  $0.0\,\text{cm}^3$  mark. Using a measuring cylinder,  $25\,\text{cm}^3$  of solution D was poured into a conical flask.

Potassium manganate(VII) solution was added to the flask until the mixture just turned permanently pink.

Use the burette diagram to record the final volume in the table and complete the table.



final reading


	burette reading
final burette reading/cm <sup>3</sup>	
initial burette reading/cm <sup>3</sup>	
difference/cm <sup>3</sup>	

[2]

### (b) Experiment 2

Experiment 1 was repeated using 25 cm<sup>3</sup> of solution **E** instead of solution **D**.

Use the burette diagrams to record the readings in the table and complete the table.



initial reading

final reading

	burette reading
final burette reading/cm <sup>3</sup>	
initial burette reading/cm <sup>3</sup>	
difference/cm <sup>3</sup>	

[2]

om	
as vn.	For Examiner's Use
 [3] vas	
[1]	
[1] on	
[1] 2. [1]	
 [2] um	

(c)	Exp	periment 3			
	obs	ueous ammonia was added to solution <b>E</b> in a test-tube. A green precipitate was served.			
	The mixture was left to stand for 5 minutes. The surface of the precipitate turned brown				
	Wh	at conclusions can you draw from these observations?			
		[3]			
(d)	(i)	What colour change was observed as potassium manganate(VII) solution was added to the flask in Experiment 1?			
		[1]			
	(ii)	Why was an indicator not added to the flask?			
		[1]			
(e)	(i)	In which experiment was the greatest volume of potassium manganate(VII) solution used?			
		[1]			
	(ii)	Compare the volumes of potassium manganate(VII) used in Experiments 1 and 2.			
		[1]			
	(iii)	Suggest an explanation for the difference in volumes.			
		[2]			
(f)		experiment 2 was repeated using 12.5 cm <sup>3</sup> of solution <b>E</b> , what volume of potassium nganate(VII) solution would be used? Explain your answer.			
		[2]			
(g)	Giv	e one advantage and one disadvantage of using a measuring cylinder for solutions <b>D E</b> .			
	adv	vantage			
	disa	advantage[2]			
		[Total: 17]			

Two liquids, **F** and **G**, were analysed. **G** was an aqueous solution of potassium chloride. The tests on the liquids and some of the observations are in the following table. Complete the observations in the table.

For Examiner's Use

tests	observations		
tests on liquid F			
(a) (i) Appearance of liquid F.	yellow solution		
The pH of the liquid was tested.	pH = 7		
(ii) An equal volume of dilute sulfuric acid was added to liquid <b>F</b> .	solution turned orange		
Excess aqueous sodium hydroxide was then added to the mixture.	solution turned from orange to yellow		
(b) Dilute sulfuric acid was added to liquid F followed by hydrogen peroxide.	rapid effervescence		
The mixture was shaken and the gas given off tested with a splint.	glowing splint relit		
tests on liquid <b>G</b>			
(c) Dilute nitric acid was added to liquid G followed by aqueous barium nitrate.	[1]		
(d) Dilute nitric acid was added to liquid G followed by aqueous silver nitrate.	[2]		
(e) What does test (a)(i) tell you about liquid F?			
	[2]		
(f) What type of reaction happened in test (a)(ii)? Explain your answer.			
type of reaction			
explanation	[2]		
(g) Identify the gas given off in test (b).			
	[1]		
	[Total: 8]		

6

## **Indigestion tablets**

For Examiner's Use

Indigestion pain is caused by too much acid in your stomach. The acid is hydrochloric acid. Indigestion tablets contain a base which neutralises the acid.

You are provided with two different brands of indigestion tablets, Painremuve and Indcure.

Plan an investigation to compare which of these brands of tablet is the most effective. You are provided with dilute hydrochloric acid and common laboratory apparatus.
[7]

[Total: 7]

# **BLANK PAGE**

# **BLANK PAGE**

12

## **BLANK PAGE**

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.