



Cambridge International Examinations
Cambridge International General Certificate of Secondary Education

CANDIDATE
NAME

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--

CHEMISTRY

0620/43

Paper 4 Theory (Extended)

October/November 2018

1 hour 15 minutes

Candidates answer on the Question Paper.

No Additional Materials are required.

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

DO **NOT** WRITE IN ANY BARCODES.

Answer **all** questions.

Electronic calculators may be used.

A copy of the Periodic Table is printed on page 12.

You may lose marks if you do not show your working or if you do not use appropriate units.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.

The syllabus is approved for use in England, Wales and Northern Ireland as a Cambridge International Level 1/Level 2 Certificate.

This document consists of **11** printed pages and **1** blank page.



- 1 Answer the following questions using only the substances in the list.
Each substance may be used once, more than once or not at all.

ammonia	bauxite	carbon dioxide	carbon monoxide
hematite	oxygen	sodium chloride	sulfur dioxide

State which substance is:

- (a) an element [1]
- (b) an ore of iron [1]
- (c) used to bleach wood pulp [1]
- (d) used to manufacture fertilisers [1]
- (e) a toxic gas produced during the incomplete combustion of hydrocarbons
..... [1]
- (f) an ionic compound [1]
- (g) a reactant in photosynthesis [1]
- (h) a product of photosynthesis. [1]

[Total: 8]

2 This question is about electrolysis.

(a) (i) What is meant by the term *electrolysis*?

.....
 [2]

(ii) Name the type of particle responsible for the conduction of electricity during electrolysis in:

the metal wires

the electrolyte [2]

(b) The table gives information about the products of the electrolysis of two electrolytes. Platinum electrodes are used in each case.

(i) Give **two** reasons why platinum is suitable to use as an electrode.

1

2 [2]

(ii) Complete the table.

electrolyte	observation at the anode (+)	name of product at the anode (+)	observation at the cathode (-)	name of product at the cathode (-)
concentrated aqueous potassium chloride			bubbles of colourless gas	
aqueous copper(II) sulfate	bubbles of colourless gas			

[6]

[Total: 12]

- 3 Tin is a metallic element in Group IV. Its main ore is cassiterite which is an impure form of tin(IV) oxide, SnO_2 .
Tin also occurs in stannite, $\text{Cu}_2\text{FeSnS}_4$.

(a) Calculate the relative formula mass, M_r , of $\text{Cu}_2\text{FeSnS}_4$.

$$M_r \text{ of } \text{Cu}_2\text{FeSnS}_4 = \dots\dots\dots [1]$$

(b) The M_r of SnO_2 is 151.

Calculate the percentage of tin by mass in SnO_2 .

$$\text{percentage of tin by mass in } \text{SnO}_2 = \dots\dots\dots [1]$$

(c) The percentage of tin by mass in $\text{Cu}_2\text{FeSnS}_4$ is 27.6%.

Use this information and your answer to (b) to suggest whether it would be better to extract tin from SnO_2 or $\text{Cu}_2\text{FeSnS}_4$.
Explain your answer.

.....
..... [1]

(d) Tin can be extracted by heating tin(IV) oxide with carbon. Carbon monoxide is the other product.

Write a chemical equation for this reaction.

..... [2]

(e) The position of tin in the reactivity series is shown.

iron	most reactive
tin	↑
copper	least reactive

A student added iron to a solution containing Sn^{2+} ions.

The student then separately added tin to a solution containing Cu^{2+} ions.

Complete the ionic equations. If there is no reaction write 'no reaction'.



[2]

(f) Copper(II) nitrate, $\text{Cu}(\text{NO}_3)_2$, decomposes when it is heated. The only solid product is copper(II) oxide, CuO . There are two gaseous products. One of the gaseous products is oxygen.

(i) Describe a test for oxygen.

test

result

[2]

(ii) Name the other gaseous product. Describe its appearance.

name

appearance

[2]

(iii) Write a chemical equation for the thermal decomposition of copper(II) nitrate.

..... [1]

(g) Iron does not rust when it is completely coated with zinc. When the zinc is scratched, the iron still does not rust.

(i) Explain why the iron does **not** rust when it is completely coated with zinc.

..... [1]

(ii) Explain why the iron still does **not** rust when the zinc is scratched.

.....

.....

.....


.....

.....

..... [3]


[Total: 16]

- 4 (a) Dilute sulfuric acid and aqueous potassium hydroxide can be used to make potassium sulfate crystals using a method that includes titration.



A student titrated 25.0 cm³ of 0.0500 mol/dm³ aqueous potassium hydroxide with dilute sulfuric acid in the presence of an indicator. The volume of dilute sulfuric acid needed to neutralise the aqueous potassium hydroxide was 20.0 cm³.

The equation for the reaction is shown.



Determine the concentration of the dilute sulfuric acid.

- Calculate the number of moles of aqueous potassium hydroxide used.

..... mol

- Calculate the number of moles of dilute sulfuric acid needed to neutralise the aqueous potassium hydroxide.

..... mol

- Calculate the concentration of the dilute sulfuric acid.

..... mol/dm³
[3]

- (b) After the titration has been completed, the conical flask contains an aqueous solution of potassium sulfate and some of the dissolved indicator.

Describe how to prepare a **pure**, dry sample of potassium sulfate crystals from new solutions of dilute sulfuric acid and aqueous potassium hydroxide of the same concentrations as used in the titration. Include a series of key steps in your answer.

.....

.....

.....

.....

.....

.....

.....

..... [5]

- (c) Potassium hydrogensulfate, KHSO_4 , is an acid salt. It dissolves in water to produce an aqueous solution, **X**, containing K^+ , H^+ and SO_4^{2-} ions.

Describe what you would see when the following experiments are done.

- (i) Magnesium ribbon is added to an excess of solution **X**.

.....

..... [2]

- (ii) A flame test is done on solution **X**.

..... [1]

- (iii) An aqueous solution containing barium ions is added to solution **X**.

..... [1]

- (d) Dilute sulfuric acid reacts with bases, metals and carbonates.

Write chemical equations for the reaction of dilute sulfuric acid with each of the following:

- (i) magnesium hydroxide

..... [2]

- (ii) zinc

..... [2]

- (iii) sodium carbonate


..... [2]

[Total: 18]

- 5 A student investigates the rate of reaction between lumps of calcium carbonate and dilute hydrochloric acid using the apparatus shown.



The calcium carbonate was in excess.



- (a) Which measurements should the student make during the reaction to determine the rate of reaction?

.....
..... [2]

- (b) What happens to the rate of reaction as the reaction proceeds? Explain your answer.

.....
.....
..... [3]

- (c) The student repeated the experiment at a higher temperature. All other conditions were kept the same. The student found that the rate of reaction increased.

Explain, in terms of collisions, why the rate of reaction increased.

.....
.....
..... [4]

- (d) Apart from using a higher temperature, suggest **two** other methods of increasing the rate of this reaction.

1

2

[2]

[Total: 11]

6 (a) Ethanol can be manufactured by fermentation and by hydration.

(i) Describe these **two** processes of ethanol manufacture.

In each case you should:

- identify the reactants
- give the reaction conditions
- write a chemical equation for the reaction which produces ethanol.

fermentation

.....

.....

.....

.....

hydration

.....

.....

.....

.....

[6]

(ii) Give **two** advantages of ethanol manufacture by fermentation compared to by hydration.

1

2

[2]


(iii) State **two** major uses of ethanol.

1

2

[2]

(b) The structures of some polymers are shown.



Answer the following questions about these polymers.
Each polymer may be used once, more than once or not at all.

State which polymer, **A**, **B**, **C**, **D** or **E**, represents:

- (i) an addition polymer [1]
- (ii) a protein [1]
- (iii) a polyester made from only **one** monomer [1]
- (iv) *Terylene* [1]
- (v) a complex carbohydrate. [1]

[Total: 15]

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cie.org.uk after the live examination series.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

The Periodic Table of Elements

		Group															
I	II	III	IV	V	VI	VII	VIII										
1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
Li lithium 7	Be beryllium 9	B boron 11	C carbon 12	N nitrogen 14	O oxygen 16	F fluorine 19	Ne neon 20										
11	12	13	14	15	16	17	18										
Na sodium 23	Mg magnesium 24	Al aluminium 27	Si silicon 28	P phosphorus 31	S sulfur 32	Cl chlorine 35.5	Ar argon 40										
19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36
K potassium 39	Ca calcium 40	Sc scandium 45	Ti titanium 48	V vanadium 51	Cr chromium 52	Mn manganese 55	Fe iron 56	Co cobalt 59	Ni nickel 59	Cu copper 64	Zn zinc 65	Ga gallium 70	Ge germanium 73	As arsenic 75	Se selenium 79	Br bromine 80	Kr krypton 84
37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54
Rb rubidium 85	Sr strontium 88	Y yttrium 89	Zr zirconium 91	Nb niobium 93	Mo molybdenum 96	Tc technetium —	Ru ruthenium 101	Rh rhodium 103	Pd palladium 106	Ag silver 108	Cd cadmium 112	In indium 115	Sn tin 119	Sb antimony 122	Te tellurium 128	I iodine 127	Xe xenon 131
55	56	57-71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86
Cs caesium 133	Ba barium 137	lanthanoids	Hf hafnium 178	Ta tantalum 181	W tungsten 184	Re rhenium 186	Os osmium 190	Ir iridium 192	Pt platinum 195	Au gold 197	Hg mercury 201	Tl thallium 204	Pb lead 207	Bi bismuth 209	Po polonium —	At astatine —	Rn radon —
87	88	89-103	104	105	106	107	108	109	110	111	112	113	114	115	116	117	118
Fr francium —	Ra radium —	actinoids	Rf rutherfordium —	Db dubnium —	Sg seaborgium —	Bh bohrium —	Hs hassium —	Mt meitnerium —	Ds darmstadtium —	Rg roentgenium —	Cn copernicium —	Nh nihonium —	Fl flerovium —	Lv livermorium —	Ts tennessine —	Og oganesson —	—

1
H
hydrogen
1

Key
atomic number
atomic symbol
name
relative atomic mass

57	58	59	60	61	62	63	64	65	66	67	68	69	70	71
La lanthanum 139	Ce cerium 140	Pr praseodymium 141	Nd neodymium 144	Pm promethium —	Sm samarium 150	Eu europium 152	Gd gadolinium 157	Tb terbium 159	Dy dysprosium 163	Ho holmium 165	Er erbium 167	Tm thulium 169	Yb ytterbium 173	Lu lutetium 175
89	90	91	92	93	94	95	96	97	98	99	100	101	102	103
Ac actinium —	Th thorium 232	Pa protactinium 231	U uranium 238	Np neptunium —	Pu plutonium —	Am americium —	Cm curium —	Bk berkelium —	Cf californium —	Es einsteinium —	Fm fermium —	Md mendelevium —	No nobelium —	Lr lawrencium —

lanthanoids

actinoids

The volume of one mole of any gas is 24 dm³ at room temperature and pressure (r.t.p.).