



Cambridge IGCSE™

CANDIDATE
NAME

CENTRE
NUMBER

| | | | | |
|--|--|--|--|--|
| | | | | |
|--|--|--|--|--|

CANDIDATE
NUMBER

| | | | |
|--|--|--|--|
| | | | |
|--|--|--|--|

* 8 9 5 5 2 3 8 2 3 1 *



CHEMISTRY

0620/62

Paper 6 Alternative to Practical

October/November 2020

1 hour

You must answer on the question paper.

No additional materials are needed.

INSTRUCTIONS


- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- You may use a calculator.
- You should show all your working and use appropriate units.

INFORMATION

- The total mark for this paper is 40.
- The number of marks for each question or part question is shown in brackets [].

This document has **12** pages. Blank pages are indicated.

- 1 The diagram shows the apparatus used to pass an electric current through concentrated hydrochloric acid. Hydrogen and chlorine were formed at the electrodes.



- (a) Name the item of apparatus labelled **A**.

..... [1]

- (b) The electrodes were made of platinum.

- (i) Give **two** reasons why platinum is a suitable material for the electrodes.

1

2

[2]

- (ii) Suggest another material suitable to use as electrodes in this experiment.

..... [1]

- (c) The teacher doing this experiment wore safety glasses, gloves, had their hair tied back and stood up throughout the experiment.

State **one** other safety precaution that should be taken when doing this experiment.
Explain your answer.

safety precaution

explanation

[2]

[Total: 6]

- 2 A student investigated the rate of a reaction between sodium metabisulfite and potassium iodate. In the reaction, starch was used as an indicator. At first the reacting mixture remained colourless but then suddenly changed to a blue-black colour.

Five experiments were done. In each experiment the total volume of liquid was 45 cm³.

Experiment 1

- Using a 10 cm³ measuring cylinder, 5 cm³ of aqueous sodium metabisulfite was poured into a beaker.
- Using another 10 cm³ measuring cylinder, 5 cm³ of aqueous starch was poured into the beaker.
- Using a 25 cm³ measuring cylinder, 15 cm³ of distilled water was poured into the beaker.
- Using another 25 cm³ measuring cylinder, 20 cm³ of aqueous potassium iodate was poured into the beaker. At the same time a stop-clock was started.
- The mixture in the beaker was stirred until a sudden colour change was seen.
- The stop-clock was immediately stopped and the time recorded.
- The beaker was rinsed with water.

Experiment 2

- Experiment 1 was repeated using 17 cm³ of distilled water and 18 cm³ of aqueous potassium iodate.

Experiment 3

- Experiment 1 was repeated using 21 cm³ of distilled water and 14 cm³ of aqueous potassium iodate.

Experiment 4

- Experiment 1 was repeated using 23 cm³ of distilled water and 12 cm³ of aqueous potassium iodate.

Experiment 5


- Experiment 1 was repeated using 25 cm³ of distilled water and 10 cm³ of aqueous potassium iodate.

(a) Use the information in the description of the experiments and the stop-clock diagrams to complete the table. Record the times in **seconds**.

| experiment | volume of aqueous sodium metabisulfite /cm ³ | volume of distilled water /cm ³ | volume of aqueous potassium iodate /cm ³ | stop-clock diagram | time to change colour /s |
|------------|---|--|---|--------------------|--------------------------|
| 1 | | | 20 | | |
| 2 | | | 18 | | |
| 3 | | | 14 | | |
| 4 | | | 12 | | |
| 5 | | | 10 | | |

[5]

- (b) Plot the results from Experiments 1 to 5 on the grid.
Draw a smooth curve of best fit.



[3]

- (c) (i) **From your graph**, predict the time to change colour if 16 cm³ of aqueous potassium iodate was used.
Show clearly **on the grid** how you worked out your answer.

time to change colour = s [2]

- (ii) Calculate the volume of distilled water required if 16 cm³ of aqueous potassium iodate was used.

volume of distilled water = cm³ [1]

- (d) Sketch **on the grid** the graph you would expect if Experiments 1 to 5 were repeated at a higher temperature. [1]

- (e) The concentration of potassium iodate in the reaction mixture in each experiment can be calculated using the equation shown.

$$\text{concentration} = \frac{0.05 \times \text{volume of aqueous potassium iodate}}{45}$$

- (i) Calculate the concentration of potassium iodate in the reaction mixture in Experiment 2.

concentration = mol/dm³ [1]

- (ii) State which experiment, 1, 2, 3, 4 or 5, had the fastest rate of reaction.

..... [1]

- (f) Suggest why the volume of distilled water added to each experiment was increased as the volume of aqueous potassium iodate was decreased.

.....
 [1]

- (g) Give **one** change you could make to the apparatus used which would improve the results. Explain your answer.

change to apparatus

.....

explanation

..... [2]

- (h) How could the reliability of the results of this investigation be checked?

.....
 [1]

[Total: 18]

- 3 Solid **Q** and solid **R** were analysed. Solid **Q** was zinc carbonate.
Tests were done on each solid.

tests on solid Q

Complete the expected observations.

- (a) Solid **Q** was placed in a boiling tube. About 10 cm³ of dilute sulfuric acid was added to the boiling tube. Any gas produced was tested.
The contents of the boiling tube were kept for (c).

observations

.....

..... [3]

- (b) Identify the gas given off in (a).

..... [1]

- (c) The reaction mixture from (a) was filtered.
The filtrate was solution **S**. 1 cm depth of solution **S** was poured into a boiling tube.

- (i) Aqueous sodium hydroxide was added dropwise and then in excess to solution **S** in the boiling tube.

observations

.....

..... [2]

- (ii) Explain why it is **not** possible to identify the cation contained in solution **S** from your observations in (c)(i).

.....

..... [1]

- (iii) Suggest an additional test that can be done on solution **S** to confirm the cation was Zn²⁺.

.....

..... [1]

tests on solid R

Tests were done and the following observations were made.

| tests on solid R | observations |
|--|--|
| test 1 A flame test was done on solid R. | yellow flame |
| Solid R was dissolved in distilled water to produce solution R. The solution was divided into two equal portions in two test-tubes. test 2 About 1 cm ³ of dilute nitric acid followed by a few drops of aqueous silver nitrate were added to the first portion of solution R. | yellow precipitate formed |
| test 3 The second portion of solution R was added to 1 cm ³ of aqueous bromine in a test-tube. | the solution changed colour from orange to brown |

(d) Identify solid R.

.....
..... [2]

[Total: 10]

4 Brass is a mixture of two metals, copper and zinc.

Copper does not react with dilute sulfuric acid. Zinc reacts with hot dilute sulfuric acid to form the soluble salt zinc sulfate.

Plan an investigation to find the percentage by mass of zinc in a sample of brass.
In your answer you should include how to calculate the percentage by mass of zinc.

You have access to normal laboratory apparatus.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

..... [6]

BLANK PAGE

BLANK PAGE

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.