UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS GCE Ordinary Level

MARK SCHEME for the May/June 2008 question paper

5070 CHEMISTRY

5070/02

Paper 2 (Theory), maximum raw mark 75

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began.

All Examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes must be read in conjunction with the question papers and the report on the examination.

• CIE will not enter into discussions or correspondence in connection with these mark schemes.

CIE is publishing the mark schemes for the May/June 2008 question papers for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level syllabuses and some Ordinary Level syllabuses.



www.dynamicpapers.com Syllabus

5070

Paper

02

		GCL O LLVLL – May/June 2000	3070 02	
A 1	(a)	carbon monoxide / CO		[1]
	(b)	ammonia / NH ₃		[1]
	(c)	argon / Ar		[1]
	(d)	carbon monoxide / CO		[1]
	(e)	oxygen / O ₂ NOT: O		[1]
			[Tota	al: 5]
A2	(a)	$36.8(\%)$ / 36.8 / $37(\%)$ (answer alone = 2 marks) (NOT 36%) $M_{\rm r}$ of iron(II) sulphate = 152 (for 1 mark)		[2]
	(b)	barium nitrate / other soluble barium salt e.g. barium chloride NOT: barium hydroxide white precipitate / solid IGNORE: incorrect name of precipitate ALLOW: this mark if nitric acid missing from 1 st marking point	·	[1] [1]
	(c)	$4Fe^{2+} + O_2 + 4H^+ \rightarrow 4Fe^{3+} + 2H_2O$ 1 mark for correct reactants and products; 1 mark for correct balance		[2]
	(d)	(i) orange to green		[1]
		(ii) green to yellow ALLOW: brown / orange / reddish brown		[1]
	(e)	(i) 0.00076 / 7.6 × 10 ⁻⁴ (moles)		[1]
		(ii) mols $Fe^{2+} = 0.00456$ ALLOW: 0.0046		[1]
		mass of iron(II) ions = $0.255 / 0.26 / 0.258$ (g)	v FC1	[1]

Mark Scheme

GCE O LEVEL - May/June 2008

Page 2

[Total: 11]

ALLOW: error carried forward [i.e. answer to moles Fe²⁺ × 56]

WWW.	.dynamicpap	ers.com
	Syllahus	Pane

Pa	Page 3 Mark Scheme Syllabus			
		GCE O LEVEL – May/June 2008	5070	Paper 02
.3 (a)	43 proto 55 neutro	ns + 43 electrons ons		[1 [1
(b)		onable, correct, isotope e.g. $^{97}_{43}$ Tc mass numbers from between 86 and 110		[1
(c)	ALLOW: electrons NOT: ch	mber of electrons and protons / same number of + and balance between the number of protons and electrons are - and protons are + arge on electron = to that on the proton arge on electron and proton is opposite		[1
(d)	variaformsolhigh(con	melting point / boiling point; able valency / oxidation state / (compounds) have ions a coloured <u>compounds</u> / form coloured <u>ions;</u> [NOT: if ution] density; appounds) form complex ions		orms coloure
	• cata	lytic activity		[2
				[Total: 7
4 (a)		alkane: (bromine) stays orange / no (colour) change / bromine colours of brown / red / orange	stays the same;	[
		alkene: (bromine) decolourised / (orange) to colourles	s	[
(b)	6 correct	ectrons between the two carbons; shared pairs between carbons and 6 hydrogen atoms dent marking points]	i	[
(c)	ALLOW:	$C_2H_4Cl_2$ etc. (up to C_2Cl_6) any order of atoms		[

[Total: 7]

[1]

[1]

ALLOW: correct graphical / displayed formulae / dot and cross diagrams

ALLOW: but-1-ene / but-2-ene / methylpropene

NOT: CH₃CH₂CH₂CH₃ / graphical formulae

ALLOW: HC1

(d) butene / butylene

NOT: C_nH_{2n}

WWW.	dynamicpap	ers.com
	Syllabus	Pape

Paper

P	age 4		Mark Scheme	Syllabus	Paper
			GCE O LEVEL – May/June 2008	5070	02
.5 (a)	(i)	P ₂ O ₅	₅ / P ₄ O ₁₀		[1
	(ii)	low in ALLO cher acidi	sical property: melting point / low boiling point / electrical insulator or o OW: white in colour / solid mical property: ic oxide / reacts with alkalis / reacts with bases / dissol : it is an acid / dissolves in water		[1
(b)	1 m	ark fo	→2KC <i>l</i> + 3O ₂ or correct reactant and products; or correct balance		[2
(c)			→ SO ₂ :: state symbols		[1
(d)			nd C _n H _{2n+2} its a general formula		[′
					[Total: 7
۱6 (a)	(i)	ALL IGN	anoes / treatment of <u>sulphide</u> ores OW: bacterial <u>oxidation</u> / <u>burning</u> natural gas ORE: unqualified burning fuels / from car engines / ke / from power stations	making sulphur	[′ ic acid / fror
	(ii)	ALL NOT	ning / car <u>engines</u> / car exhausts / <u>high temperature</u> fur OW: burning fuel in car T: from cars unqualified T: bacterial activity / from fertilizers	rnaces / explosiv	es [ˈ
(b)	(i)	carb	on dioxide / CO ₂		[
	(ii)		ium nitrite / calcium nitrate or correct formulae ORE: incorrect oxidation numbers		[
	(iii)	ALLO NOT • ALLO NOT	one of: erodes buildings / reacts with buildings or statues OW: corrodes buildings / eats away buildings : destroys buildings / damages buildings forest death / kills trees or plants / kills fish in lakes / OW: damages / destroys crops : kills animals (unless in lakes / rivers) breathing difficulties in humans OWTTE : causes pollution / harmful (unless specified) / affects		[′ als

Mark Scheme

Page 4

www.dynamicpapers.com **Syllabus**

5070

Paper

02

			GCE O LEVEL – Way/June 2000	5070	02
	(c)	correct :correct :ALLOW: lineALLOW: E for a correct :	t on left and product on right and products above rearrow and label for activation energy (even if exoth arrow and label for enthalpy change in place of arrow for activation energy and 43 kJ for ΔH rection of arrow		[1] rawn) [1] [1]
					[Total: 8]
В7	(a)	NOT: goes y			[1]
		ALLOW: oxi	→ Br ₂ + 2Cl ⁻ gained electrons gained electrons gained electrons dation number of chlorine decreases / goes from ect oxidation numbers de has gained electrons	0 to -1	[1] [1]
	(b)	with 2+ at to NOT: 2+ in r	es diagram of magnesium ion (ignore whether dots op right / near top right nucleus tten as Mg ²⁺ = 2.8	or crosses)	[1]
		dot and cros with - at top ALLOW: onl	as diagram of chloride ion (ignore whether dots or conjugated right / near top right by one chloride ion shown then as $Cl^- = 2.8.8$	crosses)	[1]
	(c)		e it / silver nitrate in water; e / add aqueous solution / from (aq) in equation		[1]
		add sole water / I ALLOW: hyd	ution of soluble chloride / named soluble chloride hydrochloric acid; drochloric acid alone without the word solution or d	lissolved in wate	[1]
		filter;	s mark if equation given with ALL state symbols co	rrect	[1]
			recipitate with water <u>and</u> leave water to evapora	ate / wash ppt v	vith water <u>and</u>
		ALLOW was	sh ppt with water <u>and</u> dry in an oven		[1]

Mark Scheme

GCE O LEVEL - May/June 2008

Page 5

[Total: 10]

[1]

ALLOW: thins ozone layer / damages ozone layer / makes hole in ozone layer

(d) depletion of ozone / destroys ozone (molecules)

NOT: increases risk / causes skin cancer

ALLOW: increases greenhouse effect / greenhouse gas

www.dynamicpapers.com

Paper

Syllabus

				GCE O LEVEL – May/June 2008	5070	02
В8	(a)			oint / volatility :: number of carbon atoms		[1]
	(b)	(i)	ALL ALL NOT	akdown of long chained hydrocarbons (into shorter / sn OW: large for long chained; alkanes / carbon chains fo OW: converting long chained alkanes to alkenes F: splitting larger fractions F: breaking down larger substances / molecules / partic	r hydrocarbons	[1]
			by <u>h</u> or b NOT	igh temperature / stated temperatures in range 400–80 y high temperature and catalyst / stated temperatures T: by heating / heat OW: aluminium oxide / silicon dioxide / zeolites in place	00°C; in range 200–80	•
		(ii)	grea	tions which are less needed / exceed demand chang ter demand; OW: idea of less useful fractions used to make more u		re needed / in [1]
				F: larger fractions / alkanes to smaller alkanes oil fraction converted to gasoline		[1]
			ALL	Oll fraction converted to gasoline OW: gas oil fraction converted to kerosene / petroleum OW: waxes converted to one of the above 3 fra	•	[1]
				verted to one of the above 3 fractions	ictions / waxes	and bitumen
	(c)	(i)	CH ₃	CH=CH ₂ (minimum structure to show double bond)		[1]
		(ii)	ALL	$H_{32} \rightarrow C_3H_6 + C_{12}H_{26}$ OW: other possible product apart from propene with country $C_3H_6 + C_9H_{20}$ on right	orrect balance	[1]
	(d)	(i)	ALL ALL	et with <u>steam</u> and <u>catalyst</u> (both required) OW: phosphoric acid (in place of the word 'catalyst') OW: water + temperature of above 100°C in place of some correct equation with correct state symbols	team	[1]
			CH ₃	Ր։ fermentation CH₂CH₂OH / CH₃CH(OH)CH₃ (as minimum) OW: full formula showing all atoms and bonds or mixtւ	ures of the two	[1]
		(ii)		$H(CH_3) - CH_2 - CH(CH_3) - CH_2 - $ or full structural for $OW: - [CH(CH_3) - CH_2]_n - $	mula	[1]
						[Total: 10]
В9	(a)		/ H₃O T: 'hy	+ vdrogen ions'		[1]
	(b)	(i)	Mg	es Mg $(0.24/24) = 0.01$ AND moles acid $(2 \times 5/100)$ in excess since requires 2 moles acid to 1 mole mag in equation	,	[1] se of 1:2 mole [1]
		(ii)	0.00	es MgC <i>l</i> ₂ (0.01/2) = 0.005; 15 × 95 = 4.75 / 0.48 g [NOT: 0.4 (g)] OW: error carried forward from directly above and fron	n part (i)	[1] [1]

Mark Scheme

Page 6

ALLOW: error carried forward from directly above and from part (i)

www.dynamicpapers.com
Syllabus Paper

[1]

		,		Cullabus	
Page 7			Mark Scheme GCE O LEVEL – May/June 2008	Syllabus 5070	Paper 02
	(iii)	ALL	'3 of: same number of moles of each acid / same amount of acid / same number of hydrogen ions which react in each of each acid at the same volumber of each acid and ethanoic acid acid is stronger than ethanoic acid ORA; hydrochloric acid fully ionised and ethanoic acid partial OW: hydrochloric acid more ionised than ethanoic acid higher concentration of hydrogen ions in hydrochloric hydrogen ions in ethanoic acid; more collisions per unit time / collision rate higher ethanoic acid ORA	ach acid; ume is a weak acid ally ionised d ORA c acid / lower co	/ hydrochlorion
(c)	(i) (ii)	bubl ALL ALL	I ₃ COOH + Na ₂ CO ₃ → 2CH ₃ COONa + CO ₂ + H ₂ O OW: correct ionic form for sodium ethanoate oles/ effervescence OW: tube gets hot / heat given off OW: sodium carbonate dissolves / disappears c: gas given off / carbon dioxide given off		[1
310 (a)	ALI neg IGN NO NO	LOW: gative NORE T: ele	pattern of positive ions; + / X ⁺ / X ²⁺ etc. for the positive ions sign / e ⁻ / e dispersed amongst the ions inequality of numbers of electrons and + charges ectrons in clumps separated from positive ions egative sign / e ⁻ / e in circles unless the circles are ions	considerably sm	[Total: 10] [1] [1] aller than the
(b)	NO NO	T: ele T: ref	s move / electrons are delocalised / sea of electrons ectrons are free (unless qualified) ference to free electrons in the outer shells / valency associated with particular atoms	electrons if it im	[1] plies that they
(c)		ALL NOT	tion is fast <u>er</u> OW: larg <u>er</u> surface area for reaction : reaction is fast (comparison needed)		[1
	(ii)	mas	es hydrogen (0.072 / 24) = 0.003 s zinc = 0.003 × 65 = 0.195 g OW: error carried forward		[1
	/···	400	50/ / 40 00/		F.4

ALLOW: error carried forward from part (ii) to give values below 100%

(iii) 16.25% / 16.3%

www.dynamicpapers.com

Page 8	Mark Scheme	Syllabus	Paper
	GCE O LEVEL – May/June 2008	5070	02

(d) three of: [3]

- (zinc gives) white precipitate (on addition of aqueous ammonia);
- (white) ppt dissolves in excess ammonia/gives colourless solution with excess ammonia;
- copper would give (light) blue ppt (on addition of aqueous ammonia);

ALLOW: ppt is not blue

 (if copper) (light) blue ppt would dissolve in excess ammonia/gives blue solution with excess ammonia;

ALLOW: no blue solution formed with excess ammonia

[Total: 10]