

Cambridge Assessment International Education Cambridge Ordinary Level

CHEMISTRY 5070/31

Paper 3 Practical Test MARK SCHEME

Maximum Mark: 40

May/June 2019

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the May/June 2019 series for most Cambridge IGCSE™, Cambridge International A and AS Level and Cambridge Pre-U components, and some Cambridge O Level components.

Cambridge O Level – Mark Scheme PUBLISHED

Generic Marking Principles

These general marking principles must be applied by all examiners when marking candidate answers. They should be applied alongside the specific content of the mark scheme or generic level descriptors for a question. Each question paper and mark scheme will also comply with these marking principles.

GENERIC MARKING PRINCIPLE 1:

Marks must be awarded in line with:

- the specific content of the mark scheme or the generic level descriptors for the question
- the specific skills defined in the mark scheme or in the generic level descriptors for the question
- the standard of response required by a candidate as exemplified by the standardisation scripts.

GENERIC MARKING PRINCIPLE 2:

Marks awarded are always whole marks (not half marks, or other fractions).

GENERIC MARKING PRINCIPLE 3:

Marks must be awarded **positively**:

- marks are awarded for correct/valid answers, as defined in the mark scheme. However, credit is given for valid answers which go beyond the scope of the syllabus and mark scheme, referring to your Team Leader as appropriate
- marks are awarded when candidates clearly demonstrate what they know and can do
- · marks are not deducted for errors
- marks are not deducted for omissions
- answers should only be judged on the quality of spelling, punctuation and grammar when these features are specifically assessed by the question as indicated by the mark scheme. The meaning, however, should be unambiguous.

GENERIC MARKING PRINCIPLE 4:

Rules must be applied consistently e.g. in situations where candidates have not followed instructions or in the application of generic level descriptors.

© UCLES 2019 Page 2 of 7

GENERIC MARKING PRINCIPLE 5:

Marks should be awarded using the full range of marks defined in the mark scheme for the question (however; the use of the full mark range may be limited according to the quality of the candidate responses seen).

GENERIC MARKING PRINCIPLE 6:

Marks awarded are based solely on the requirements as defined in the mark scheme. Marks should not be awarded with grade thresholds or grade descriptors in mind.

© UCLES 2019 Page 3 of 7

Question	Answer	Marks
1(a)	Titration Measurements (1) Both readings i.e. initial and final are present for each titration and the readings are recorded to 1dp.	12
	Titres (1) All the titres are calculated correctly i.e. no subtraction errors and all recorded in the correct row i.e. volume of Q used	
	Accuracy (6) For the two best titres give: 3 marks for a titre within 0.2 cm³ of the Supervisor's value. 2 marks for a titre within 0.3 cm³ of the Supervisor's value. 1 mark for a titre within 0.4 cm³ of the Supervisor's value. No marks for a titre more than 0.4 cm³ from the Supervisor's value. Concordance (3) Give 3 marks if all the ticked values are within 0.2 cm³. Give 2 marks if all the ticked values are within 0.3 cm³. Give 1 marks if all the ticked values are within 0.4 cm³.	
	Average (1) Give 1 mark if the candidate calculates a correct average of selected titres.	
1(b)	Pipette volume = 25 cm^3 and the average volume of Q used = 20.2 cm^3 Moles of potassium manganate(VII) in the average volume = $(20.2 \times 0.0192) / 1000$ = 0.000388	1
1(c)	Answer from (b) \times 5/2 = 0.000388 \times 5/2 = 0.000970	1
1(d)	Answer from (c) × 1000 / 25 (or 40) = 0.000970 × 1000 / 25 = 0.0388	1

© UCLES 2019 Page 4 of 7

Question	Answer	Marks
1(e)	Answer from (d) × 126 = 0.0388 × 126 = 4.88 g	1
1(f)	The volume of acid added simply needs to be enough to ensure the solution is acidic/acidified. OWTTE.	1

© UCLES 2019 Page 5 of 7

Question	Answer	Marks
For ppt: acce Ignore cloudy Ignore solution For gases: to For the evolutions: co		
2	R (test 1) (a) white ppt (1) (b) soluble in excess (1) colourless solution (1)	21
	R (test 2) (a) white ppt (1) (b) soluble in excess (1) colourless solution (1) (c) no reaction (1)	
	R (test 3) (a) no reaction (1) (b) white ppt (1) (c) no reaction (1)	
	S (test 1) (a) off-white / cream / yellow/brown ppt (1) (b) insoluble in excess (1)	

© UCLES 2019 Page 6 of 7

Question	Answer	Marks
2	S (test 2) (a) off-white / cream / yellow / brown ppt (1) (b) insoluble in excess (1) (c) bubbles (1) gas relights a glowing splint (1) oxygen (1) (darker) brown/black solid (1)	
	S (test 3) (a) no reaction (1) (b) no reaction (1) (c) white ppt (1)	
	Conclusions R is zinc sulfate / ZnSO ₄ (1) The anion in S is chloride/C l ⁻ (1)	2

© UCLES 2019 Page 7 of 7