

BLANK PAGE



Question 2 continued

Leave
blank

Lined writing area for the answer to Question 2.

(Total 6 marks)

Q2



Leave blank

3. A cricket ball of mass 0.5 kg is struck by a bat. Immediately before being struck, the velocity of the ball is $(-30\mathbf{i}) \text{ m s}^{-1}$. Immediately after being struck, the velocity of the ball is $(16\mathbf{i} + 20\mathbf{j}) \text{ m s}^{-1}$.

- (a) Find the magnitude of the impulse exerted on the ball by the bat. **(4)**

In the subsequent motion, the position vector of the ball is \mathbf{r} metres at time t seconds. In a model of the situation, it is assumed that $\mathbf{r} = [16t\mathbf{i} + (20t - 5t^2)\mathbf{j}]$. Using this model,

- (b) find the speed of the ball when $t = 3$. **(4)**



Question 3 continued

[Handwriting area with 32 horizontal lines]

Leave
blank

Q3

(Total 8 marks)



Leave blank

4.

Figure 1




Figure 1 shows four uniform rods joined to form a rigid rectangular framework $ABCD$, where $AB = CD = 2a$, and $BC = AD = 3a$. Each rod has mass m . Particles, of mass $6m$ and $2m$, are attached to the framework at points C and D respectively.

(a) Find the distance of the centre of mass of the loaded framework from

(i) AB ,

(ii) AD .

(7)

The loaded framework is freely suspended from B and hangs in equilibrium.

(b) Find the angle which BC makes with the vertical.

(3)



Question 5 continued

Leave
blank

Lined area for writing the answer to Question 5.

(Total 8 marks)

Q5




N 2 2 3 3 2 A 0 1 1 2 0

Leave blank

6.

Figure 2



A wooden plank AB has mass $4m$ and length $4a$. The end A of the plank lies on rough horizontal ground. A small stone of mass m is attached to the plank at B . The plank is resting on a small smooth horizontal peg C , where $BC = a$, as shown in Figure 2. The plank is in equilibrium making an angle α with the horizontal, where $\tan \alpha = \frac{3}{4}$. The coefficient of friction between the plank and the ground is μ . The plank is modelled as a uniform rod lying in a vertical plane perpendicular to the peg, and the stone as a particle. Show that

(a) the reaction of the peg on the plank has magnitude $\frac{16}{5} mg$, (3)

(b) $\mu \geq \frac{48}{61}$. (6)

(c) State how you have used the information that the peg is smooth. (1)



Question 6 continued

Lined writing area for Question 6 continued.

Leave blank

(Total 10 marks)

Q6



Question 7 continued

Leave
blank

Handwritten area with horizontal lines for writing.

(Total 12 marks)

Q7



Leave blank

8. Two particles A and B move on a smooth horizontal table. The mass of A is m , and the mass of B is $4m$. Initially A is moving with speed u when it collides directly with B , which is at rest on the table. As a result of the collision, the direction of motion of A is reversed. The coefficient of restitution between the particles is e .

(a) Find expressions for the speed of A and the speed of B immediately after the collision. (7)

In the subsequent motion, B strikes a smooth vertical wall and rebounds. The wall is perpendicular to the direction of motion of B . The coefficient of restitution between B and the wall is $\frac{4}{5}$. Given that there is a second collision between A and B ,

(b) show that $\frac{1}{4} < e < \frac{9}{16}$. (5)

Given that $e = \frac{1}{2}$,

(c) find the total kinetic energy lost in the first collision between A and B . (3)



BLANK PAGE



BLANK PAGE

