



Cambridge International AS & A Level

CANDIDATE
NAME

--

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--

CHEMISTRY

9701/32

Paper 3 Advanced Practical Skills 2

May/June 2023

2 hours

You must answer on the question paper.

You will need: The materials and apparatus listed in the confidential instructions

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- You may use a calculator.
- You should show all your working and use appropriate units.

INFORMATION

- The total mark for this paper is 40.
- The number of marks for each question or part question is shown in brackets [].
- The Periodic Table is printed in the question paper.
- Important values, constants and standards are printed in the question paper.
- Notes for use in qualitative analysis are provided in the question paper.

Session

Laboratory

For Examiner's Use

1

2

3

Total

This document has **12** pages.

Quantitative Analysis

Read through the whole method before starting any practical work. Where appropriate, prepare a table for your results in the space provided.

Show the precision of the apparatus you used in the data you record.

Show your working and appropriate significant figures in the final answer to **each** step of your calculations.

- 1 In this experiment you will determine the relative atomic mass, A_r , of metal **M** by thermal decomposition of its basic carbonate, $\text{MCO}_3 \cdot \text{M(OH)}_2$.

FB 1 is the basic metal carbonate, $\text{MCO}_3 \cdot \text{M(OH)}_2$.

(a) Method

- Weigh the empty crucible with its lid. Record the mass.
- Transfer all of the **FB 1** from the container into the crucible.
- Weigh the crucible, lid and **FB 1**. Record the mass.
- Calculate and record the mass of **FB 1** used.
- Place the crucible and contents on a pipe-clay triangle.
- Heat the crucible gently, with the lid on, for approximately 1 minute.
- Heat strongly, with the lid off, for a further 4 minutes.
- Replace the lid and leave the crucible to cool for at least 5 minutes.

During the cooling period, you may wish to begin work on Question 3.

- When the crucible is cool, weigh the crucible with its lid and contents. Record the mass.
- Place the crucible and contents on the pipe-clay triangle. Remove the lid.
- Heat the crucible strongly for a further 2 minutes.
- Replace the lid and leave the crucible to cool for at least 5 minutes.
- When the crucible is cool, reweigh the crucible with its lid and contents. Record the mass.
- Calculate and record the mass of residue obtained.

Results

I	
II	
III	
IV	
V	

[5]

(b) Calculations

- (i) When **FB 1** undergoes thermal decomposition, the products are the metal oxide, **MO**, carbon dioxide and water vapour.
Give the equation for the thermal decomposition of **FB 1**. Include state symbols.

..... [1]

- (ii) The amount, in mol, of carbon dioxide produced is given by the following formula.

$$\text{amount of CO}_2 = \frac{\text{mass loss during heating}}{(M_r \text{ of CO}_2 + M_r \text{ of water})}$$

Calculate the amount, in mol, of carbon dioxide produced in (a).

amount of CO₂ = mol [1]

- (iii) Calculate the relative formula mass, M_r , of the basic metal carbonate.
Show your working.

M_r of $\text{MCO}_3 \cdot \text{M}(\text{OH})_2$ = [1]

- (iv) Calculate the relative atomic mass of metal **M**.

A_r of **M** = [1]

- (c) A student accidentally spilt a little of the residue before carrying out the final weighing.
Predict whether the calculated value of the relative atomic mass of **M** will be higher or lower
as a result of this mistake.
Explain your answer.

The A_r of **M** will be

explanation

.....

..... [1]

- (d) A student suggested that addition of sulfuric acid to the residue from (a) would show whether
the basic metal carbonate had decomposed fully.
State whether the student is correct.
Explain your answer.

.....

.....

..... [1]

[Total: 11]

- 2 In this experiment you will determine the relative atomic mass, A_r , of another metal, **X**, by a titration method using the metal carbonate, X_2CO_3 .

FB 2 is $0.0460 \text{ mol dm}^{-3}$ hydrochloric acid, HCl .

FB 3 is the metal carbonate, X_2CO_3 .

FB 4 is methyl orange indicator.

(a) **Method**

Preparing a solution of FB 3

- Weigh the stoppered container of **FB 3**. Record the mass in the space below.
- Tip all of the **FB 3** into the 250 cm^3 beaker.
- Reweigh the container with its stopper. Record the mass.
- Calculate and record the mass of **FB 3** used.
- Add approximately 100 cm^3 of distilled water to **FB 3** in the beaker.
- Stir the mixture with a glass rod until all the **FB 3** has dissolved.
- Transfer this solution into the 250 cm^3 volumetric flask.
- Wash the beaker with distilled water and transfer the washings to the volumetric flask.
- Rinse the glass rod with distilled water and transfer the washings to the volumetric flask.
- Make up the solution in the volumetric flask to the mark using distilled water.
- Shake the flask thoroughly.
- This solution of X_2CO_3 is **FB 5**. Label the flask **FB 5**.

Titration

- Fill the burette with **FB 2**.
- Pipette 25.0 cm^3 of **FB 5** into a conical flask.
- Add several drops of **FB 4** to the conical flask.
- Perform a rough titration and record your burette readings in the space below.

The rough titre is cm^3 .

- Carry out as many accurate titrations as you think necessary to obtain consistent results.
- Make sure any recorded results show the precision of your practical work.
- Record, in a suitable form below, all of your burette readings and the volume of **FB 2** added in each accurate titration.

I	
II	
III	
IV	
V	
VI	
VII	
VIII	

[8]

- (b) From your accurate titration results, calculate a suitable mean value to use in your calculations. Show clearly how you obtain the mean value.

25.0 cm³ of **FB 5** required cm³ of **FB 2**. [1]

(c) Calculations

- (i) Give your answers to **(c)(ii)**, **(c)(iv)**, **(c)(v)** and **(c)(vi)** to an appropriate number of significant figures. [1]
- (ii) Calculate the amount, in mol, of hydrochloric acid present in the volume of **FB 2** in **(b)**.

amount of HCl = mol [1]

- (iii) Give the ionic equation for the reaction of hydrochloric acid with the metal carbonate during the titration. Include state symbols.

.....CO₃²⁻..... + → + [1]

- (iv) Calculate the concentration of **X₂CO₃**, in mol dm⁻³, in **FB 5**.

concentration of **X₂CO₃** in **FB 5** = mol dm⁻³ [1]

(v) Calculate the relative formula mass, M_r , of X_2CO_3 .

M_r of X_2CO_3 = [1]

(vi) Calculate the relative atomic mass of **X**.

A_r of **X** = [1]

(vii) Identify **X**.

X is [1]

[Total: 16]

Qualitative Analysis

For each test you should record all your observations in the spaces provided.

Examples of observations include:

- colour changes seen
- the formation of any precipitate and its solubility (where appropriate) in an excess of the reagent added
- the formation of any gas and its identification (where appropriate) by a suitable test.

You should record clearly at what stage in a test an observation is made.

Where no change is observed you should write 'no change'.

Where reagents are selected for use in a test, the name or correct formula of the element or compound must be given.

If any solution is warmed, a boiling tube must be used.

Rinse and reuse test-tubes and boiling tubes where possible.

No additional tests should be attempted.

- 3 (a) FB 6** is a solution containing one cation listed in the Qualitative analysis notes.
The anion contains sulfur.

- (i) State the reagents you would use to identify the cation in **FB 6**.

reagents

Use your selected reagents to test **FB 6**.

Use 1 cm depth of **FB 6** in a test-tube for each test.

Record your observations in the space below.

[2]

- (ii) Identify the anion in **FB 6**.
Include a description of your procedure and the observations you make.

anion in **FB 6** [2]

- (iii) Deduce the formula of **FB 6**.

formula of **FB 6** [1]

- (b) You will devise chemical tests to distinguish between the two possible identities given for each of compounds **FB 7**, **FB 8**, **FB 9** and **FB 10**.

In each case, you should:

- name the reagent or reagents you will use to identify the compound
- state any necessary conditions for your test
- use a 1 cm depth of the solution of the unknown compound and use a boiling tube if you need to warm a mixture
- carry out your test and record the observations you make (if any)
- state your conclusion about the identity of the compound.

- (i) **FB 7** is either aqueous sodium nitrate or aqueous sodium nitrite.

FB 7 is [2]

- (ii) **FB 8** is either aqueous sodium nitrate or aqueous silver nitrate.

FB 8 is [2]

- (iii) **FB 9** is either aqueous ethanol or aqueous propan-1-ol.
(In your test, do **not** heat but you may need to leave your reaction mixture to stand.)

FB 9 is [2]

- (iv) **FB 10** is either aqueous methanol or aqueous ethanoic acid.

FB 10 is [2]

[Total: 13]

Qualitative analysis notes

1 Reactions of cations

cation	reaction with	
	NaOH(aq)	NH ₃ (aq)
aluminium, Al ³⁺ (aq)	white ppt. soluble in excess	white ppt. insoluble in excess
ammonium, NH ₄ ⁺ (aq)	no ppt. ammonia produced on warming	–
barium, Ba ²⁺ (aq)	faint white ppt. is observed unless [Ba ²⁺ (aq)] is very low	no ppt.
calcium, Ca ²⁺ (aq)	white ppt. unless [Ca ²⁺ (aq)] is very low	no ppt.
chromium(III), Cr ³⁺ (aq)	grey-green ppt. soluble in excess giving dark green solution	grey-green ppt. insoluble in excess
copper(II), Cu ²⁺ (aq)	pale blue ppt. insoluble in excess	pale blue ppt. soluble in excess giving dark blue solution
iron(II), Fe ²⁺ (aq)	green ppt. turning brown on contact with air insoluble in excess	green ppt. turning brown on contact with air insoluble in excess
iron(III), Fe ³⁺ (aq)	red-brown ppt. insoluble in excess	red-brown ppt. insoluble in excess
magnesium, Mg ²⁺ (aq)	white ppt. insoluble in excess	white ppt. insoluble in excess
manganese(II), Mn ²⁺ (aq)	off-white ppt. rapidly turning brown on contact with air insoluble in excess	off-white ppt. rapidly turning brown on contact with air insoluble in excess
zinc, Zn ²⁺ (aq)	white ppt. soluble in excess	white ppt. soluble in excess

2 Reactions of anions

anion	reaction
carbonate, CO ₃ ²⁻	CO ₂ liberated by dilute acids
chloride, Cl ⁻ (aq)	gives white ppt. with Ag ⁺ (aq) (soluble in NH ₃ (aq))
bromide, Br ⁻ (aq)	gives cream/off-white ppt. with Ag ⁺ (aq) (partially soluble in NH ₃ (aq))
iodide, I ⁻ (aq)	gives pale yellow ppt. with Ag ⁺ (aq) (insoluble in NH ₃ (aq))
nitrate, NO ₃ ⁻ (aq)	NH ₃ liberated on heating with OH ⁻ (aq) and Al foil
nitrite, NO ₂ ⁻ (aq)	NH ₃ liberated on heating with OH ⁻ (aq) and Al foil; decolourises acidified aqueous KMnO ₄
sulfate, SO ₄ ²⁻ (aq)	gives white ppt. with Ba ²⁺ (aq) (insoluble in excess dilute strong acids); gives white ppt. with high [Ca ²⁺ (aq)]
sulfite, SO ₃ ²⁻ (aq)	gives white ppt. with Ba ²⁺ (aq) (soluble in excess dilute strong acids); decolourises acidified aqueous KMnO ₄
thiosulfate, S ₂ O ₃ ²⁻ (aq)	gives off-white/pale yellow ppt. slowly with H ⁺

3 Tests for gases

gas	test and test result
ammonia, NH_3	turns damp red litmus paper blue
carbon dioxide, CO_2	gives a white ppt. with limewater
hydrogen, H_2	'pops' with a lighted splint
oxygen, O_2	relights a glowing splint

4 Tests for elements

element	test and test result
iodine, I_2	gives blue-black colour on addition of starch solution

Important values, constants and standards

molar gas constant	$R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$
Faraday constant	$F = 9.65 \times 10^4 \text{ C mol}^{-1}$
Avogadro constant	$L = 6.022 \times 10^{23} \text{ mol}^{-1}$
electronic charge	$e = -1.60 \times 10^{-19} \text{ C}$
molar volume of gas	$V_m = 22.4 \text{ dm}^3 \text{ mol}^{-1}$ at s.t.p. (101 kPa and 273 K) $V_m = 24.0 \text{ dm}^3 \text{ mol}^{-1}$ at room conditions
ionic product of water	$K_w = 1.00 \times 10^{-14} \text{ mol}^2 \text{ dm}^{-6}$ (at 298 K (25 °C))
specific heat capacity of water	$c = 4.18 \text{ kJ kg}^{-1} \text{ K}^{-1}$ ($4.18 \text{ J g}^{-1} \text{ K}^{-1}$)

Group																																			
1	2		Key												13	14	15	16	17	18															
			atomic number atomic symbol name relative atomic mass																																
3	Li lithium 6.9	4 Be beryllium 9.0																																	
11	Na sodium 23.0	12 Mg magnesium 24.3	3	21	Sc scandium 45.0	22	Ti titanium 47.9	23	V vanadium 50.9	24	Cr chromium 52.0	25	Mn manganese 54.9	26	Fe iron 55.8	27	Co cobalt 58.9	28	Ni nickel 58.7	29	Cu copper 63.5	30	Zn zinc 65.4	31	Ga gallium 69.7	32	Ge germanium 72.6	33	As arsenic 74.9	34	Se selenium 79.0	35	Br bromine 79.9	36	Kr krypton 83.8
37	Rb rubidium 85.5	38 Sr strontium 87.6	39	Y yttrium 88.9	40	Zr zirconium 91.2	41	Nb niobium 92.9	42	Mo molybdenum 95.9	43	Tc technetium —	44	Ru ruthenium 101.1	45	Rh rhodium 102.9	46	Pd palladium 106.4	47	Ag silver 107.9	48	Cd cadmium 112.4	49	In indium 114.8	50	Sn tin 118.7	51	Sb antimony 121.8	52	Te tellurium 127.6	53	I iodine 126.9	54	Xe xenon 131.3	
55	Cs caesium 132.9	56 Ba barium 137.3	57–71	lanthanoids	72	Hf hafnium 178.5	73	Ta tantalum 180.9	74	W tungsten 183.8	75	Re rhenium 186.2	76	Os osmium 190.2	77	Ir iridium 192.2	78	Pt platinum 195.1	79	Au gold 197.0	80	Hg mercury 200.6	81	Tl thallium 204.4	82	Pb lead 207.2	83	Bi bismuth 209.0	84	Po polonium —	85	At astatine —	86	Rn radon —	
87	Fr francium —	88 Ra radium —	89–103	actinoids	104	Rf rutherfordium —	105	Db dubnium —	106	Sg seaborgium —	107	Bh bohrium —	108	Hs hassium —	109	Mt meitnerium —	110	Ds darmstadtium —	111	Rg roentgenium —	112	Cn copernicium —	113	Nh nihonium —	114	Fl flerovium —	115	Mc moscovium —	116	Lv livermorium —	117	Ts tennessine —	118	Og oganeson —	

lanthanoids

57	La	lanthanum	138.9
58	Ce	cerium	140.1
59	Pr	praseodymium	140.9
60	Nd	neodymium	144.4
61	Pm	promethium	—
62	Sm	samarium	150.4
63	Eu	europlum	152.0
64	Gd	gadolinium	157.3
65	Tb	terbium	158.9
66	Dy	dysprosium	162.5
67	Ho	holmium	164.9
68	Er	erbium	167.3
69	Tm	thulium	168.9
70	Yb	ytterbium	173.1
71	Lu	lutetium	175.0
72	Hf	hafnium	178.5
73	Ta	tantalum	180.9
74	W	tungsten	183.8
75	Re	rhenium	186.2
76	Os	osmium	190.2
77	Ir	iridium	192.2
78	Pt	platinum	195.1
79	Au	gold	197.0
80	Hg	mercury	200.6
81	Tl	thallium	204.4
82	Pb	lead	207.2
83	Bi	bismuth	209.0
84	Po	polonium	209.0
85	At	astatine	210.0
86	Rn	radon	222.0
87	Fr	francium	223.0
88	Ra	radium	226.0
89	Ac	actinium	227.0
90	Th	thorium	232.0
91	Pa	protactinium	231.0
92	U	uranium	238.0
93	Np	neptunium	237.0
94	Pu	plutonium	244.1
95	Am	americium	243.1
96	Cm	curium	247.1
97	Bk	berkelium	247.1
98	Cf	californium	251.1
99	Es	einsteinium	252.1
100	Fm	fermium	257.1
101	Md	mendelevium	258.1
102	No	nobelium	259.1
103	Lr	lawrencium	262.1

actinoids