
CHEMISTRY

9701/51

Paper 5 Planning, Analysis and Evaluation

October/November 2017

MARK SCHEME

Maximum Mark: 30

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge International will not enter into discussions about these mark schemes.



Cambridge International is publishing the mark schemes for the October/November 2017 series for most Cambridge IGCSE[®], Cambridge International A and AS Level components and some Cambridge O Level components.

Question	Answer	Marks
1(a)(i)	mass = $228.2 \times 1.00 \times (250/1000)$ = 57.1 g	1
1(a)(ii)	Distilled/deionised water must be mentioned somewhere for 2 marks to be given. Dissolve (all) the solid in a (suitable container) with (distilled) water	1
	Transfer / add to a <u>250 cm³</u> volumetric flask AND make to mark with (distilled) water	1
1(a)(iii)	(starch) gives a sharp 'end-point' / turns blue sharply / goes blue with volume of I ₂ invisible to naked eye	1
1(b)	volumes of (NH ₄) ₂ S ₂ O ₈ (aq) constant	1
	volumes of I ⁻ varying with range	1
	total volume constant, made up by water	1
1(c)(i)	mol I ₂ (aq) = $V(\text{S}_2\text{O}_3^{2-}) \times [\text{S}_2\text{O}_3^{2-}] / 2$ = $\left(\frac{0.005 \times 0.0050}{2} \right)$ = 1.25×10^{-5}	1
	$[\text{I}_2(\text{aq})] = \frac{\text{moles of I}_2(\text{aq})}{V_{\text{total}}} = \frac{1.25 \times 10^{-5}}{0.021} = 5.95 \times 10^{-4}$	1
	rate = $[\text{I}_2(\text{aq})] / \text{time}$ = $\frac{5.95 \times 10^{-4}}{134} = 4.44 \times 10^{-6}$	1
1(c)(ii)	repeat the experiment (and take average)	1
1(c)(iii)	% error = $\frac{2 \times 0.05}{5.0} \times 100\% = 2(.0) \%$	1

PUBLISHED

Question	Answer	Marks
1(d)	No thiosulfate had been added	1
1(e)	Ammonium persulfate must be stated along with its hazard and linked to the precaution. Ammonium persulfate is a skin irritant so wear gloves OR Ammonium persulfate is an irritant to the respiratory system; do the experiment in fume cupboard/face mask OR Ammonium persulfate is harmful if swallowed so avoid mouth contact/wear face mask OR Ammonium persulfate is oxidising so avoid contact with flammable /combustible materials. OR Ammonium persulfate is harmful /hazardous to the environment so do not dispose of down the drain /use (large quantities) of water to dilute before disposal	1

Question	Answer				Marks																																												
2(a)	<table border="1" data-bbox="342 220 1108 863"> <thead> <tr> <th data-bbox="342 220 533 355">time /s</th> <th data-bbox="533 220 723 355">burette reading /cm³</th> <th data-bbox="723 220 913 355">volume (of hydrogen) /cm³</th> <th data-bbox="913 220 1108 355">charge /C</th> </tr> </thead> <tbody> <tr><td>0</td><td>46.20</td><td>0.00</td><td>0</td></tr> <tr><td>50</td><td>41.20</td><td>5.00</td><td>40</td></tr> <tr><td>100</td><td>36.20</td><td>10.00</td><td>80</td></tr> <tr><td>150</td><td>31.45</td><td>14.75</td><td>120</td></tr> <tr><td>200</td><td>25.80</td><td>20.40</td><td>160</td></tr> <tr><td>250</td><td>20.80</td><td>25.40</td><td>200</td></tr> <tr><td>300</td><td>16.40</td><td>29.80</td><td>240</td></tr> <tr><td>350</td><td>11.45</td><td>34.75</td><td>280</td></tr> <tr><td>400</td><td>6.80</td><td>39.40</td><td>320</td></tr> <tr><td>450</td><td>1.50</td><td>44.70</td><td>360</td></tr> </tbody> </table> <p data-bbox="342 898 831 962">volumes of hydrogen correct to 2 d.p. charge correct</p>				time /s	burette reading /cm ³	volume (of hydrogen) /cm ³	charge /C	0	46.20	0.00	0	50	41.20	5.00	40	100	36.20	10.00	80	150	31.45	14.75	120	200	25.80	20.40	160	250	20.80	25.40	200	300	16.40	29.80	240	350	11.45	34.75	280	400	6.80	39.40	320	450	1.50	44.70	360	2
time /s	burette reading /cm ³	volume (of hydrogen) /cm ³	charge /C																																														
0	46.20	0.00	0																																														
50	41.20	5.00	40																																														
100	36.20	10.00	80																																														
150	31.45	14.75	120																																														
200	25.80	20.40	160																																														
250	20.80	25.40	200																																														
300	16.40	29.80	240																																														
350	11.45	34.75	280																																														
400	6.80	39.40	320																																														
450	1.50	44.70	360																																														
2(b)	All ten points plotted correctly				1																																												
	Best-fit straight line drawn				1																																												
2(c)	Yes, (the data is reliable because) most of the points are on the line OR only a few points are not on the line.				1																																												
2(d)(i)	co-ordinates read and recorded correctly				1																																												
	gradient determined				1																																												
2(d)(ii)	= (i) ÷ 24000				1																																												
2(d)(iii)	= 1 ÷ (2 × (ii))				1																																												

Question	Answer	Marks
2(e)(i)	 <p>straight line from origin to (300, 9.0)</p>	1
2(e)(ii)	Oxygen is (slightly) soluble in water	1
2(f)(i)	 <p>straight line with negative gradient</p>	1
2(f)(ii)	Faraday constant will be <u>lower</u> (than calculated) because the volume / V_m larger	1

PUBLISHED

Question	Answer	Marks
2(g)	No effect at cathode	1
	Less gas produced at anode	1
	Copper anode will dissolve / is (an) active (anode) OR copper has lower / more negative E^\ominus	1