

Cambridge  
International  
AS & A Level

**Cambridge International Examinations**  
Cambridge International Advanced Subsidiary and Advanced Level

CANDIDATE  
NAME

CENTRE  
NUMBER

--	--	--	--	--

CANDIDATE  
NUMBER

--	--	--	--

\* 5 8 2 4 8 8 7 6 2 5 9 \*



**CHEMISTRY**

**9701/22**

Paper 2 AS Level Structured Questions

**October/November 2018**

**1 hour 15 minutes**

Candidates answer on the Question Paper.

Additional Materials: Data Booklet

**READ THESE INSTRUCTIONS FIRST**

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

DO **NOT** WRITE IN ANY BARCODES.

Answer **all** questions.

Electronic calculators may be used.

You may lose marks if you do not show your working or if you do not use appropriate units.

A Data Booklet is provided.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [ ] at the end of each question or part question.

This document consists of **13** printed pages and **3** blank pages.

Answer **all** the questions in the spaces provided.

- 1 The model of the nuclear atom was first proposed by Ernest Rutherford. He developed this model on the basis of results obtained from an experiment using gold metal foil.

(a) Complete the table with information for two of the particles in an atom of  $^{197}\text{Au}$ .

particle	relative mass	relative charge	location within atom	total number in an atom of $^{197}\text{Au}$
electron	0.0005	-1		79
neutron			nucleus	

[4]

(b) State the type of bonding in gold.

..... [1]

(c) A sample of gold found in the earth consists of only one isotope.

(i) Explain what is meant by the term *isotopes*.


.....  
 .....  
 ..... [2]

(ii) A different sample of gold contains more than one isotope.

Suggest why this different sample of gold has the same **chemical** properties as the sample found in the earth.

.....  
 ..... [1]

- (d) *Tumbaga* is an alloy of copper and gold. A sample of tumbaga was analysed. The mass spectrum of the sample is shown.



- (i) Calculate the percentage abundance of gold,  $x$ , in the sample of tumbaga.

$$x = \dots\dots\dots \% \quad [1]$$

- (ii) Calculate the relative atomic mass,  $A_r$ , of the copper present in this sample. Give your answer to **two** decimal places.

$$A_r(\text{Cu}) = \dots\dots\dots [2]$$


[Total: 11]

2 The table gives some data for elements in the third period and some of their compounds.


element	Na	Mg	Al	Si	P	S
type of bonding	metallic				covalent	covalent
formula of oxide					P <sub>4</sub> O <sub>10</sub>	SO <sub>2</sub>
formula of chloride	NaCl	MgCl <sub>2</sub>				SCl <sub>2</sub>

(a) Complete the table to show the bonding in the elements, and the formulae of their oxides and chlorides. [3]

(b) SCl<sub>2</sub> is formed in the following reaction.




(i) Complete the 'dot-and-cross' diagram to show the bonding in a molecule of SCl<sub>2</sub>. Show outer electrons only.




[1]

(ii) Complete and fully label the reaction pathway diagram for the reaction between S<sub>2</sub>Cl<sub>2</sub> and Cl<sub>2</sub>. Include labels for activation energy, E<sub>a</sub>, and enthalpy change of the forward reaction, ΔH.



[2]

(c) (i) On the axes, sketch the trend in melting point of the elements Na to S.



[1]

(ii) Give three statements to explain your sketch.

- 1 .....
- .....
- 2 .....
- .....
- 3 .....
- .....

[3]

(d) Write an equation for the reaction of  $P_4O_{10}$  with water.

..... [1]

(e)  $SO_2$  can be released into the atmosphere when fossil fuels containing sulfur are burnt.

State and explain one environmental consequence of the release of  $SO_2$  into the atmosphere.

.....

.....

.....

..... [2]

- (f) The elements in the third period show a general increase in their first ionisation energies from left to right.

Identify **two** pairs of successive elements in the third period that do **not** agree with this statement.

For each pair, explain why the change in ionisation energy does **not** agree with this statement.

*Use of the Data Booklet may help you to answer this question.*

pair 1 .....

explanation .....

.....

.....

.....

.....

pair 2 .....

explanation .....

.....

.....

.....


.....

[4]

[Total: 17]

- 3 Trihalomethanes are organic molecules in which three of the hydrogen atoms of methane are replaced by halogen atoms, for example  $\text{CHF}_3$ .

(a) The equation shows a reaction to produce  $\text{CHF}_3$ .




Use the data to calculate the enthalpy change of reaction,  $\Delta H_r$ , for this formation of  $\text{CHF}_3$ .

compound	enthalpy change of formation, $\Delta H_f / \text{kJ mol}^{-1}$
$\text{CHI}_3(\text{s})$	-182.1
$\text{CHF}_3(\text{g})$	-692.9
$\text{AgF}(\text{s})$	-204.6
$\text{AgI}(\text{s})$	-61.8

enthalpy change of reaction,  $\Delta H_r = \dots\dots\dots \text{kJ mol}^{-1}$  [3]

(b) The graph shows the relationship between  $pV$  and  $p$  at a given temperature for  $\text{CHF}_3$  and an ideal gas.



(i)  $\text{CHF}_3$  is not an ideal gas.

State **three** basic assumptions that scientists make about the properties of ideal gases.

- 1 .....
- 2 .....
- 3 .....

[3]


(ii) Explain why  $\text{CHF}_3$  deviates from the properties of an ideal gas at pressures greater than 300 atm.

- .....
- .....
- .....
- .....

[2]



- (c) A different trihalomethane,  $\text{CHCl}_3$ , reacts with  $\text{O}_2$  to produce carbonyl dichloride.  $\text{HCl}(\text{g})$  is also released as a product of this reaction.




carbonyl dichloride

- (i) Write an equation for this reaction of  $\text{CHCl}_3$  with  $\text{O}_2$ .

..... [1]

- (ii) The conversion of  $\text{CHCl}_3$  to carbonyl dichloride can be monitored by infra-red spectroscopy. The infra-red spectrum of carbonyl dichloride is shown.



On the infra-red spectrum of carbonyl dichloride identify with an **X** the absorption that would **not** be present in an infra-red spectrum of  $\text{CHCl}_3$ .

Explain your answer.


.....  
 ..... [2]

- (iii) Suggest another difference between the infra-red spectra of  $\text{CHCl}_3$  and carbonyl dichloride.

.....  
 ..... [1]

[Total: 12]

4 The diagram shows a reaction sequence starting from ethanal.



(a) (i) Draw the **displayed** formula of **P**.

[1]

(ii) Name the type of chemical reaction that occurs in reaction 3.

..... [1]

(iii) Write an equation to represent reaction 4.

Use [O] to represent the oxidising agent.

..... [1]

(iv) State the reagents and conditions for reaction 4.

..... [1]

(b) Compound **Q** is formed as a mixture of two optical isomers.

(i) Explain what is meant by the term *optical isomers*.

.....

.....


..... [1]

(ii) Draw the **two** optical isomers of **Q**, showing clearly their three-dimensional structures.



[2]

(c) **R** can be used to make a polymer, **W**, in two steps.





Draw one repeat unit of **W**.

[3]

(d) Compound **Z**,  $\text{H}_2\text{C}=\text{CHCH}_3$ , is produced from **R**.

**Z** can be used in a two-step process to produce 2-aminopropane.

(i) In the first step, **Z** reacts with  $\text{HBr}$  to form two products. The structure of the product depends on which intermediate is formed, intermediate **I** or intermediate **II**.

intermediate **I**intermediate **II**

Explain why intermediate **I** is more likely to form than intermediate **II**.

.....


.....

.....

..... [2]


(ii) When intermediate **I** forms, the product of the first step is **T**.

Complete the diagram to show the mechanism for the conversion of **Z** to **T**. Include all relevant charges, partial charges, curly arrows and lone pairs.



[3]

(iii) **T** can then be converted to 2-aminopropane.




Name the mechanism for this conversion.

..... [1]


- (e) (i) Compound **S**,  $\text{CH}_3\text{COCO}_2\text{H}$ , can be reduced by  $\text{LiAlH}_4$ .

Complete the equation using structural formulae to represent this reaction.  
Use [H] to represent the reducing agent.



Other reducing agents containing Group 1 metal cations include  $\text{LiBH}_4$ ,  $\text{NaBH}_4$  and  $\text{KBH}_4$ .  
The strength of the reducing agent depends on the size of its cation.

- (ii) Give the electronic configuration of the  $\text{Na}^+$  cation.



- (iii) Suggest why ionic radius increases down Group 1.

.....  
.....  
..... [1]

[Total: 20]

**BLANK PAGE**

**BLANK PAGE**

**BLANK PAGE**

---

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at [www.cie.org.uk](http://www.cie.org.uk) after the live examination series.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.