
PHYSICS

9702/21

Paper 2 AS Level Structured Questions

May/June 2017

MARK SCHEME

Maximum Mark: 60

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the May/June 2017 series for most Cambridge IGCSE[®], Cambridge International A and AS Level and Cambridge Pre-U components, and some Cambridge O Level components.

PUBLISHED

Question	Answer	Marks
1(a)	(stress =) force / area or $\text{kg m s}^{-2} / \text{m}^2$	B1
	$= \text{kg m}^{-1} \text{s}^{-2}$	A1
1(b)(i)	$0.58 = 2\pi \times [(4 \times 0.500 \times 0.600^3) / (E \times 0.0300 \times 0.00500^3)]^{0.5}$	C1
	$E = [4\pi^2 \times 4 \times 0.500 \times (0.600)^3] / [(0.58)^2 \times 0.0300 \times (0.00500)^3]$ $= 1.35 \times 10^{10} \text{ (Pa)}$	C1
	$= 14 \text{ (13.5) GPa}$	A1
1(b)(ii)1.	(accuracy determined by) the closeness of the value(s)/measurement(s) to the true value	B1
	(precision determined by) the range of the values/measurements	B1
1(b)(ii)2.	l is (cubed so) $3 \times$ (percentage/fractional) uncertainty and T is (squared so) $2 \times$ (percentage / fractional) uncertainty and (so) l contributes more	B1

PUBLISHED

Question	Answer	Marks
2(a)	resultant force (in any direction) is zero	B1
	resultant torque/moment (about any point) is zero	B1
2(b)(i)	$a = (v - u) / t$ or gradient or $\Delta v / (\Delta)t$	C1
	e.g. $a = (8.8 - 4.6) / (7.0 - 4.0) = 1.4 \text{ m s}^{-2}$	A1
2(b)(ii)	$s = 4.6 \times 4 + [(8.8 + 4.6) / 2] \times 3$	C1
	$= 18.4 + 20.1$	A1
	$= 39 \text{ (38.5) m}$	
2(b)(iii)	$\Delta E = \frac{1}{2} \times 95 [(8.8)^2 - (4.6)^2]$	C1
	$= 3678 - 1005$	A1
	$= 2700 \text{ (2673) J}$	
2(b)(iv)1.	weight = 95×9.81 (= 932 N)	C1
	vertical tension force = $280 \sin 25^\circ$ or $280 \cos 65^\circ$ (=118.3 N)	C1
	$F = 932 + 118$ $= 1100 \text{ (1050) N}$	A1
2(b)(iv)2.	horizontal tension force = $280 \cos 25^\circ$ or $280 \sin 65^\circ$ (= 253.8 N)	C1
	resultant force = 95×1.4 (= 133 N)	C1
	$133 = 253.8 - R$ $R = 120 \text{ (120.8) N}$	A1

PUBLISHED

Question	Answer	Marks
3(a)	$\rho = m / V$	C1
	$V = \pi d^2 L / 4$ or $\pi r^2 L$	C1
	weight = $2.7 \times 10^3 \times \pi (1.2 \times 10^{-2})^2 \times 5.0 \times 10^{-2} \times 9.81 = 0.60 \text{ N}$	A1
3(b)(i)	the point from where (all) the weight (of a body) seems to act	B1
3(b)(ii)	$W \times 12$	C1
	$(0.25 \times 8) + (0.6 \times 38)$	C1
	$W = (2 + 22.8) / 12$ $= 2.1 \text{ (2.07) N}$	A1
3(c)(i)	pressure changes with depth (in water) or pressure on bottom (of cylinder) different from pressure on top	B1
	pressure on bottom of cylinder <u>greater than</u> pressure on top or force (up) on bottom of cylinder <u>greater than</u> force (down) on top	B1
3(c)(ii)	anticlockwise moment reduced and reducing the weight of X reduces clockwise moment or anticlockwise moment reduced so clockwise moment now greater than (total) anticlockwise moment	B1

PUBLISHED

Question	Answer	Marks
4(a)	(two) waves travelling (at same speed) in opposite directions overlap	B1
	waves (are same type and) have same frequency/wavelength	B1
4(b)(i)	$\lambda = 12 / 250 (= 0.048 \text{ m})$	C1
	distance = 1.5×0.048 = 0.072 m	A1
4(b)(ii)	$T = 1 / 250$ = 0.004 (s) or 4 (ms)	C1
	1. curve drawn is mirror image of that in Fig. 4.2 and labelled P	A1
	2. horizontal line drawn between A and B and labelled Q	A1

Question	Answer	Marks
5(a)	observed frequency is different to source frequency when source moves relative to observer	B1
5(b)	$360 = (400 \times 340) / (340 \pm v)$	C1
	$v = 38 (37.8) \text{ m s}^{-1}$	A1
	away (from the observer)	B1

PUBLISHED

Question	Answer	Marks
6(a)	volt / ampere	B1
6(b)(i)	$R_T = [1/3.0 + 1/6.0]^{-1} + 4.0 (= 6.0 \Omega)$	C1
	$I = 1.5 / 6.0$	C1
	$= 0.25 \text{ A}$	A1
6(b)(ii)	$V_B = 0.5 \text{ V}$	A1
	$I = 0.5 / 3.0$ $= 0.17 (0.167) \text{ A}$	
6(b)(iii)	$P = I^2 R$ or VI or V^2/R	C1
	ratio = $(0.167^2 \times 3.0) / (0.25^2 \times 4.0)$ $= 0.33$	A1
6(c)(i)	vary/change/different radius/diameter/ <u>cross-sectional</u> area (of wire)	B1
6(c)(ii)	$v = I / Ane$	C1
	ratio = $\frac{(I_B / A_B)}{(I_C / A_C)}$ or $\frac{I_B \times A_C}{I_C \times A_B}$	
	$(R \propto 1/A \text{ so})$ ratio = $\frac{I_B \times R_B}{I_C \times R_C} = \frac{0.167 \times 3.0}{0.25 \times 4.0}$ $= 0.50$	A1
6(d)(i)	0.25 A to 0.13 (0.125) A or halved	A1
6(d)(ii)	no change	A1

PUBLISHED

Question	Answer	Marks									
7(a)(i)	(proton is uud so) $(2/3)e + (2/3)e - (1/3)e = e$	B1									
7(a)(ii)	(neutron is udd so) $(2/3)e - (1/3)e - (1/3)e = 0$	B1									
7(b)(i)	<table border="1" data-bbox="808 347 1429 501"> <tbody> <tr> <td></td> <td>β^-</td> <td>β^+</td> </tr> <tr> <td>nucleon number</td> <td>90</td> <td>64</td> </tr> <tr> <td>proton number</td> <td>39</td> <td>28</td> </tr> </tbody> </table> <p><i>all correct</i></p>		β^-	β^+	nucleon number	90	64	proton number	39	28	B1
	β^-	β^+									
nucleon number	90	64									
proton number	39	28									
7(b)(ii)	weak (nuclear force/interaction)	B1									
7(b)(iii)	β^- decay: electron and (electron) antineutrino β^+ decay: positron and (electron) neutrino <i>all correct</i>	B1									