

**CAMBRIDGE**  
INTERNATIONAL EXAMINATIONS

**NOVEMBER 2002**

**GCE Advanced Subsidiary Level**

<b>MARK SCHEME</b>
<b>MAXIMUM MARK : 50</b>
<b>SYLLABUS/COMPONENT : 9709 / 2</b> <b>MATHEMATICS</b> <b>(Pure 2)</b>



Page 1	Mark Scheme	Syllabus	Paper
	AS Level Examinations – November 2002	9709	2

1	<i>EITHER:</i> State or imply non-modular inequality $(2x - 1)^2 < (3x)^2$ , or corresponding equation	B1	4
	Expand and make reasonable solution attempt at <del>2/3</del> 3-term quadratic, or equivalent	M1	
	Obtain critical values $-1$ and $\frac{1}{5}$	A1	
	State correct answer $x < -1, x > \frac{1}{5}$	A1	
	<i>OR:</i> State <del>one</del> correct equation for a critical value e.g. $2x - 1 = 3x$	M1	
	State two relevant equations separately e.g. $2x - 1 = 3x$ and $2x - 1 = -3x$	A1	
	Obtain critical values $-1$ and $\frac{1}{5}$	A1	
	State correct answer $x < -1, x > \frac{1}{5}$	A1	
	<i>OR:</i> State one critical value (probably $x = -1$ ), from a graphical method or by inspection or by solving a linear inequality	B1	
	State the other critical value correctly	B2	
State correct answer $x < -1, x > \frac{1}{5}$	B1		
[The answer $\frac{1}{5} < x < -1$ scores B0.]			
<hr/>			
2	State or obtain $-2 + a + b = 0$ , or equivalent	B1	5
	Substitute $x = -2$ and equate to $-5$	M1	
	Obtain 3-term equation, or equivalent	A1	
	Solve a relevant pair of equations, obtaining $a$ or $b$	M1	
	Obtain both answers $a = 3$ and $b = -1$	A1	
<hr/>			
3	(i) State or imply that $9^x = y^2$	B1	5
	(ii) Carry out recognisable solution method for quadratic in $y$	M1	
	Obtain $y = \frac{1}{2}$ and $y = 3$ from $2y^2 - 7y + 3 = 0$	A1	
	Use log method to solve an equation of the form $3^x = k$	M1	
	Obtain answer $x = -\frac{\ln 2}{\ln 3}$ , or exact equivalent {To ANY base}	A1	
	State exact answer $x = 1$ (no penalty if logs used)	B1	
<hr/>			
4	(i) Make recognisable sketches over the given range of a suitable pair of graphs e.g. $y = \sin x$ and $y = \frac{1}{x^2}$	B1	3
	State or imply connection between intersections and roots and justify given statement	B1	
	(ii) Calculate values (or signs) of $\sin x - \frac{1}{x^2}$ at $x = 1$ and $x = 1.5$	M1	
	Derive given result correctly	A1	
	(iii) Rearrange $\sin x = \frac{1}{x^2}$ and obtain given answer	B1	
	(iv) Use the iterative formula correctly with $1 \leq x_n \leq 1.5$	M1	
	Obtain final answer 1.07	A1	
	Show sufficient iterations to justify its accuracy to 3d.p., or show there is a sign change in the interval (1.065, 1.075)	A1	

Page 2	Mark Scheme	Syllabus	Paper
	AS Level Examinations – November 2002	9709	2

- 5 (i) Use relevant formulae for  $\cos(x - 30^\circ)$  and  $\sin(x - 60^\circ)$  { allow ONE sign error } M1\*  $\odot$   
 Use  $\sin 30^\circ = \cos 60^\circ = \frac{1}{2}$  and  $\sin 60^\circ = \cos 30^\circ = \frac{\sqrt{3}}{2}$  M1(dep\*)  
 Collect terms and obtain given answer correctly A1 3  
 (ii) Carry out correct processes to evaluate a single trig ratio M1  
 Obtain answer  $73.9^\circ$  A1  
 Obtain second answer  $253.9^\circ$  and no others A1 ✓ 3  
 (iii) State or imply that  $\cos^2 x = \frac{1}{13}$  or  $\sin^2 x = \frac{12}{13}$  B1  
 Use a relevant trig formula to evaluate  $\cos 2x$  M1  
 Obtain exact answer  $-\frac{11}{13}$  correctly A1 3  
 [Use of only say  $\cos x = +\frac{1}{\sqrt{13}}$ , probably from a right triangle, can earn B1M1A0.]

- 6 (a) Obtain indefinite integral  $-\frac{1}{2} \cos 2x + \sin x$  B1 + B1  
 Use limits with attempted integral M1  
 Obtain answer 2 correctly with no errors A1 4  
 (b) (i) Identify  $R$  with correct definite integral and attempt to integrate M1  
 Obtain indefinite integral  $\ln(x + 1)$  B1  
 Obtain answer  $R = \ln(p + 1) - \ln 2$  A1 3  
 (ii) Use exponential method to solve an equation of the form  $\ln x = k$  M1  
 Obtain answer  $p = 13.8$  A1 2

- 7 (i) State  $6y \frac{dy}{dx}$  as the derivative of  $3y^2$  B1  
 State  $\pm 2x \frac{dy}{dx} \pm 2y$  as the derivative of  $-2xy$  (allow any combination of signs here) B1  
 Equate attempted derivative of LHS to 0 (or 10) and solve for  $\frac{dy}{dx}$  M1  
 Obtain the given answer correctly A1 4  
 [The M1 is dependent on at least one of the B marks being earned.]  
 (ii) State or imply the points lie on  $y - 2x = 0$  or  $(y - 2x) / (3y - 2x) = 0$  B1  $\odot$   
 Carry out complete method for finding one coordinate of a point of intersection of  $y = kx$  with the given curve M1  
 Obtain  $10x^2 = 10$  or  $2\frac{1}{2}y^2 = 10$  or 2-term equivalent A1  
 Obtain one correct point e.g. (1, 2) or 2 values of  $x$  (or  $y$ ) A1  
 Obtain a second correct point e.g. (-1, -2) A1  $\sqrt{7}$  5  $\odot$