

**CAMBRIDGE INTERNATIONAL EXAMINATIONS**

**Cambridge International Advanced Subsidiary and Advanced Level**

**MARK SCHEME for the October/November 2014 series**

**9709 MATHEMATICS**

**9709/61**

Paper 6, maximum raw mark 50

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the October/November 2014 series for most Cambridge IGCSE®, Cambridge International A and AS Level components and some Cambridge O Level components.

® IGCSE is the registered trademark of Cambridge International Examinations.

Page 2	Mark Scheme	Syllabus	Paper
	Cambridge International AS/A Level – October/November 2014	9709	61

## Mark Scheme Notes

Marks are of the following three types:

- M Method mark, awarded for a valid method applied to the problem. Method marks are not lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. Correct application of a formula without the formula being quoted obviously earns the M mark and in some cases an M mark can be implied from a correct answer.
  - A Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated method mark is earned (or implied).
  - B Mark for a correct result or statement independent of method marks.
- When a part of a question has two or more “method” steps, the M marks are generally independent unless the scheme specifically says otherwise; and similarly when there are several B marks allocated. The notation DM or DB (or dep\*) is used to indicate that a particular M or B mark is dependent on an earlier M or B (asterisked) mark in the scheme. When two or more steps are run together by the candidate, the earlier marks are implied and full credit is given.
- The symbol  $\dagger$  implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A or B marks are given for correct work only. A and B marks are not given for fortuitously “correct” answers or results obtained from incorrect working.
- Note: B2 or A2 means that the candidate can earn 2 or 0.  
B2/1/0 means that the candidate can earn anything from 0 to 2.
- The marks indicated in the scheme may not be subdivided. If there is genuine doubt whether a candidate has earned a mark, allow the candidate the benefit of the doubt. Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored.
- Wrong or missing units in an answer should not lead to the loss of a mark unless the scheme specifically indicates otherwise.
- For a numerical answer, allow the A or B mark if a value is obtained which is correct to 3 s.f., or which would be correct to 3 s.f. if rounded (1 d.p. in the case of an angle). As stated above, an A or B mark is not given if a correct numerical answer arises fortuitously from incorrect working. For Mechanics questions, allow A or B marks for correct answers which arise from taking  $g$  equal to 9.8 or 9.81 instead of 10.

Page 3	Mark Scheme	Syllabus	Paper
	Cambridge International AS/A Level – October/November 2014	9709	61



The following abbreviations may be used in a mark scheme or used on the scripts:

- AEF Any Equivalent Form (of answer is equally acceptable)
- AG Answer Given on the question paper (so extra checking is needed to ensure that the detailed working leading to the result is valid)
- BOD Benefit of Doubt (allowed when the validity of a solution may not be absolutely clear)
- CAO Correct Answer Only (emphasising that no “follow through” from a previous error is allowed)
- CWO Correct Working Only – often written by a ‘fortuitous’ answer
- ISW Ignore Subsequent Working
- MR Misread
- PA Premature Approximation (resulting in basically correct work that is insufficiently accurate)
- SOS See Other Solution (the candidate makes a better attempt at the same question)
- SR Special Ruling (detailing the mark to be given for a specific wrong solution, or a case where some standard marking practice is to be varied in the light of a particular circumstance)

### Penalties

- MR –1 A penalty of MR –1 is deducted from A or B marks when the data of a question or part question are genuinely misread and the object and difficulty of the question remain unaltered. In this case all A and B marks then become “follow through” marks. MR is not applied when the candidate misreads his own figures – this is regarded as an error in accuracy. An MR –2 penalty may be applied in particular cases if agreed at the coordination meeting.
- PA –1 This is deducted from A or B marks in the case of premature approximation. The PA –1 penalty is usually discussed at the meeting.

Page 4	Mark Scheme	Syllabus	Paper
	Cambridge International AS/A Level – October/November 2014	9709	61

<b>1</b>	mean = $(5 + (-2) + 12 + 7 + (-3) + 2 + (-6) + 4 + 0 + 8) / 10$ = 2.7 var = $(5^2 + (-2)^2 + \dots + 8^2) / 10 - 2.7^2 = 35.1 - 2.7^2$ = 27.8	B1  M1  A1	<b>3</b>	Subst in correct var formula must have – mean <sup>2</sup> Correct answer
<b>2</b>	(i) $0.24 + 0.35 + 2k + k + 0.05 = 1$ $k = 0.12$  (ii) model number is 1  (iii) mean = $1 \times 0.35 + 2 \times 0.24 + 3 \times 0.12 + 4 \times 0.05$ $P(>1.39) = P(2, 3, 4) = 0.41$	M1  A1  B1  B1  M1  B1	<b>2</b>  <b>1</b>  <b>3</b>	Summing probs = 1 Correct answer  1.39 seen Finding $P(X > \text{their mean})$ Correct ans following mean or mode only
<b>3</b>	$P(8) = P(H \ 4 \ 4) + P(T \ 2 \ 4) + P(T \ 4 \ 2)$  $= \frac{1}{3} \times \frac{1}{16} + \frac{2}{3} \times \frac{1}{16} + \frac{2}{3} \times \frac{1}{16}$  $= \frac{5}{48}$  $P(H   8) = \frac{P(H \cap 8)}{P(8)}$  $= \frac{\frac{1}{48}}{\frac{5}{48}} = \frac{1}{5}$	M1  M1  A1  B1  A1		$\frac{1}{3}$ or $\frac{2}{3}$ mult by dice related prob, seen anywhere  Summing two or three 2-factor probs involving $\frac{1}{3}$ and $\frac{2}{3}$  $\frac{5}{48}$ oe seen as num or denom of a fraction  $\frac{1}{48}$ oe seen as num or denom of a fraction  Correct ans
<b>4</b>	(i) median A = 0.52 LQ = 0.41 UQ = 0.79  (ii)      	B1  B1  B1ft  B1  B1  B1	<b>3</b>	ft wrong units  2 correct boxes ft (i) OK if superimposed  2 pairs correct whiskers lines up to box not inside  Correct uniform scale need at least 4 values on it. No scale no marks unless perfect A and B with all 10 values shown, in which case score B1B1B0

Page 5	Mark Scheme	Syllabus	Paper
	Cambridge International AS/A Level – October/November 2014	9709	61

(iii) Smartphone B is quicker, slightly less variable, etc.	B1   	1	oe sensible answer
5 (i) $1.2 = 15p$ $p = 0.08$ $\text{Var} = npq = 15 \times 0.08 \times 0.92 = 1.104$ <b>AG</b>	M1   A1	2	Attempt to find $p$ using $1.2 = 15p$  Correct answer
(ii) $P(0, 1, 2) = \frac{(0.92)^{15}}{15!} + {}^{15}C_1(0.08)(0.92)^{14}$ $= {}^{15}C_2(0.08)^2(0.92)^{13}$ $= 0.887$	M1   A1	3	Binomial expression ${}^{15}C_x p^x (1-p)^{15-x}$ $0 < p < 1$ Correct unsimplified expression for $P(0, 1, 2)$ Correct answer
(iii) $P(\text{at least 1 faulty screw}) = 1 - P(0) = 1 - (0.92)^{15}$ $= 0.7137\dots$ $P(\text{at least 1 faulty screw in 7 packets}) = {}^8C_7(0.713\dots)^7(0.2863\dots)$ $= 0.216$	M1   A1 M1  A1	4	Attempt at $P(0)$ or $1 - P(0)$  Rounding to 0.71 Binomial expression ${}^8C_7 p^7 (1-p)$ $0 < p < 1$ Correct answer
6 (i) $z_1 = \frac{70 - 66.4}{5.6} = 0.6429$ $z_2 = \frac{72.5 - 66.4}{5.6} = 1.089$ $\Phi(1.089) - \Phi(0.643) = 0.8620 - 0.7399$ $= 0.1221$ $0.1221 \times 250 = 30.5$ 30 or 31 sheep	M1   A1  M1 A1ft	5	Standardising one variable, no cc, no sq rt  Correct area $\Phi_2 - \Phi_1$  Correct answer rounding to 0.12  Mult by 250 Correct answer ft their 0.1221
(ii) $66.4 - 59.2 = 7.2$ $66.4 + 7.2 = 73.6$	M1 A1	2	Subt from 66.4 Correct answer
(iii) $z = 0.674$ $\frac{67.5 - \mu}{4.92} = 0.674$ $\mu = 64.2$	B1   M1 A1	3	$\pm 0.674$ or 0.675 seen  Standardising with a $z$ -value no cc no sq rt Correct answer
7 (i) W(8) M(5) $4 \quad 2 = {}^8C_4 \times {}^5C_2 = 700$ $5 \quad 1 = {}^8C_5 \times {}^5C_1 = 280$ $6 \quad 0 = {}^8C_6 \times {}^5C_0 = 28$ Total = 1008	M1   A1 A1	4	Mult 2 combs, ${}^8C_x \times {}^5C_y$ Summing 2 or 3 options 2 correct options unsimplified Correct answer
(ii) M1 and MMWWW = ${}^3C_2 \times {}^8C_3 = 168$ M2 and MMWWW = ${}^3C_2 \times {}^8C_3 = 168$ Neither and MMMWWW = ${}^3C_1 \times {}^8C_3 = 56$ Total = 392	M1   B1 A1	3	Summing 3 options  One correct option  Correct answer
OR total, no restrictions = ${}^5C_3 \times {}^8C_3 = 560$ M1M2 and MWWW = ${}^3C_1 \times {}^8C_3 = 168$ $560 - 168 = 392$	M1   B1 A1		Subt 2 men together from no restrictions  One correct of 560 or 168 Correct answer

Page 6	Mark Scheme	Syllabus	Paper
	<b>Cambridge International AS/A Level – October/November 2014</b>	<b>9709</b>	<b>61</b>

<p>(iii) e.g. WWMWWW  <math>= 5! \text{ (women)} \times 4 = 480</math></p>	M1 M1 A1	<b>3</b>	5! Seen mult by integer $\geq 1$ Mult by 4 Correct answer
OR $6! - MWWWWW - WWWWWM$ $= 6! - 5! - 5!$ $= 480$	M1 M1 A1		6! seen with a subtraction 5! or $2 \times 5!$ Seen subtracted Correct answer