


Question Number	Scheme	Marks
<p>4. (a)</p> <p>(b)</p> <p>(c)</p>	$M(B), N 2a \cos \theta = W a \cos \theta + \frac{1}{4} W \frac{3a}{2} \sin \theta$ $N = \frac{7W}{8}$ $R = \frac{1}{4} W; \quad F + N = W$ $F \leq \mu R \text{ or } F = \mu R$ $\frac{1}{2} \leq \mu^* \text{ (exact)}$ <p>It does not bend</p> <p>Or has negligible thickness</p>	<p>M1 A2 (-1 e.e.)</p> <p>dep. M1 A1 (5)</p> <p>B1; B1</p> <p>M1</p> <p>A1 c.s.o. (5)</p> <p>B1 (1)</p> <p style="text-align: right;">(10 marks)</p>
<p>5. (a)</p> <p>(b)</p> <p>(c)</p> <p>OR</p>	$2ut = 735$ $0 = 3ut - \frac{1}{2} gt^2$ <p>eliminating t</p> $u = 24.5^*$ $t = \frac{735}{49} = 15$ <p>Initially: $v^2 = (2u)^2 + (3u)^2$</p> <p>(7803.25)</p> $\frac{1}{2} mv^2 - \frac{1}{2} m 65^2 = mgh$ $h = 180 \text{ m (183 m)}$ <p>(1824)</p> $v_y^2 = 65^2 - (2u)^2$ $v_y^2 = (3u)^2 - 2gh$ $h = 180 \text{ m (183 m)}$	<p>M1 A1</p> <p>M1 A1</p> <p>dep. M1</p> <p>A1 (6)</p> <p>M1 A1 (2)</p> <p>M1</p> <p>M1 A1</p> <p>A1 (4)</p> <p>M1</p> <p>M1 A1</p> <p>A1 (4)</p> <p style="text-align: right;">(12 marks)</p>

(ft = follow through mark; cao = correct answer only; (*) indicates final line is given on the paper)

Question Number	Scheme	Marks
6. (a)	$u \rightarrow \rightarrow 0$ CLM: $mu = mv_1 + 3mv_2$ $m \quad 3m$ NIL: $eu = -v_1 + v_2$ $v_1 \rightarrow \quad v_2 \rightarrow$ solving, $v_2 = \frac{u}{4}(1 + e)^*$	B1 M1 A1 dep. M1 A1 (5)
(b)	Solving for v_1 ; $\left \frac{u}{4}(1 - 3e) \right $	M1 A1 (2)
(c)	$\frac{1}{2} m \frac{u^2}{16} (1 - 3e)^2 + \frac{1}{2} 3m \frac{u^2}{16} (1 + e)^2 = \frac{1}{6} mu^2$ $e^2 = \frac{1}{9}$ $e = \frac{1}{3}$	M1 A1 f.t. A1 dep. M1 A1 A1 (6)
(d)	$v_1 = \frac{u}{4} (1 - 3 \times \frac{1}{3}) = 0 \Rightarrow$ at rest.	A1 c.s.o. (1) (14 marks)

Question Number	Scheme	Marks
7. (a)	$AD: 10m\bar{x} = 3m \frac{5a}{2} + 3m \times 5a$ $\bar{x} = 2.25a *$	M1 A1 A1 (3)
(b)	$AB: 10m\bar{y} = 2m \times 2a + 3m \times a$ $\bar{y} = 0.7a$	M1 A1 (2)
(c)	$\tan \theta = \frac{2.5a - \bar{x}}{\bar{y}}$ $\theta = 20^\circ$	M1 A1 f.t. A1 (3)
(e)	 $M(0), 10mg \times \frac{a}{4} = P \times 2a$ (OR: $4mg \times \frac{5a}{2} - 3mg \times \frac{5a}{2} = P \times 2a$) $P = \frac{5mg}{4} *$ (exact) $S = \frac{5mg}{4}; R = 10mg$ $F = \sqrt{S^2 + R^2} = \frac{5mg\sqrt{65}}{4}$ (10.1 mg)	M1 A1 A1 A1 (4) B1; B1 M1 A1 (4) (16 marks)