

# Mark Scheme (Results)

June 2011

GCE Mechanics M1 (6677) Paper 1

Edexcel is one of the leading examining and awarding bodies in the UK and throughout the world. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers.

Through a network of UK and overseas offices, Edexcel's centres receive the support they need to help them deliver their education and training programmes to learners.

For further information, please call our GCE line on 0844 576 0025, our GCSE team on 0844 576 0027, or visit our website at [www.edexcel.com](http://www.edexcel.com).

If you have any subject specific questions about the content of this Mark Scheme that require the help of a subject specialist, you may find our **Ask The Expert** email service helpful.

Ask The Expert can be accessed online at the following link:  
<http://www.edexcel.com/Aboutus/contact-us/>

June 2011

Publications Code UA028437

All the material in this publication is copyright

© Edexcel Ltd 2011

## EDEXCEL GCE MATHEMATICS

### General Instructions for Marking


1. The total number of marks for the paper is 75.
2. The Edexcel Mathematics mark schemes use the following types of marks:
  - **M** marks: method marks are awarded for 'knowing a method and attempting to apply it', unless otherwise indicated.
  - **A** marks: Accuracy marks can only be awarded if the relevant method (M) marks have been earned.
  - **B** marks are unconditional accuracy marks (independent of M marks)
  - Marks should not be subdivided.


3. Abbreviations


These are some of the traditional marking abbreviations that will appear in the mark schemes and can be used if you are using the annotation facility on ePEN.

- bod – benefit of doubt
- ft – follow through
- the symbol  $\frac{\Delta}{\Delta}$  will be used for correct ft
- cao – correct answer only
- cso - correct solution only. There must be no errors in this part of the question to obtain this mark
- isw – ignore subsequent working
- awrt – answers which round to
- SC: special case
- oe – or equivalent (and appropriate)
- dep – dependent
- indep – independent
- dp decimal places
- sf significant figures
- \* The answer is printed on the paper
- $\square$  The second mark is dependent on gaining the first mark

June 2011  
 Mechanics M1 6677  
 Mark Scheme

Question Number	Scheme	Marks
<b>1.</b>		
<b>(a)</b>	$0^2 = u^2 - 2 \times 9.8 \times 40$ $u = 28 \text{ m s}^{-1}$ ** GIVEN ANSWER	M1 A1 A1 (3)
<b>(b)</b>	$33.6 = 28t - \frac{1}{2}9.8t^2$ $4.9t^2 - 28t + 33.6 = 0$ $t = \frac{28 \pm \sqrt{28^2 - 4 \times 4.9 \times 33.6}}{9.8}$ $= 4 \text{ s or } (1.7 \text{ s or } 1.71 \text{ s})$	M1 A1  M1 A1 A1 (5) <b>8</b>
<b>2.</b>		
<b>(a)</b>	 CLM: $3 \times 3 - 2 \times 2 = 3v + 2(v+1)$ $v_P = 0.6 \text{ m s}^{-1}; v_Q = 1.6 \text{ m s}^{-1}$	M1 A1 M1A1 (A1 ft) (5)
<b>(b)</b>	$3(v-3) \quad \text{OR} \quad 2(v+1-2)$ $= 7.2 \text{ N s} \quad \quad = 7.2 \text{ N s}$	M1 A1 ft A1 (3) <b>8</b>

Question Number	Scheme	Marks
3.	$\begin{aligned} \nearrow & 4\cos\alpha + F = W\sin\alpha \\ \nwarrow & R = 4\sin\alpha + W\cos\alpha \\ & F = 0.5R \\ & \cos\alpha = 0.8 \quad \text{or} \quad \sin\alpha = 0.6 \\ & R = 20\text{N} \quad ** \quad \text{GIVEN ANSWER} \\ & W = 22\text{N} \end{aligned}$	M1 A1 M1 A1 B1 B1 M1 A1 A1 (9)
<u>OR</u>	$\begin{aligned} \rightarrow & R\sin\alpha = 4 + F\cos\alpha \\ \uparrow & R\cos\alpha + F\sin\alpha = W \\ & F = 0.5R \\ & \cos\alpha = 0.8 \quad \text{or} \quad \sin\alpha = 0.6 \\ & R = 20\text{N} \quad ** \quad \text{GIVEN ANSWER} \\ & W = 22\text{N} \end{aligned}$	M1 A1 M1 A1 B1 B1 M1 A1 A1 (9)
4.		9
(a)		B1 shape B1 figs (2)
(b)	$\left(\frac{1}{2} \times 4 \times 5\right) + 60 \times 5$ $= 310$	M1 A1 A1 (3)
(c)	$\frac{(5+V)}{2} \times 20 = (400-310)$ $V = 4$	M1 A2 ft DM1 A1 (5)
(d)	$\frac{5-4}{20} = 0.05 \text{ ms}^{-2}$	M1 A1 (2)
		12

Question Number	Scheme	Marks
5.		
(a)		
(i)	<b>EITHER</b> $M(R), 8X + 2X = 40g \times 6 + 20g \times 4$ solving for $X$ , $X = 32g = 314$ or $310$ N	M1 A2 M1 A1
(ii)	( $\uparrow$ ) $X + X = 40g + 20g + Mg$ (or another moments equation) solving for $M$ , $M = 4$	M1 A2 M1 A1
(i)	<b>OR</b> $M(P), 6X = 40g \times 2 + 20g \times 4 + Mg \times 8$ solving for $X$ , $X = 32g = 314$ or $310$ N	M1 A2 M1 A1
(ii)	( $\uparrow$ ) $X + X = 40g + 20g + Mg$ (or another moments equation) solving for $M$ , $M = 4$	M1 A2 M1 A1
		(10)
(b)	Masses concentrated at a point or weights act at a point	B1 (1)
		<b>11</b>
6.		
(a)	$R = 0.3g \cos \alpha$ $= 0.24g = 2.35$ (3sf)=2.4 (2sf)	M1 A1 (2)
(b)	$mg - T = 1.4m$ $T - 0.3g \sin \alpha - F = 0.3 \times 1.4$ $F = 0.5R$ Eliminating $R$ and $T$ $m = 0.4$	M1 A1 M1 A2 M1 <b>DM1</b> A1 (8)
(c)	$v = 1.4 \times 0.5$ $-0.3g \sin \alpha - F = 0.3a$ $a = -9.8$ $0 = 0.7 - 9.8t$ $t = 0.071$ s or $0.0714$ s (1/14 A0)	B1 M1 A1 A1 M1 A1 (6)
		<b>16</b>

Question Number	Scheme	Marks
7.		
(a)	$\tan \theta = \frac{3}{4}$ ; bearing is $37^\circ$ ( <b>nearest degree</b> )	M1; A1 (2)
(b)		
(i)	$\mathbf{p} = (\mathbf{i} + \mathbf{j}) + t(2\mathbf{i} - 3\mathbf{j})$	M1 A1
(ii)	$\mathbf{q} = (-2\mathbf{j}) + t(3\mathbf{i} + 4\mathbf{j})$	A1
(iii)	$\mathbf{PQ} = \mathbf{q} - \mathbf{p} = (-\mathbf{i} - 3\mathbf{j}) + t(\mathbf{i} + 7\mathbf{j})$	M1 A1 (5)
(c)		
(i)	$-1 + t = 0$ $t = 1$ or 3pm	M1 A1
(ii)	$-1 + t = -(-3 + 7t)$ $t = \frac{1}{2}$ or 2.30 pm	M1 A1 (4) <b>11</b>

Further copies of this publication are available from  
Edexcel Publications, Adamsway, Mansfield, Notts, NG18 4FN

Telephone 01623 467467

Fax 01623 450481

Email [publication.orders@edexcel.com](mailto:publication.orders@edexcel.com)

Order Code UA028437 June 2011

For more information on Edexcel qualifications, please visit  
[www.edexcel.com/quals](http://www.edexcel.com/quals)

Pearson Education Limited. Registered company number 872828  
with its registered office at Edinburgh Gate, Harlow, Essex CM20 2JE



Llywodraeth Cynulliad Cymru  
Welsh Assembly Government

