

GCE Examinations
Advanced Subsidiary

Core Mathematics C3

Paper C

MARKING GUIDE

This guide is intended to be as helpful as possible to teachers by providing concise solutions and indicating how marks could be awarded. There are obviously alternative methods that would also gain full marks.

Method marks (M) are awarded for knowing and using a method.

Accuracy marks (A) can only be awarded when a correct method has been used.

(B) marks are independent of method marks.



Written by Shaun Armstrong

© *Solomon Press*

These sheets may be copied for use solely by the purchaser's institute.

C3 Paper C – Marking Guide

1. (a)
$$= \frac{x+4}{(2x+1)(x+1)} - \frac{2}{2x+1}$$

$$= \frac{(x+4) - 2(x+1)}{(2x+1)(x+1)}$$

$$= \frac{2-x}{(2x+1)(x+1)}$$
M1
M1
A1

(b)
$$\frac{2-x}{(2x+1)(x+1)} = \frac{1}{2}$$

$$2(2-x) = 2x^2 + 3x + 1$$

$$2x^2 + 5x - 3 = 0$$

$$(2x-1)(x+3) = 0$$

$$x = -3, \frac{1}{2}$$
M1
M1
A1 (6)

2. (a) if $\theta = \frac{\pi}{2}$, $\sin \theta = 1$, $\operatorname{cosec} \theta = 1$

$$\therefore \operatorname{cosec} \theta - \sin \theta = 1 - 1 = 0$$

$$\therefore \text{statement is false}$$
M1
A1

(b)
$$1 - \sin^2 \theta = 2 \sin \theta$$

$$\sin^2 \theta + 2 \sin \theta - 1 = 0$$

$$\sin \theta = \frac{-2 \pm \sqrt{4+4}}{2} = -1 - \sqrt{2} \text{ (no solutions) or } -1 + \sqrt{2}$$

$$\theta = 0.4271, \pi - 0.4271$$

$$\theta = 0.43, 2.71 \text{ (2dp)}$$
M1
M1
M1 A1
M1
A1 (7)

3. (a)
$$2x - 3 = e$$

$$x = \frac{1}{2}(e + 3)$$
M1
M1 A1

(b)
$$3e^{2y} - 16e^y + 5 = 0$$

$$(3e^y - 1)(e^y - 5) = 0$$


$$e^y = \frac{1}{3}, 5$$


$$y = \ln \frac{1}{3}, \ln 5$$
M1
M1
A1
M1 A1 (8)

4. (a)
$$= \frac{1}{3x-2} \times 3 = \frac{3}{3x-2}$$
M1 A1

(b)
$$= \frac{2 \times (1-x) - (2x+1) \times (-1)}{(1-x)^2} = \frac{3}{(1-x)^2}$$
M1 A2

(c)
$$= \frac{3}{2}x^{\frac{1}{2}} \times e^{2x} + x^{\frac{3}{2}} \times 2e^{2x} = \frac{1}{2}x^{\frac{1}{2}} e^{2x}(3 + 4x)$$
M1 A2 (8)



6. (a) $4 - \ln 3x = 0, \quad \ln 3x = 4, \quad x = \frac{1}{3}e^4$ M1 A1
- (b)  B2
- (c) $y = 4 - \ln 3x$
 $\ln 3x = 4 - y$ M1
 $x = \frac{1}{3}e^{4-y}$ M1
 $\therefore f^{-1}(x) = \frac{1}{3}e^{4-x}$ A1
- (d) $fg(x) = 4 - \ln 3e^{2-x}$
 $= 4 - (\ln 3 + \ln e^{2-x})$ M1
 $= 4 - \ln 3 - (2 - x)$ M1
 $= x + 2 - \ln 3 \quad [a = 2, b = 3]$ A1 (10)

7. (a) $4 \sin x + 3 \cos x = R \sin x \cos \alpha + R \cos x \sin \alpha$
 $R \cos \alpha = 4, \quad R \sin \alpha = 3$
 $\therefore R = \sqrt{4^2 + 3^2} = 5$ M1 A1
 $\tan \alpha = \frac{3}{4}, \quad \alpha = 0.644 \text{ (3sf)}$ M1 A1
 $\therefore 4 \sin x + 3 \cos x = 5 \sin(x + 0.644)$
- (b) minimum = -5 B1
occurs when $x + 0.6435 = \frac{3\pi}{2}, \quad x = 4.07 \text{ (3sf)}$ M1 A1
- (c) $5 \sin(2\theta + 0.6435) = 2$
 $\sin(2\theta + 0.6435) = 0.4$ M1
 $2\theta + 0.6435 = \pi - 0.4115, 2\pi + 0.4115$ B1 M1
 $2\theta = 2.087, 6.051$ M1
 $\theta = 1.04, 3.03 \text{ (2dp)}$ A2 (13)

8. (a) $\frac{dy}{dx} = \frac{1}{2}x^{-\frac{1}{2}} - 4e^{1-4x}$ M1
grad = -3, grad of normal = $\frac{1}{3}$ A1
 $\therefore y - \frac{3}{2} = \frac{1}{3}(x - \frac{1}{4}) \quad [4x - 12y + 17 = 0]$ M1 A1
- (b) SP: $\frac{1}{2}x^{-\frac{1}{2}} - 4e^{1-4x} = 0, \quad \frac{1}{2\sqrt{x}} = 4e^{1-4x}$
 $\frac{1}{8\sqrt{x}} = e^{1-4x}$ M1
 $8\sqrt{x} = e^{4x-1}$
 $4x - 1 = \ln 8\sqrt{x}$ M1
 $x = \frac{1}{4}(1 + \ln 8\sqrt{x})$ A1
- (c) $x_1 = 0.7699, x_2 = 0.7372, x_3 = 0.7317, x_4 = 0.7308 = 0.731 \text{ (3dp)}$ M1 A2
- (d) let $f(x) = \frac{1}{2}x^{-\frac{1}{2}} - 4e^{1-4x}$
 $f(0.7305) = -0.00025, \quad f(0.7315) = 0.0017$ M1
sign change, $f(x)$ continuous \therefore root A1
- (e) $x_1 = 6.304, x_2 = 1.683 \times 10^{19}$
diverges rapidly away from root B2 (14)

Total (75)

Performance Record – C3 Paper C

Question no.	1	2	3	4	5	6	7	8	Total
Topic(s)	rational expressions	trigonometry	exponentials and logarithms	differentiation	functions	functions	trigonometry	differentiation, numerical methods	
Marks	6	7	8	8	9	10	13	14	75
Student									