

GCE Examinations
Advanced Subsidiary / Advanced Level
Statistics
Module S1

Paper F

MARKING GUIDE

This guide is intended to be as helpful as possible to teachers by providing concise solutions and indicating how marks should be awarded. There are obviously alternative methods that would also gain full marks.

Method marks (M) are awarded for knowing and using a method.

Accuracy marks (A) can only be awarded when a correct method has been used.

(B) marks are independent of method marks.



Written by Shaun Armstrong & Chris Huffer


© *Solomon Press*

These sheets may be copied for use solely by the purchaser's institute.


S1 Paper F – Marking Guide

1.	(a)	mean = $\frac{1145.3}{15} = 76.4$ kg	M1 A1	
		variance = $\frac{88042.14}{15} - 76.353^2 = 39.6$ kg ²	M2 A1	
	(b)	mean lower as replacement weighs less	B2	
		variance higher as replacement's weight further from mean	B2	(9)
<hr/>				
2.	(a)	$a + b + \frac{1}{4} + 2a + \frac{1}{8} = 1$	M1	
		$3a + b = \frac{5}{8}; b = \frac{5}{8} - 3a$	M1 A1	
	(b)	$\sum xP(x) = a + 2b + \frac{3}{4} + 8a + \frac{5}{8}$	M1	
		$= 9a + 2(\frac{5}{8} - 3a) + \frac{11}{8} = 3a + \frac{21}{8}$	M1 A1	
	(c)	$3a + \frac{21}{8} = \frac{45}{16}$	M1	
		$3a = \frac{45}{16} - \frac{21}{8} = \frac{3}{16}$	M1	
		$a = \frac{1}{16}, b = \frac{7}{16}$	A2	(10)
<hr/>				
3.	(a)	$P(Z < \frac{25-21.5}{2.2}) = P(Z < 1.59) = 0.9441$	M2 A1	
	(b)	$P(Z > \frac{19-21.5}{2.2}) = P(Z > -1.14) = 0.8729 \therefore 87.3\%$	M2 A1	
	(c)	$P(Z < \frac{20-21.5}{2.2}) = P(Z < -0.68) = 0.2483$	M1 A1	
		$P(2 \text{ of } 3 < 20) = 3 \times 0.2483^2 \times 0.7517 = 0.139$	M2 A1	(11)
<hr/>				
4.	(a)	$0.76 = 0.5 + 0.42 - P(A \cap B)$	M1	
		$P(A \cap B) = 0.92 - 0.76 = 0.16$	M1 A1	
	(b)	$(1 - 0.5) + 0.16 = 0.66$	M2 A1	
	(c)	$= \frac{P(B \cap A')}{P(A')} = \frac{0.42 - 0.16}{1 - 0.5} = 0.52$	M2 A1	
	(d)	$P(A) \times P(B) = 0.5 \times 0.42 = 0.21$	M1 A1	
		$\neq P(A \cap B) \therefore$ not independent	A1	(12)

5. (a) $n = 31$, median = 29
 $Q_1 = 23$
 $Q_3 = 34$
 $IQR = Q_3 - Q_1 = 34 - 23 = 11$ A1
A1
A1
M1 A1
- (b) $Q_2 - Q_1 = 6$; $Q_3 - Q_2 = 5$
 $\therefore Q_2 - Q_1 > Q_3 - Q_2 \therefore$ slight +ve skew M1
M1 A1
- (c) e.g. recommend mean and std. dev. as they take account of all values and there is little skew / few extreme values B2
- (d) $Q_1 - 2s = 2.4$; $Q_3 + 2s = 54.6 \therefore$ outliers are 0, 2 M1 A1



B4 (16)



B4

- (b) $S_{np} = 873 - \frac{21 \times 240.1}{6} = 32.65$ M1
 $S_{nn} = 91 - \frac{21^2}{6} = 17.5$ M1
 $b = \frac{32.65}{17.5} = 1.8657$ M1 A1
 $a = \frac{240.1}{6} - 1.8657 \times \frac{21}{6} = 33.4867$ M1 A1
 $p = 33.5 + 1.87n$ A1
line on graph above B2

- (c) $S_{pp} = 9675.41 - \frac{240.1^2}{6} = 67.4083$ M1
 $r = \frac{32.65}{\sqrt{17.5 \times 67.4083}} = 0.9506$ M1 A1
 r strongly +ve supporting linear model B1 (17)

Total (75)

