

GCE Examinations

Advanced Subsidiary / Advanced Level

Mechanics
Module M1

Paper J

MARKING GUIDE

This guide is intended to be as helpful as possible to teachers by providing concise solutions and indicating how marks should be awarded. There are obviously alternative methods that would also gain full marks.

Method marks (M) are awarded for knowing and using a method.

Accuracy marks (A) can only be awarded when a correct method has been used.

(B) marks are independent of method marks.




Written by Shaun Armstrong & Chris Huffer

© *Solomon Press*

These sheets may be copied for use solely by the purchaser's institute.


M1 Paper J – Marking Guide


1. (a) speed = 17 = mag. of vel. = $\sqrt{8^2 + \lambda^2}$ M1
 $\lambda^2 = 289 - 64 = 225$; $\lambda > 0$ so $\lambda = 15$ M1 A1
- (b) $\mathbf{a} = \frac{\Delta \mathbf{v}}{t} = \frac{1}{5} [(3\mathbf{i} + 5\mathbf{j}) - (8\mathbf{i} + 15\mathbf{j})] = -\mathbf{i} - 2\mathbf{j}$ M2 A1
 $\mathbf{F} = m\mathbf{a} = 2(-\mathbf{i} - 2\mathbf{j}) = -2(\mathbf{i} + 2\mathbf{j})$ so $\mu = -2$ M1 A1 **(8)**

2. (a) 
- resolve \uparrow : $R - P\cos 60^\circ - 0.5g = 0 \therefore R = 0.5g + P\cos 60^\circ$ M1 A1
 resolve \rightarrow : $P\sin 60^\circ - F = 0$ M1
 $F = \mu R = \frac{1}{\sqrt{3}}(0.5g + 0.5P)$ M1 A1
 sub. in giving $\frac{\sqrt{3}}{2}P - \frac{1}{\sqrt{3}}(0.5g + 0.5P) = 0$ M1
 $3P - P - g = 0 \therefore 2P = g$ so $P = \frac{g}{2}$ A1
- (b) brush is moved slowly so very little air resistance B1 **(8)**

3. (a) cons. of mom: $1500(2) + 0 = (1500 + 750)V$ M1
 $3000 = 2250V \therefore V = \frac{4}{3}$ M1 A1
- (b) impulse = Δ mom = $750(\frac{4}{3} - 0) = 1000$ Ns M1 A1
- (c) car has $(27 + 9)$ m in which to stop and travels 18 m in first second M1
 must stop from 18 ms^{-1} in 18 m A1
 $u = 18, s = 18, v = 0, a = -f$ M1
 $v^2 = u^2 + 2as$, so $0 = 324 - 36f$ M1
 $f = 9$ so to stop before hitting other car, $f > 9$ A1 **(10)**

4. (a) eqn. of motion for A: $T - 6g = 6a$ (1) M1
 eqn. of motion for B & C: $8g - T = 8a$ (2) M1
 (1) + (2) gives $2g = 14a$ i.e. $a = \frac{g}{7} \text{ ms}^{-2}$ M1 A1
- (b) sub. a into (1) to get $T = 6a + 6g = \frac{6g}{7} + 6g$ M1
 force on pulley = $2T = \frac{96g}{7}$ M1 A1
- (c) resolve \downarrow for C: $3g - R = 3 \times \frac{g}{7}$ M1
 $R = 3g - \frac{3g}{7} = \frac{18g}{7}$ M1 A1 **(10)**

5. (a) For Q : $a = \frac{\Delta v}{t} = \frac{6-0}{6} = 1$ M1
 $u = 0, v = 4$, use $v = u + at$: $4 = 0 + 1t$ i.e. $t = 4$ seconds M1 A1
- (b)  B3
- (c) Q will catch P when area under Q graph = area under P graph
 $\therefore \frac{1}{2} (6)(6) + 6(t-6) = 4t$ M1
 i.e. $18 + 6t - 36 = 4t \therefore 2t = 18 \therefore t = 9$ M1 A1
 after 9 seconds, P has travelled $4 \times 9 = 36$ cm,
 $\therefore Q$ reaches top first if $x > 36$ M1 A1 (11)

6. (a) as rock moves further from A , tension at A decreases linearly and is a minimum when rock reaches B .
-  B3
- (b) max. tension when rock at one end (A , say) B1
 moments about B : $50g(4) + Mg(8) - T_A(8) = 0$ M1
 $8Mg = 8T_A - 200g \therefore Mg = T_A - 25g$ M1
 given $T_A \leq 40g$; $Mg \leq 40g - 25g (= 15g)$ M1
 i.e. $M \leq 15$ A1
- (c) assume rock placed as close to A as poss. so that $T_A = 40g$
 resolve (\uparrow): $T_A + T_B = 50g + 20g = 70g \therefore T_B = 30g$ M1 A1
 moments about centre of plank: $T_A(4) - T_B(4) - 20g(d) = 0$ M1
 $160g - 120g - 20gd = 0 \therefore d = 2$ M1 A1
 rock can be 2 m either side of centre i.e. 4 m out of 8 m = $\frac{1}{2}$ plank A1 (14)

7. (a) cargo ship travels $(9t\mathbf{i} - 6t\mathbf{j})$ km in t hours
 posⁿ vector after t hours is $[(7\mathbf{i} + 56\mathbf{j}) + (9t\mathbf{i} - 6t\mathbf{j})]$ km M1
 $= [(7 + 9t)\mathbf{i} + (56 - 6t)\mathbf{j}]$ km A1
 posⁿ vector of ferry after t hours is $(12t\mathbf{i} + 18t\mathbf{j})$ km A1
- (b) they will collide if coeffs. of \mathbf{i} and \mathbf{j} in posⁿ vectors are equal B1
 $7 + 9t = 12t$ and $56 - 6t = 18t$ are both satisfied when $t = \frac{7}{3}$ M1 A1
 collision after $\frac{7}{3}$ hrs or 2 hrs 20 mins i.e. at 8:20 a.m. A1
 posⁿ vector = $12(\frac{7}{3})\mathbf{i} + 18(\frac{7}{3})\mathbf{j} = (28\mathbf{i} + 42\mathbf{j})$ M1 A1
- (c) at 8 a.m. ferry at $(24\mathbf{i} + 36\mathbf{j})$
 $\frac{1}{3}$ hr at $21\mathbf{i} + 6\mathbf{j} = 7\mathbf{i} + 2\mathbf{j}$ so at 8:20 a.m. ferry is at $31\mathbf{i} + 38\mathbf{j}$ M2 A1
 at 8:20 a.m cargo ship is at $(28\mathbf{i} + 42\mathbf{j})$
 dist. between = $\sqrt{(3^2 + 4^2)} = 5$ km M1 A1 (14)

Total (75)

Performance Record – M1 Paper J

Question no.	1	2	3	4	5	6	7	Total
Topic(s)	i, j, $F = ma$	statics, friction	cons. of mom., impulse, uniform accel.	connected bodies	speed - time graph, uniform accel.	moments	rel. posn. i, j	
Marks	8	8	10	10	11	14	14	75
Student								