

GCE Examinations  
Advanced Subsidiary

# Core Mathematics C1

Paper L

## MARKING GUIDE

This guide is intended to be as helpful as possible to teachers by providing concise solutions and indicating how marks could be awarded. There are obviously alternative methods that would also gain full marks.

Method marks (M) are awarded for knowing and using a method.

Accuracy marks (A) can only be awarded when a correct method has been used.

(B) marks are independent of method marks.




*Written by Shaun Armstrong*




© *Solomon Press*

*These sheets may be copied for use solely by the purchaser's institute.*

## C1 Paper L – Marking Guide

1.	$= \sqrt{49} + (\sqrt[3]{8})^2 = 7 + 2^2$ $= 11$	B1 M1 A1	(3)
2.	(a) $u_4 = \frac{5+1}{3} = 2$  (b) $5 = \frac{u_2+1}{3}, u_2 = 14$ $14 = \frac{u_1+1}{3}, u_1 = 41$	B1  M1 A1  A1	(4)
3.	(a) $b^2 - 4ac = 12^2 - (4 \times 4 \times 9) = 0$ $\therefore 1$ real root  (b) $4x^2 + 12x + 9 = 8$ $4x^2 + 12x + 1 = 0$ $x = \frac{-12 \pm \sqrt{144 - 16}}{8}$ $= \frac{-12 \pm 8\sqrt{2}}{8}$ $= -\frac{3}{2} \pm \sqrt{2}$	M1 A1   M1  M1  A2	(6)
4.	(a) $5x > 15$ $x > 3$  (b) $(x+2)(x-8) < 0$ $-2 < x < 8$  (c) $3 < x < 8$	M1 A1    M1 M1 A1  B1	(6)
5.	(a) $(2 - \sqrt{x})^2 = 0$ $\sqrt{x} = 2$ $x = 4$  (b) $= (2 - \sqrt{3})^2 = 4 - 4\sqrt{3} + 3 = 7 - 4\sqrt{3}$  (c) $= \int (2 - \sqrt{x})^2 dx$ $= \int (4 - 4\sqrt{x} + x) dx$ $= 4x - \frac{8}{3}x^{\frac{3}{2}} + \frac{1}{2}x^2 + c$	M1 A1   M1 A1    B1  M1 A2	(8)
6.	(a) $\text{grad} = \frac{-4-6}{1-(-3)} = -\frac{5}{2}$ $\therefore y - 6 = -\frac{5}{2}(x + 3)$ $2y - 12 = -5x - 15$ $5x + 2y + 3 = 0$  (b) $m: y = -\frac{2}{k}x - \frac{7}{k} \therefore \text{grad} = -\frac{2}{k}$  $l$ and $m$ perp. $\therefore -\frac{5}{2} \times -\frac{2}{k} = -1$ $k = -5$	M1 A1  M1  A1    M1 A1  M1  A1	(8)

7. (a)  $f(x) = \int (5 + \frac{4}{x^2}) dx$   
 $f(x) = 5x - 4x^{-1} + c$  M1 A2
- (b)  $f(1) = 5 - 4 + c = 1 + c$  M1  
 $f(2) = 10 - 2 + c = 8 + c$   
 $f(2) = 2f(1) \therefore 8 + c = 2(1 + c)$  M1  
 $c = 6$  A1  
 $f(x) = 5x - 4x^{-1} + 6$   
 $f(4) = 20 - 1 + 6 = 25$  M1 A1 (8)

8. (a) LHS =  $(x + 1)(x^2 - 7x + 12)$  M1  
 $= x^3 - 7x^2 + 12x + x^2 - 7x + 12$  M1  
 $= x^3 - 6x^2 + 5x + 12 = \text{RHS}$  A1
- (b)  B3
- (c) (i)  (ii)  B2 B2
- (10)

9. (a) (i)  $= (t^2 - 5) - (t - 1) = t^2 - t - 4$  M1 A1  
(ii)  $= (t^2 - 5) + (t^2 - t - 4) = 2t^2 - t - 9$  M1 A1
- (b)  $2t^2 - t - 9 = 19$   
 $2t^2 - t - 28 = 0$   
 $(2t + 7)(t - 4) = 0$  M1  
 $t > 0 \therefore t = 4$  A1
- (c)  $a = 4 - 1 = 3, d = 16 - 4 - 4 = 8$  B1  
 $u_{10} = 3 + (9 \times 8) = 3 + 72 = 75$  M1 A1
- (d)  $= \frac{40}{2} [6 + (39 \times 8)] = 20 \times 318 = 6360$  M1 A1 (11)

10. (a) A (0, 2) B1  
 $\frac{dy}{dx} = 3 - 2x$  M1 A1  
grad = 3 M1  
 $\therefore y = 3x + 2$  A1
- (b) grad of  $m = 3$   
grad of curve at B =  $\frac{-1}{3} = -\frac{1}{3}$  M1 A1  
at B:  $3 - 2x = -\frac{1}{3}$   
 $x = \frac{5}{3}$  M1 A1  
 $y = 2 + 3(\frac{5}{3}) - (\frac{5}{3})^2 = 4\frac{2}{9} \therefore B (1\frac{2}{3}, 4\frac{2}{9})$  M1 A1 (11)

Total (75)

### Performance Record – C1 Paper L

Question no.	1	2	3	4	5	6	7	8	9	10	Total
Topic(s)	indices	recur. relation	quad. formula	inequals	surds, integr.	straight lines	integr.	curve sketch, transform.	AP	diff., tangent	
Marks	3	4	6	6	8	8	8	10	11	11	75
Student											