

Mark Scheme (Results)

Summer 2013

International GCSE
Chemistry (4CH0) Paper 1CR

Science Double Award (4SC0)
Paper 1CR

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications come from Pearson, the world's leading learning company. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk for our BTEC qualifications.

Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

If you have any subject specific questions about this specification that require the help of a subject specialist, you can speak directly to the subject team at Pearson.

Their contact details can be found on this link: www.edexcel.com/teachingservices.

You can also use our online Ask the Expert service at www.edexcel.com/ask. You will need an Edexcel username and password to access this service.

Pearson: helping people progress, everywhere

Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

Summer 2013

Publications Code UG035595

All the material in this publication is copyright

© Pearson Education Ltd 2013

Question number	Answer	Accept	Reject	Marks
1 (a)	can <u>easily/quickly</u> identify each gas OR <u>less likely</u> to make a mistake in identification IGNORE just to identify the gas			1
(b) (i)	argon/Ar <u>and</u> helium/He			1
(ii)	oxygen/O ₂ IGNORE O			1
(iii)	air/it is a mixture (of gases) OR air/it is not a single substance IGNORE mixture of elements			1
(iv)	not flammable/not explosive/does not burn			
(c) (i)	hydrogen/H ₂ IGNORE H	air		1
(ii)	carbon dioxide/CO ₂			1
(iii)	carbon dioxide/CO ₂			1
			Total	8

Question number	Answer	Accept	Reject	Marks
2 (a)	D			1
(b)	<p>M1 before heating – colourless (solution/liquid) IGNORE clear/transparent/looks like water</p> <p>M2 after heating – milky/chalky/cloudy/white (precipitate)/turbid</p> <p>IGNORE references to goes clear OWTTE</p>	no colour	white solution/liquid any colour other than white	1 1
(c)	<p>M1 (sulfur dioxide/it) dissolves in/reacts with (rain) water</p> <p>M2 to form an acidic solution/an acid/sulfurous acid /acid rain IGNORE references to any other products whether correct or not</p> <p>M3 which reacts with/corrodes the marble/calcium carbonate</p> <p>IGNORE erodes / weathers / melts / eats into</p>	$\text{SO}_2 + \text{H}_2\text{O} \rightarrow \text{H}_2\text{SO}_3$ OR $\text{SO}_2 + \text{H}_2\text{O} + \frac{1}{2}\text{O}_2 \rightarrow \text{H}_2\text{SO}_4$ for both M1 and M2 sulfuric acid <u>chemical</u> weathering dissolves correct equation for reaction with either sulfurous or sulfuric acid SO_2 reacts with marble for M3 only		1 1 1
			Total	6

Question number	Answer	Accept	Reject	Marks																				
3 (a)	<table border="1"> <thead> <tr> <th>Name of barium salt</th> <th>Formula of barium salt</th> <th>Solubility in water</th> <th>Poisonous</th> </tr> </thead> <tbody> <tr> <td>barium chloride</td> <td>BaCl₂</td> <td></td> <td></td> </tr> <tr> <td>barium nitrate</td> <td></td> <td></td> <td></td> </tr> <tr> <td>barium carbonate</td> <td>BaCO₃</td> <td></td> <td></td> </tr> <tr> <td>barium sulfate</td> <td></td> <td></td> <td></td> </tr> </tbody> </table>	Name of barium salt	Formula of barium salt	Solubility in water	Poisonous	barium chloride	BaCl ₂			barium nitrate				barium carbonate	BaCO ₃			barium sulfate						1
	Name of barium salt	Formula of barium salt	Solubility in water	Poisonous																				
	barium chloride	BaCl ₂																						
	barium nitrate																							
	barium carbonate	BaCO ₃																						
barium sulfate																								
				1																				
(b)	<p>M1 (it forms) barium chloride/BaCl₂/a soluble (barium) salt</p> <p>M2 by reaction/with hydrochloric acid/stomach acid</p>	<p>by neutralisation</p> <p>word or chemical equation for 2 marks (equation can be unbalanced)</p>	<p>any suggestion that barium chloride is reacting</p>	1																				
(c)	barium sulfate/BaSO ₄			1																				

Question number	Answer	Accept	Reject	Marks
3 (d)	<p>M1 barium sulfate is formed</p> <p>M2 which is not poisonous/not toxic/harmless IGNORE references to magnesium hydroxide not poisonous</p> <p>M2 dep on M1</p> <p>M3 barium hydroxide + magnesium sulfate → barium sulfate + magnesium hydroxide</p> <p>OR</p> <p>barium ions + sulfate ions → barium sulfate</p>	<p>'products', provided shown correctly in word equation</p> <p>is insoluble</p> <p>$\text{Ba}(\text{OH})_2 + \text{MgSO}_4 \rightarrow \text{BaSO}_4 + \text{Mg}(\text{OH})_2$</p> <p>OR</p> <p>$\text{Ba}^{2+} + \text{SO}_4^{2-} \rightarrow \text{BaSO}_4$</p>		<p>1</p> <p>1</p> <p>1</p>
(e) (i)	<p>M1 water – (reacts) <u>very/extremely</u> quickly/more quickly <u>than strontium/quickest</u> IGNORE rapidly/vigorously</p> <p>M2 air – (reacts) <u>very/extremely</u> quickly/more quickly <u>than strontium/quickest</u> (without heating) IGNORE rapidly/vigorously</p> <p>(ii)</p> <p>in/under any one of the following: (paraffin/mineral) oil/petroleum (oil)/(liquid) paraffin</p> <p>(iii) IGNORE in an air tight container</p> <p>reactivity <u>increases</u> as atomic number <u>increases</u></p>	<p>explosively/violently</p> <p>explosively/violently</p> <p>in a vacuum</p> <p>reactivity increases with atomic number/down the group OWTTE reverse argument</p>		<p>1</p> <p>1</p> <p>1</p> <p>1</p>

positive correlation

			Total	12

Question number	Answer	Accept	Reject	Marks
4 (a)	M1 (negative electrode) – graphite M2 (positive electrode) – graphite	carbon carbon		2
(b) (i)	it/aluminium oxide/alumina has a (very) high m.pt IGNORE high b.pt/references to strong bonding/bauxite has a high m.pt/lot of energy needed to melt it		aluminium has a high melting point	1
(ii)	aluminium oxide/alumina is dissolved in/mixed with (molten/liquid) cryolite IGNORE cryolite lowers the m.pt of aluminium oxide/alumina	added to Na_3AlF_6 for cryolite cryolite is used as the solvent (for aluminium oxide/alumina)	aluminium is dissolved in cryolite	1
(c)	M1 reduction M2 (it/aluminium ions/ Al^{3+}) gain of electron(s) IGNORE references to loss of oxygen M2 dep on M1	reacts with/combines with decrease in oxidation number/oxidation number changes from +3 to 0	redox for M1 only Al/aluminium gains electrons	1 1
(d)	M1 oxygen formed/produced (at the positive electrode/anode) IGNORE oxygen from the aluminium oxide M2 reacts with the carbon/the (positive) electrode M2 not dep on M1 , but must mention oxygen	oxygen from the electrolysis anode / graphite	any indication that the oxygen is from the air for M1 only cathode/negative electrode	1 1
(e)	Any two from: M1 malleable M2 low density	easy to shape/easy to bend/easy to extrude bend		2

	M3 does not react <u>with food/drink(s)</u> IGNORE light(er)/high strength to weight ratio/references to cost/lightweight/does not rust	non-toxic/does not corrode		
			Total	10

Question number	Answer	Accept	Reject	Marks
5 (a)	M1 (molecules/compounds/substances) with the same <u>molecular</u> formula/number of each type of atoms IGNORE chemical formula/same compound M2 (but) different structural formulae/different displayed formulae/different structures	hydrocarbons atoms arranged differently	elements/atoms general formula/empirical formula for M1 only	1 1
(b)	D			1
(c) (i)	M1 C _n H _{2n}	letters other than n, e.g. x	C _n +H _{2n}	1
(ii)	M1 double bond between two left hand end carbon atoms M2 single bond between each pair of rest of carbon atoms Penalise max 1 mark for any extra bond shown			1 1
(d)	M1 addition M2 orange M3 colourless IGNORE clear/transparent/looks like water	additional yellow/brown	red, either on its own or in combination with any other colour	1 1 1
(e)	M1 saturated – <u>all</u> (carbon to carbon) bonds are single /contains <u>only</u> (carbon to carbon)	does not contain any multiple/double bonds		1

	single bonds			1
	M2 unsaturated - contains (carbon to carbon) double/multiple bond(s)			
			Total	11

Question number	Answer	Accept	Reject	Marks
6 (a) (i)	7			1
(ii)	M1 solid M2 black	<u>very</u> dark grey		1 1
(iii)	M1 (formula) – HAt M2 (name) – hydrogen astatide	AtH astatine hydride	hydrogen astatine	1 1
(iv)	M1 – (astatine/it/At) is less reactive (than iodine, I) IGNORE astatine is unreactive M2 – elements get less reactive with <u>increasing</u> atomic number/as group is <u>descended</u> /the lower they are in the group	<u>iodine</u> is more reactive reverse argument Astatine (atom) has more (electron) shells/outer shell of astatine is further from nucleus so attracts an <u>electron</u> less readily	any references to astatide or iodide	1 1
(b) (i)	4 (1) (1) 2 (1)	multiples/halves		1
(ii)	(paper) turns white/bleaches IGNORE turns red	(litmus) turns colourless		1
(c) (i)	acid IGNORE hydrogen ions/names of acids	correct formula		1
(ii)	to displace (all of) the <u>bromine</u> / to react all of the <u>bromide</u> (ions)	bromine (an) <u>ions</u> for bromide to complete the reaction		1
(iii)	$\text{Br}_2 + \text{SO}_2 + 2\text{H}_2\text{O} \rightarrow 2\text{HBr} + \text{H}_2\text{SO}_4$ M1 all formulae correct M2 balanced	multiples and halves		2
(iv)	$2\text{HBr} + \text{Cl}_2 \rightarrow \text{Br}_2 + 2\text{HCl}$	multiples and halves		1

(d)	M1 colourless IGNORE clear/transparent/looks like water M2 brown (solution) / (dark) grey/black <u>solid/precipitate</u>	red- brown/orange/orange- brown	red on its own	1
			Total	16

Question number	Answer	Accept	Reject	Marks
7 (a)	M1 (reactants) s aq M2 (products) aq l g	capital letters		1 1
(b) (i)	to prevent acid escaping/spraying out/spitting out IGNORE to prevent water escaping	solution/liquid/HCl		1
(ii)	C			1
(c) (i)	M1 A M2 gas produced/collected more quickly / experiment over in shorter time / (gradient of) line steeper M2 dep on M1	reaction is faster		1 1 1
(ii)	M1 0.1(0) M2 volume of gas is half/40 ÷ 80 = ½ / 80 = 40 x 2 M2 dep on M1	Half the products are produced		1
(d) (i)	M1 & M2 - all points plotted to nearest gridline deduct 1 mark for each incorrect plot up to a max. of 2			2 1
(ii)	M3 suitable straight line of best fit (csq on plotted points) (must be drawn with the aid of a ruler). Line need not be extrapolated. M1 as concentration increases rate increases	(show a) <u>positive</u> correlation as one doubles the other doubles/directly proportional		1 1

(iii)	<p>M2 proportional / in proportion</p> <p>M1 more ions/particles (in a given volume) IGNORE more reactants</p> <p>M2 collide (successfully)</p> <p>M3 more per second/more frequently</p> <p>Must be reference to frequency or number of collisions per unit time IGNORE greater chance of collision</p>	for 2 marks	<p>molecules/atoms</p> <p>any reference to greater energy</p>	<p>1</p> <p>1</p> <p>1</p>
			Total	16

Question number	Answer	Accept	Reject	Marks
8 (a) (i)	Impurities/chemicals/substances may affect the colour/flame IGNORE affect the result/test			1
(ii)	colour can (easily) be seen (in a non-luminous flame) IGNORE references to temperature	a luminous flame may mask the colour		1
(iii)	yellow/orange/gold(en)	any combination of the acceptable colours, e.g. golden-yellow		1
(b) (i)	Li ⁺ and Ca ²⁺	lithium and calcium/Li and Ca	Ca ⁺ etc	1
(ii)	M1 – ammonia/NH ₃ M2 – (water is needed) to form hydroxide ions/OH ⁻	to form an alkali/an alkaline solution/ammonium hydroxide		1
(iii)	M1 – iron(III)/Fe ³⁺ M2 – ammonium/NH ₄ ⁺ If both names and formulae given both must be correct	to dissolve the ammonia ammonia needs to be aqueous ferric	any other oxidation states/ferrous ammonia	1 1
			Total	8

Question number	Answer	Accept	Reject	Marks
9 (a)	(i) measuring cylinder			1
	(ii) M1 44	answers in other correct units, e.g. 0.044 dm ³ ml		1
	M2 cm ³			1
	(iii)	M1 $\frac{44 \times 0.01(0)}{1000}$		
M2 0.00044(0) Mark csq on answer to (a)(ii)		0.44 for 1 mark only correct answer with no working for 2 marks	0.0004	1
(b)	<u>zinc</u> because M1 1 mol zinc reacts with 2 mol HCl M2 only 0.005 mol of zinc are needed M1 is standalone M2 is dep on zinc given as being in excess			1 1
(c)	(i) (rate) increases/faster reaction	less time for reaction to take place	faster time	1
	(ii) no effect/same volume (of hydrogen) produced	none/no change		1
			Total	9

Question number	Answer	Accept	Reject	Marks
10 (a)	<p>any two from:</p> <ul style="list-style-type: none"> • forward and backward reactions (still) occurring • concentrations/amounts of reactants/products/components remain constant • rate of forward reaction = rate of reverse reaction <p>IGNORE concentrations/amounts of reactants and products are the same IGNORE reaction is reversible/goes both ways, OWTTE IGNORE references to le Chatelier</p>	<p>both reactions (still) occurring</p> <p>stay the same in place of remain constant</p>		2
(b) (i)	<p>M1 – (increase in temperature) decrease(s)</p> <p>M2 – (increase in pressure) increase(s)</p>	<p>less/<u>low</u>er(s)/drop(s)/gets smaller</p>		1
(ii)	<p>M1 – (forward) reaction is exothermic/gives out heat</p> <p>OR</p> <p><u>reverse</u> reaction is endothermic/takes in heat</p> <p>M2 – fewer (gas) molecules/particles on right hand side</p> <p>OR fewer moles (of gas) on right hand side</p> <p>IGNORE references to volumes IGNORE references to le Chatelier's principle IGNORE references to reverse reaction lowers the temperature IGNORE references to forward reaction reduces the pressure</p>	<p>more/raise(s)/high<u>er</u>/gets bigger</p> <p>reverse argument shifts to side with fewer (gas) molecules/fewer moles (of gas)</p>	atoms	1

10 (c)	(i)	(the position of) equilibrium is not established/reached		1
	(ii)	M1 – (the mixture of gases is) cooled	temperature is decreased	1
	(iii)	M2 – ammonia liquefies / condenses M3 recycled / <u>re</u> used / recirculated	put (back) into the reaction chamber used <u>again</u> (in the process)	1
(d)	<u>heat(ing)</u> / <u>energy</u> costs would be higher	yield (of ammonia) would decrease		1
(e)	(i)	M1 $M_r(\text{N}_2) = 28$ M2 $112\,000 \div 28 (= 4\,000) / 112\,000 \div$ M1 M3 $8\,000 / \text{M2} \times 2$	28 anywhere in the calculation $112 \div 28) \times 2 = 8$ for 2 marks $(112\,000 \div 14) \times 2 = 16\,000$ for 2 marks Correct final answer without working for 3 marks	1 1 1
	(ii)	1 200 / 15% of M3		1
			Total	15

Question number	Answer	Accept	Reject	Marks
11 (a)	(produces) <u>most</u> heat/energy <u>per gram</u> / <u>per unit mass</u>	<u>highest</u> temperature rise <u>per gram</u> / <u>per unit mass</u> <u>most</u> energy for <u>smallest</u> number of <u>grams</u> / <u>mass</u>	per amount	1
(b)	(produces) <u>most</u> heat/energy <u>per mole</u> / <u>per molecule</u> / <u>per amount</u>	<u>highest</u> temperature rise <u>per mole</u> / <u>per molecule</u> <u>most</u> energy for <u>smallest</u> number of <u>moles</u> / <u>molecules</u> / <u>amount</u>		1
(c)	Any two from: <ul style="list-style-type: none"> • heat/energy losses (e.g. by convection, by conduction, to air, to surroundings) • incomplete combustion • evaporation of water • copper / can / beaker / thermometer /apparatus absorbs heat • flame moves around because of draughts 	<ul style="list-style-type: none"> • non-standard conditions 		2
(d) (i)	A			1
(ii)	B			1
(e)	<p>M1 breaking bonds is endothermic / takes in heat/energy</p> <p>M2 making bonds is exothermic / gives out heat/energy</p> <p>M3 more heat/energy given out than taken in</p>	<p>more energy is given out when bonds are made than is taken in when bonds are broken for 3 marks</p> <p>more energy is given out when bonds are made than when bonds are broken for 1 mark</p>		1 1 1

	IGNORE references to numbers/strengths of bonds			
			Total	9

Further copies of this publication are available from
Edexcel Publications, Adamsway, Mansfield, Notts, NG18 4FN

Telephone 01623 467467

Fax 01623 450481

Email publication.orders@edexcel.com

Order Code xxxxxxxx Summer 2013

For more information on Edexcel qualifications, please visit our website
www.edexcel.com

Pearson Education Limited. Registered company number 872828
with its registered office at Edinburgh Gate, Harlow, Essex CM20 2JE

Ofqual
■■■■■■■■■■



Llywodraeth Cynulliad Cymru
Welsh Assembly Government

