

Mark Scheme (Results)

June 2011

GCE Physics (6PH04) Paper 01
Physics on the Move

Edexcel is one of the leading examining and awarding bodies in the UK and throughout the world. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers.

Through a network of UK and overseas offices, Edexcel's centres receive the support they need to help them deliver their education and training programmes to learners.

For further information, please call our GCE line on 0844 576 0025, our GCSE team on 0844 576 0027, or visit our website at www.edexcel.com.

If you have any subject specific questions about the content of this Mark Scheme that require the help of a subject specialist, you may find our **Ask The Expert** email service helpful.

Ask The Expert can be accessed online at the following link:
<http://www.edexcel.com/Aboutus/contact-us/>

Alternatively, you can contact our GCE Science Advisor directly by sending an email to ScienceSubjectAdvisor@EdexcelExperts.co.uk.

You can also telephone 0844 576 0037 to speak to a member of our subject advisor team.

June 2011

Publications Code UA028549

All the material in this publication is copyright

© Edexcel Ltd 2011

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

Physics Specific Marking Guidance

Underlying principle

The mark scheme will clearly indicate the concept that is being rewarded, backed up by examples. It is not a set of model answers.

For example:

Horizontal force of hinge on table top

66.3 (N) or 66 (N) **and** correct indication of direction [no ue]

[Some examples of direction: acting from right (to left) / to the left / West / opposite direction to horizontal. May show direction by arrow. Do not accept a minus sign in front of number as direction.]

This has a clear statement of the principle for awarding the mark, supported by some examples illustrating acceptable boundaries.

Mark scheme format

- Bold lower case will be used for emphasis.
- Round brackets () indicate words that are not essential e.g. "(hence) distance is increased".
- Square brackets [] indicate advice to examiners or examples e.g. [Do not accept gravity] [ecf].

Unit error penalties

- A separate mark is not usually given for a unit but a missing or incorrect unit will normally cause the final calculation mark to be lost.
- Incorrect use of case e.g. 'Watt' or 'w' will not be penalised.
- There will be no unit penalty applied in 'show that' questions or in any other question where the units to be used have been given.
- The same missing or incorrect unit will not be penalised more than once within one question but may be penalised again in another question.
- Occasionally, it may be decided not to penalise a missing or incorrect unit e.g. the candidate may be calculating the gradient of a graph, resulting in a unit that is not one that should be known and is complex.
- The mark scheme will indicate if no unit error penalty is to be applied by means of [no ue].

Significant figures

- Use of an inappropriate number of significant figures in the theory papers will normally only be penalised in 'show that' questions where use of too few significant figures has resulted in the candidate not demonstrating the validity of the given answer.
- Use of an inappropriate number of significant figures will normally be penalised in the practical examinations or coursework.
- Using $g = 10 \text{ m s}^{-2}$ **will** be penalised.

Calculations

- Bald (i.e. no working shown) correct answers score full marks unless in a 'show that' question.
- Rounding errors will not be penalised.
- If a 'show that' question is worth 2 marks then both marks will be available for a reverse working; if it is worth 3 marks then only 2 will be available.
- use of the formula means that the candidate demonstrates substitution of physically correct values, although there may be conversion errors e.g. power of 10 error.
- recall of the correct formula will be awarded when the formula is seen or implied by substitution.
- The mark scheme will show a correctly worked answer for illustration only.

Quality of Written Communication

- Indicated by QoWC in mark scheme. QWC – Work must be clear and organised in a logical manner using technical wording where appropriate.
- Usually it is part of a max mark, the final mark not being awarded unless the QoWC condition has been satisfied.

Question Number	Answer	Mark
1	D	1
2	A	1
3	C	1
4	C	1
5	D	1
6	B	1
7	B	1
8	B	1
9	A	1
10	C	1

Question Number	Answer	Mark
11(a)	<p>Identifying the equations $E_k = p^2 / 2m$ and $\lambda = h/p$ OR $\lambda = h/p, p = mv$ and $E_k = \frac{1}{2} mv^2$ (1)</p> <p>Any combination or rearrangement (conditional mark) (1)</p> <p>(do not give 2nd mark just for quoting equation given in question)</p> <p>(Do not credit a reverse argument i.e. starting with the given equation.)</p> <p><u>Example of derivation</u> $p = \sqrt{2mE_k}$ $\lambda = h/\sqrt{2mE_k}$</p>	2
11(b)	<p>Correct sub of h^2 and m (1) Use of $E_k = eV$ (1) $\lambda = 2.5 \times 10^{-11} \text{ m}$ (1)</p> <p>OR Use of $E_k = \frac{1}{2} mv^2$ (to find $v = 3.0 \times 10^7 \text{ (m s}^{-1}\text{)})$ (1) Use of $\lambda = h/p$ with correct substitution for h and m (1) $\lambda = 2.5 \times 10^{-11} \text{ m}$ (1)</p> <p><u>Example of calculation</u></p> $\lambda = \frac{(6.63 \times 10^{-34} \text{ J s})^2}{\sqrt{2(9.11 \times 10^{-31} \text{ kg})(2500 \text{ V})(1.6 \times 10^{-19} \text{ C})}}$ <p>$\lambda = 2.46 \times 10^{-11} \text{ m}$</p> <p>OR</p> $v = \sqrt{\frac{2(2500 \text{ V})(1.6 \times 10^{-19} \text{ C})}{9.1 \times 10^{-31} \text{ kg}}} = 3.0 \times 10^7$ <p>$\lambda = 6.63 \times 10^{-34} \text{ J s} / (9.1 \times 10^{-31} \text{ kg})(3.0 \times 10^7 \text{ m s}^{-1})$</p>	3
	Total for question 11	5

Question Number	Answer	Mark
12(a)	Third column completed 4.04 and 3.50 Points plotted correctly and straight line drawn (ecf error in calculation for points plotted)	(1) (1) 2
12(b)	Any evidence of gradient (look at graph) Value between 0.061 and 0.066 (cm^{-1}) (ignore – sign) Or value between 6.1 and 6.6 (m^{-1})	(1) (1) 2
Total for question 12		4


Question Number	Answer	Mark
13(a)	Method marks only Use of $Q=CV$ with $V=16\text{ V}$ Max value of $C = 12000\ (\mu\text{F})$ μF means 10^{-6} conversion of μF to F	(1) (1) (1) 3
<u>Example of calculation</u> $C_{\text{max}} = 1.20 \times 10000 = 12000\text{ F}$ $C_{\text{max}} = 12000\text{ F} \times 16\text{ V}$ $Q_{\text{max}} = 0.192\text{ C}$		
13(b)	Either use of $\frac{1}{2} QV$ or $\frac{1}{2} CV^2$ Energy = 1.5 J	(1) (1) 2
<u>Example of calculation</u> $W = \frac{1}{2} 0.192\text{ C} \times 16\text{ V}$ Energy = 1.54 J		
Total for question 13		5

Question Number	Answer	Mark
* 14	(QWC- Work must be clear and organised in a logical manner using technical wording where appropriate)	
Max 4		
<u>Uniform</u> electric field (between plates)		(1)
Force due to E or idea of attraction/repulsion		(1)
(Ball has an) <u>acceleration</u> (not an increasing velocity)		(1)
Which is constant/uniform (can be with reference to increasing velocity)		(1)
Vertical line/ + and – values shows change in direction		(1)
Inelastic collision/less energy after impact		(1)
Total for question 14		4

Question Number	Answer	Mark
15(a)	Use of $\frac{1}{2}mv^2$ and mgh (1) (do not credit use of $v^2 = 2as$ since a not constant, scores 0/2) velocity = $0.77 \text{ (m s}^{-1}\text{)}$ (1)	2
15(b)	Use of mv (1) Correct momentum conservation statement (1) Speed = 0.53 m s^{-1} (accept 0.56 m s^{-1} from use of show that value) (1) Assumption: no external forces/ no air resistance/ no force on pivot/negligible resistance (1) <u>Example of calculation</u> $320 \text{ g} \times 0.77 \text{ m s}^{-1} = (320 \text{ g} \times v) + (55 \text{ g} \times 1.4 \text{ m s}^{-1})$ Speed of bat = 0.53 m s^{-1}	4
15(c)	Use of $E_k = \frac{1}{2}mv^2$ (allow mass in g or kg) (1) Correct calculation of E_k before and after (95, 45, 54 to any power of ten (see below) (If 0.8 m s^{-1} and 0.56 m s^{-1} , values are 102, 50 and 54) (1) Elastic. (1)	3
15(d)	Max 2 x measured to nearest cm (1) uncertainty in x gives an uncertainty in GPE / speed (of ball) (1) difficulty of measuring a moving object (1) some energy to sound (1) comments on $E_{(k)}$ after > before (1) relates uncertainty to conclusion made in (c) (1)	2
Total for Question 15		11

Question Number	Answer	Mark
16(a)	Radial lines (at least 4) most touching nucleus but not going through it (straight by eye) (1) Equispaced (1) Arrow pointing away from circle (1)	3
16(b) (i)	$F = Q_1 Q_2 / 4\pi\epsilon r^2$ or $F = k Q_1 Q_2 / r^2$ (1) Charges are $79 \times 1.6 \times 10^{-19}$ and $2 \times 1.6 \times 10^{-19}$ (1) (values are 1.23×10^{-17} and 3.2×10^{-19})	2
16(b) (ii)	Marks can be scored for use of symbols, cell annotation or values $F = \Delta p / (\Delta)t$ (1) $(\Delta)v = (\Delta)p/m$ (1) $(D5) = D4 + \frac{(-)B5 \times C5}{6.64 \times 10^{-27}}$ (1) $(D5) = 1.24 \times 10^7 + \frac{(-)20.2 \times 1 \times 10^{-21}}{6.64 \times 10^{-27}}$ $F/m = (-) 3.04 \times 10^{27}$ OR (1) $a = F/m$ (1) $v = (u) + at$ (1) $(D5) = D4 + \frac{(-)B5 \times C5}{6.64 \times 10^{-27}}$ $(D5) = 1.24 \times 10^7 + \frac{(-)20.2 \times 1 \times 10^{-21}}{6.64 \times 10^{-27}}$ $F/m = (-) 3.04 \times 10^{27}$	3
16(b) (iii)	$s = \frac{1}{2} (u+v) t$ accept $s = vt$ (with either D5 or D6) Or $s = ut + \frac{1}{2} at^2$ (1) $(s) = \frac{1}{2} (D5 + D6) * C6$ or values or other correct equations (1)	2
16(b) (iv)	Value in range 2.00 - 2.49(x 10^{-14} m) (1)	1
*16(c)	(QWC- Work must be clear and organised in a logical manner using technical wording where appropriate.) Atom mainly <u>empty</u> space (1) Charge is concentrated in the centre/in a nucleus/nucleus is charged (1) Mass is concentrated (at the centre) Or Dense/massive nucleus (1)	3
Total for question 16		14

Question Number	Answer	Mark
17(a)	(Magnetic) force acts at right angles to ion motion/current Force is the centripetal force or causing centripetal acceleration or direction of acceleration/force is to centre (of circle)	(1) (1) 2
17(b)	See $F = BQv$ or $r = p/BQ$ $F = mv^2/r$ or $p = mv$ $f = v/2\pi r$ or $f = \omega/2\pi$ or $T = 2\pi r/v$ or $T = 2\pi/\omega$	(1) (1) (1) 3
17(c)(i)	Identifies positive (field) above and below (the ion) which repels the ion	(1) (1) 2
17(c)(ii)	$3 \times 32.0645 / 10 \times (10^6)$ $= 0.0000096(u)$	(1) (1) 2
17(c)(iii)	Convert MeV to J Convert J to kg Convert kg to u Mass loss = 0.0024(u) (and this is more than 0.00001u) <u>Example of calculation</u> mass loss = $2.2 \text{ MeV} \times 1.6 \times 10^{-13} \text{ J}$ J to kg $3.52 \times 10^{-13} / 9 \times 10^{16} \text{ kg}$ kg to u $3.91 \times 10^{-30} / 1.66 \times 10^{-27} \text{ u}$	(1) (1) (1) (1) 4
	Total for question 17	13

Question Number	Answer	Mark
18(a)	to keep the time spent in each tube the same Or so that frequency of <u>alternating</u> pd/voltage constant (do not accept reference to ac currents)	(1) 1
18(b) (i)	At top of Λ 	(1) 1
18(b) (ii)	No <u>track/trail</u> to this point (as no charge) Then two tracks (as two charged particles)	(1) (1) 2
18(b) (iii)	$D^0 \rightarrow k^+ + \pi^-$ Correct symbols, do not accept pi as a word Correct charge symbols as above, top right of each term. Wrong position or extra ones loses this mark (any extra particles or gamma 0/2)	(1) (1) 2
* 18(b) (iv)	(QWC- Work must be clear and organised in a logical manner using technical wording where appropriate.) Charge: +1 -1 = 0 (the positive sign must be used, 1 -1 = 0 is not enough) Or reference to positive and negative charges cancelling. Energy/mass/mass-energy: of D is equal to that of the new particles Momentum: Of D is equal to sum/total of momentum of new particles	(1) (1) (1) (1) (1) (1) 6
18(c) (i)	$c \bar{u}$	(1) 1
18(c) (ii)	$u\bar{d}$ or $u\bar{s}$ or $c\bar{d}$ or $c\bar{s}$ (allow words)	(1) 1
Total for question 18		14

Further copies of this publication are available from
Edexcel Publications, Adamsway, Mansfield, Notts, NG18 4FN

Telephone 01623 467467

Fax 01623 450481

Email publication.orders@edexcel.com

Order Code UA028549 June 2011

For more information on Edexcel qualifications, please visit
www.edexcel.com/quals

Pearson Education Limited. Registered company number 872828
with its registered office at Edinburgh Gate, Harlow, Essex CM20 2JE

Ofqual
.....



Llywodraeth Cynulliad Cymru
Welsh Assembly Government

