

UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS International General Certificate of Secondary Education

	CANDIDATE NAME					
	CENTRE NUMBER		CANDIDATE NUMBER			
*						
	MATHEMATICS		0580/41			
1914	Paper 4 (Extended	)	May/June 2013			
404320*			2 hours 30 minutes			
	Candidates answer	r on the Question Paper.				
	Additional Materials	s: Electronic calculator Tracing paper (optional)	Geometrical instruments			

#### READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use a pencil for any diagrams or graphs.

Do not use staples, paper clips, highlighters, glue or correction fluid.

DO **NOT** WRITE IN ANY BARCODES.

Answer **all** questions.

If working is needed for any question it must be shown below that question.

Electronic calculators should be used.

If the degree of accuracy is not specified in the question, and if the answer is not exact, give the answer to three significant figures. Give answers in degrees to one decimal place. For  $\pi$ , use either your calculator value or 3.142.

At the end of the examination, fasten all your work securely together. The number of marks is given in brackets [] at the end of each question or part question. The total of the marks for this paper is 130.

This document consists of **19** printed pages and **1** blank page.



1	(a)		e day, Maria took 27 minutes to walk 1.8 km to school. left home at 0748.	For Examiner's Use
		(i)	Write down the time Maria arrived at school.	
			Answer(a)(i) [1]	
		( <b>ii</b> )	Show that Maria's average walking speed was 4 km/h.	
			Answer(a)(ii)	
			[2]	
	(b)	And	other day, Maria cycled the 1.8 km to school at an average speed of 15 km/h.	
		(i)	Calculate the percentage <b>increase</b> that 15 km/h is on Maria's walking speed of 4 km/h.	
			Answer(b)(i) % [3]	
		( <b>ii</b> )	Calculate the percentage <b>decrease</b> that Maria's cycling time is on her walking time of 27 minutes.	

Answer(b)(ii) ...... % [3]


Calculate the time Maria takes to walk to her friend's home.

For Examiner's Use

*Answer(b)*(iii) ..... min [2]


$$f(x) = 3 - x - x^2$$
  $g(x) = 3^x$ 

-1.5 -1 -0.5 0 1 1.5 0.5 х 3 1 **f**(*x*) 2.25 3.25 2.25 -0.75-1.5-1 -0.5 0 0.5 1 1.5 x 0.19 0.58 3 5.20 g(x)1.73



2

<b>(b)</b>	On the grid, drav	w the graphs of	y = f(x) and	nd $y = g(x)$ for	$-1.5 \leq x \leq 1.5$ .
------------	-------------------	-----------------	--------------	-------------------	--------------------------



For

Examiner's


Use

[3]

(c)	For	$-1.5 \le x \le 1.5$ , use your graphs to solve	For Examiner's
	(i)	$\mathbf{f}(x) = 0,$	Use
	(ii)	$Answer(c)(i) \ x =$	
	(iii)	Answer(c)(ii) x =	
		$Answer(c)(iii) x = \dots [1]$	
( <b>d</b> )	Byo	drawing a suitable tangent, find an estimate of the gradient of the graph of $y = f(x)$ when $x = 0.5$ .	
		Answer(d)	


Examiner's

Use



3 200 students estimate the mass (*m* grams) of a coin. The cumulative frequency diagram shows the results.

				7	,			
Fine	1							For Examiner's
(i)	the median,							Use
(ii)	the upper qu	artile,		Answer(a)(	i)		g [1]	
				Answer(a)(i	i)		g [1]	
(111)	the 80th perc	centile,						
				Answer(a)(ii	i)		g [1]	
(iv)	the number of	of students who	ose estimate is	7 g or less.				
				Answer(a)(iv	7)		[1]	
(i)	Use the cum	ulative frequer	ncy diagram to	complete the	frequency tabl	le.		
Mass	s ( <i>m</i> grams)	$0 < m \le 2$	$2 < m \leq 4$	$4 < m \le 6$	$6 < m \le 8$	$8 < m \le 10$		
Freq	uency	40				2		
(ii)	The probabil	lity that the stu		s that the mass	is greater than	n <i>M</i> grams is 0.	[2] 3.	
			Ar	uswer(b)(ii) M	=		[2]	
	(i) (ii) (iii) (iv) (i) Mass Freq	<ul> <li>(ii) the upper quint is the source of the source</li></ul>	<ul> <li>(i) the median,</li> <li>(ii) the upper quartile,</li> <li>(iii) the 80th percentile,</li> <li>(iv) the number of students when</li> <li>(i) Use the cumulative frequer</li> <li>Mass (m grams) 0 &lt; m ≤ 2</li> <li>Frequency 40</li> </ul>	(i) the median, (ii) the upper quartile, (iii) the 80th percentile, (iv) the number of students whose estimate is (i) Use the cumulative frequency diagram to Mass (m grams) $0 < m \le 2$ $2 < m \le 4$ Frequency 40 (ii) A student is chosen at random. The probability that the student estimates Find the value of M.	Find         (i) the median,         (ii) the upper quartile,         (iii) the upper quartile,         (iii) the 80th percentile,         (iv) the number of students whose estimate is 7g or less.         (iv) the number of students whose estimate is 7g or less.         (ii) Use the cumulative frequency diagram to complete the $Mass (m \text{ grams})$ $0 < m \le 2$ $2 < m \le 4$ $4 < m \le 6$ Frequency       40 $u$ $u$ $u$ (ii) A student is chosen at random. The probability that the student estimates that the mass Find the value of $M$ . $Mast (M = Mast (M = Ma$	Find (i) the median, (ii) the upper quartile, (ii) the upper quartile, (iii) the 80th percentile, (iv) the number of students whose estimate is 7 g or less. (iv) the number of students whose estimate is 7 g or less. (i) Use the cumulative frequency diagram to complete the frequency table $Mass (m \text{ grams})$ $0 < m \le 2$ $2 < m \le 4$ $4 < m \le 6$ $6 < m \le 8$ Frequency (i) A student is chosen at random. The probability that the student estimates that the mass is greater than Find the value of $M$ .	Find (i) the median, (ii) the upper quartile. (ii) the upper quartile. (iii) the upper quartile. (iv) the number of students whose estimate is 7 g or less. (iv) the number of students whose estimate is 7 g or less. (iv) the number of students whose estimate is 7 g or less. (iv) the number of students whose estimate is 7 g or less. (iv) the number of students whose estimate is 7 g or less. (iv) the number of students whose estimate is 7 g or less. (iv) the number of students whose estimate is 7 g or less. (ii) Use the cumulative frequency diagram to complete the frequency table. (iii) Assudent is chosen at random. The probability that the student estimates that the mass is greater than M grams is 0. Find the value of M.	Find (i) the median, $Answer(a)(i) \qquad \qquad$




Examiner's Use

Height ( <i>h</i> cm)	$150 < h \le 160$	$160 < h \le 165$	$165 < h \le 180$	$180 < h \le 190$
Frequency	5	9	18	10

The table shows information about the heights of a group of 42 students.

(a) Using mid-interval values, calculate an estimate of the mean height of the students. Show your working.




5

For

Examiner's

Use



In the diagram, *BCD* is a straight line and *ABDE* is a quadrilateral. Angle  $BAC = 90^{\circ}$ , angle  $ABC = 30^{\circ}$  and angle  $CAE = 52^{\circ}$ . AC = 15.7 cm, CE = 16.5 cm and CD = 23.4 cm.

(a) Calculate *BC*.

6

Answer(a)  $BC = \dots$  [3]

(b) Use the sine rule to calculate angle AEC.Show that it rounds to 48.57°, correct to 2 decimal places.

Answer(b)

# (c) (i) Show that angle $ECD = 40.6^{\circ}$ , correct to 1 decimal place. Answer(c)(i)

(ii) Calculate DE.

 $Answer(c)(ii) DE = \dots cm$  [4]

(d) Calculate the area of the quadrilateral *ABDE*.

Answer(d) .....  $cm^2$  [4]

For Examiner's Use

[2]

Examiner's

Use

(a)  $\begin{pmatrix}
A \\
(x+2) cm \\
B \\
(x+2) cm \\
B \\
(x+2) cm \\
C
\end{pmatrix}$ NOT TO SCALE NOT TO SCALE NOT TO SCALE NOT TO SCALE C The triangle *ABC*, *AB* = (x+2) cm and *AC* = (2x+3) cm. sin *ACB* =  $\frac{9}{16}$ Find the length of *BC*.

Answer(a) BC = ..... cm [6]

- (b) A bag contains 7 white beads and 5 red beads.
  - (i) The mass of a red bead is 2.5 grams more than the mass of a white bead. The total mass of all the 12 beads is 114.5 grams.

Find the mass of a white bead and the mass of a red bead.

Answer(b)(i) White ..... g


Red ..... g [5]

7

( <b>ii</b> )	Two	beads are taken out of the bag at random, without replacement.	For Examiner's Use				
	Finc	Find the probability that					
	(a)	they are both white,					
		<i>Answer(b)</i> (ii)(a)	]				
	(b)	one is white and one is red.					
		<i>Answer(b)</i> (ii)(b)	]				

[Turn over

8 (a)



(ii) BC = CD. Calculate angle *CBD*.


 $Answer(a)(ii) Angle CBD = \dots [1]$ 

(iii) This pentagon also has one line of symmetry. Calculate angle *ADB*.


Answer(a)(iii) Angle  $ADB = \dots$ [1]

(b) A, B and C lie on a circle centre O. Angle  $AOC = 3y^{\circ}$  and angle  $ABC = (4y + 4)^{\circ}$ .

Find the value of *y*.



For Examiner's Use



$$f(x) = x^2 + x - 3$$
  $g(x) = 2x + 7$   $h(x) = 2^x$ 

For Examiner's Use

(a) Solve the equation f(x) = 0. Show all your working and give your answers correct to 2 decimal places.


*Answer*(*a*) x = ...... [4]


**(b)**  $fg(x) = px^2 + qx + r$ 

Find the values of p, q and r.

Answer(b)  $p = \dots$  $q = \dots$ 

 $r = \dots [3]$ 





(c) The total number of dots in the first *n* stars is given by the expression  $5n^2 + 6n$ .

(i) Show that this expression is correct when n = 3.

Answer(c)(i)

[2]

For

Examiner's Use

(ii) Find the total number of dots in the first 10 stars.

*Answer*(*c*)(ii) ..... [1]

(d) The total number of dots in the first *n* stars is  $5n^2 + 6n$ . The number of dots in the (n + 1)th star is 10(n + 1) + 1.

Add these two expressions to show that the total number of dots in the first (n + 1) stars is

$$5(n+1)^2 + 6(n+1)$$
.

You must show each step of your working.

Answer(d)

### **BLANK PAGE**

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.