



Cambridge Assessment International Education
Cambridge International General Certificate of Secondary Education

BIOLOGY

0610/61

Paper 6 Alternative to Practical

May/June 2019

MARK SCHEME

Maximum Mark: 40

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the May/June 2019 series for most Cambridge IGCSE™, Cambridge International A and AS Level and Cambridge Pre-U components, and some Cambridge O Level components.

This syllabus is regulated for use in England, Wales and Northern Ireland as a Cambridge International Level 1/Level 2 Certificate.

This document consists of **8** printed pages.

Generic Marking Principles

These general marking principles must be applied by all examiners when marking candidate answers. They should be applied alongside the specific content of the mark scheme or generic level descriptors for a question. Each question paper and mark scheme will also comply with these marking principles.

GENERIC MARKING PRINCIPLE 1:

Marks must be awarded in line with:

- the specific content of the mark scheme or the generic level descriptors for the question
- the specific skills defined in the mark scheme or in the generic level descriptors for the question
- the standard of response required by a candidate as exemplified by the standardisation scripts.

GENERIC MARKING PRINCIPLE 2:

Marks awarded are always **whole marks** (not half marks, or other fractions).

GENERIC MARKING PRINCIPLE 3:

Marks must be awarded **positively**:

- marks are awarded for correct/valid answers, as defined in the mark scheme. However, credit is given for valid answers which go beyond the scope of the syllabus and mark scheme, referring to your Team Leader as appropriate
- marks are awarded when candidates clearly demonstrate what they know and can do
- marks are not deducted for errors
- marks are not deducted for omissions
- answers should only be judged on the quality of spelling, punctuation and grammar when these features are specifically assessed by the question as indicated by the mark scheme. The meaning, however, should be unambiguous.

GENERIC MARKING PRINCIPLE 4:

Rules must be applied consistently e.g. in situations where candidates have not followed instructions or in the application of generic level descriptors.

GENERIC MARKING PRINCIPLE 5:

Marks should be awarded using the full range of marks defined in the mark scheme for the question (however; the use of the full mark range may be limited according to the quality of the candidate responses seen).

GENERIC MARKING PRINCIPLE 6:

Marks awarded are based solely on the requirements as defined in the mark scheme. Marks should not be awarded with grade thresholds or grade descriptors in mind.

Question	Answer	Marks	Guidance
1(a)(i)	table drawn with a minimum of two columns and a line separating headings from data ; headings with correct units: percentage concentration of citric acid (solution) and distance (travelled) / cm or mm ; three correct distances recorded ;	3	
1(a)(ii)	from (edge of) hole to (edge of) the (red or yellow) circle ; diameter / radius of (red or yellow) circle ;	1	
1(a)(iii)	the higher the concentration (of solution) the further (the acid) moves / AW;	1	
1(a)(iv)	6.5 ;;	2	
1(a)(v)	<i>green agar pH: 7 ;</i> <i>red agar pH: 1 / 2 ;</i>	2	A value in range 7–8 / neutral A value lower than 7/ acidic
1(b)(i)	number of drops / three drops (of citric acid / solution in the holes) ; agar / agar concentration / depth of agar / volume of agar ; size of, holes / wells / AW ; type of indicator / concentration or volume of indicator ; time / 30 minutes ; temperature ; AVP ; e.g. one type of acid / same stock solution of citric acid	2	

Question	Answer	Marks	Guidance												
1(b)(ii)	<table border="1"> <thead> <tr> <th data-bbox="344 220 721 284"><i>error</i></th> <th data-bbox="721 220 1252 284"><i>effect on results</i></th> </tr> </thead> <tbody> <tr> <td data-bbox="344 284 721 384">drop sizes vary ;</td> <td data-bbox="721 284 1252 384">larger volume produces greater diffusion distance ; ora</td> </tr> <tr> <td data-bbox="344 384 721 448">no repeats ;</td> <td data-bbox="721 384 1252 448">unable to identify anomalous results ;</td> </tr> <tr> <td data-bbox="344 448 721 584">one dropping pipette used for all three solutions / contamination;</td> <td data-bbox="721 448 1252 584">a weaker solution would produce a smaller diffusion distance ;</td> </tr> <tr> <td data-bbox="344 584 721 748">longer diffusion time / citric acid added at different times / circles measured at different times ;</td> <td data-bbox="721 584 1252 748">greater / lesser distance travelled ;</td> </tr> <tr> <td data-bbox="344 748 721 884">AVP; e.g. edge of colour change difficult to judge / subjective</td> <td data-bbox="721 748 1252 884">matching AVP effect; e.g. over/under estimation of diffusion distance</td> </tr> </tbody> </table>	<i>error</i>	<i>effect on results</i>	drop sizes vary ;	larger volume produces greater diffusion distance ; ora	no repeats ;	unable to identify anomalous results ;	one dropping pipette used for all three solutions / contamination;	a weaker solution would produce a smaller diffusion distance ;	longer diffusion time / citric acid added at different times / circles measured at different times ;	greater / lesser distance travelled ;	AVP; e.g. edge of colour change difficult to judge / subjective	matching AVP effect; e.g. over/under estimation of diffusion distance	2	AW throughout mark as a pair, effect must match the source of error
<i>error</i>	<i>effect on results</i>														
drop sizes vary ;	larger volume produces greater diffusion distance ; ora														
no repeats ;	unable to identify anomalous results ;														
one dropping pipette used for all three solutions / contamination;	a weaker solution would produce a smaller diffusion distance ;														
longer diffusion time / citric acid added at different times / circles measured at different times ;	greater / lesser distance travelled ;														
AVP; e.g. edge of colour change difficult to judge / subjective	matching AVP effect; e.g. over/under estimation of diffusion distance														

Question	Answer	Marks	Guidance
1(c)	<p><i>any two from given method ;;</i></p> <ul style="list-style-type: none"> • agar plates used • indicator (in agar) • holes made in agar • (citric) acid added • left for fixed (stated) time • measure distance moved/diameter/radius <p><i>any three from novel method ;;;</i></p> <ul style="list-style-type: none"> • using range of at least 2 temperatures • temperatures specified and all less than 70 (°C) • idea of keeping temperature constant • ref to temperature equilibration time prior to adding citric acid • measure time taken to reach specified diameter <p><i>any additional points:</i> same concentration of citric acid ; wear gloves / goggles ; repeating the investigation at least twice / three holes in each plate / use three dishes at each temperature ; AVP ;</p>	6	

Question	Answer	Marks	Guidance
2(a)(i)	<i>line:</i> clear single continuous lines without shading ; <i>size:</i> occupies at least half the space available ; <i>detail:</i> layers in correct proportions ;	3	
2(a)(ii)	length of line AB = 40 mm ; actual length = 0.4 mm ;;	3	A ±1 mm A ±0.01 mm
2(a)(iii)	X written on the spongy mesophyll of drawing ;	1	
2(b)(i)	to obtain a representative leaf size ; to identify anomalous results ;	1	
2(b)(ii)	to avoid bias / gain a representative sample / AW ; so that a comparison can be made (between the different light intensities) ;	1	A leaves at different heights may be different sizes
2(b)(iii)	draw round the outline of the leaf on a grid / place leaf under a (transparent) grid ; count the squares ; include any squares more than half covered / other valid method described ;	2	
2(b)(iv)	light intensity ;	1	
2(c)(i)	67(%) ;;	2	

Question	Answer	Marks	Guidance
2(c)(ii)	<p><i>axes labelled with units:</i> light intensity / arbitrary units or au and average leaf area / mm² and species A and B labelled / key given ;</p> <p><i>scale and size:</i> even scale for leaf area sequential for x-axis bars/plotting area to occupy at least half the grid in both directions ;</p> <p><i>plots:</i> 6 values plotted accurately $\pm \frac{1}{2}$ small square ;</p> <p>4 bars: bars the same width (at least 1 small square wide) gaps present between bars / pairs of bars ;</p>	4	
2(c)(iii)	<p><i>species A:</i> as the light intensity decreases the (average) leaf area increases / ; ora</p> <p><i>species B:</i> (average) leaf area increases with increasing light intensity (to maximum at 50 au) and then decreases ; ora</p>	2	
2(c)(iv)	measure leaf growth at a narrower range of light intensities around 50 (au) ;	1	