



UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS International General Certificate of Secondary Education

CANDIDATE NAME		
CENTRE NUMBER		CANDIDATE NUMBER
CHEMISTRY		0620/06
Paper 6 Alterna	tive to Practical	October/November 2007

Candidates answer on the Question Paper.

No additional materials are required.

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use a pencil for any diagrams, graphs or rough working.

Do not use staples, paper clips, highlighters, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

Answer all questions.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.


For Examiner's Use		
1		
2		
3		
4		
5		
6		
7		
Total		

1 hour

This document consists of 13 printed pages and 3 blank pages.



1 Hydrated copper sulphate crystals, CuSO₄.5H₂O were heated in the apparatus shown below.



- (a) Indicate on the diagram using arrows
 - (i) where the copper sulphate crystals are placed,
 - (ii) where heat is applied.

[2]


(b) What is the purpose of the ice?

[1	1
 -	•

(c) The crystals changed colour from ______to ____[2]

[Total: 5]

2 An experiment was set up to investigate the rusting of iron.



(a)	Describe the	appearance	of the	iron	after	rusting.
-----	--------------	------------	--------	------	-------	----------

Γ	11	1
	ı	ı
		4

(b)	(i)	Why does	the water	r rise up	the tube?
-----	-----	----------	-----------	-----------	-----------

Γ1	1
 r.	4

(11)	Calculate the percentage change in the volume of air in the tube.	

۲ 1 1	ı
נין	

- (c) What difference would be observed if
 - (i) an iron nail was suspended in the tube instead of using iron filings,

га	-	
П	. 1	
r.	7	

(ii) the water contained salt?

[11]

[Total: 5]

[Total: 6]

3	The information	in the box is	about the	preparation o	of zinc nitrate crystals	
---	-----------------	---------------	-----------	---------------	--------------------------	--

Step 1: Add a small amount of zinc oxide to some hot dilute nitric acid, and stir.

Step 2: Keep adding zinc oxide until it is in excess.

Step 3: Remove the excess zinc oxide to leave colourless zinc nitrate solution.

Step 4: Evaporate the zinc nitrate solution until it is saturated.

Step 5: Leave the *saturated solution* to cool. White crystals form on cooling.

Step 6: Remove the crystals from the remaining solution.

Step 7: Dry the crystals on a piece of filter paper.

(a)	Sug	gest a reason for using excess zinc oxide in Step 2.	
			 [1]
/h\	Cuar		1
(D)	Sug	gest how the excess zinc oxide can be removed from the solution in Step 3.	[1]
(c)	(i)	What is meant by the term saturated solution?	
	(ii)	What practical method could show the solution to be saturated?	[2]
	į		[1]
(d)	Why	are the crystals dried in Step 7 using filter paper instead of by heating?	
			[1]


4 A student investigated the reaction of dilute hydrochloric acid with two different solids, calcium carbonate (marble) and calcium oxide. Four experiments were carried out.

Experiment 1

By using a measuring cylinder, $50 \, \text{cm}^3$ of dilute hydrochloric acid was poured into a polystyrene cup and the initial temperature of the acid was measured. 2.5 g of small marble chips were added to the cup and the mixture stirred with the thermometer.

The temperature of the mixture was measured after 2 minutes.

Use the thermometer diagrams to record the temperatures in the table of results on page 6.




Experiment 2

Experiment 1 was repeated using 2.5g of powdered calcium carbonate. Use the thermometer diagrams to record the results in the table.



Experiment 3

Experiment 1 was repeated using 1.5 g of lumps of calcium oxide. Use the thermometer diagrams to record the temperatures in the table.



Experiment 4

Experiment 1 was repeated using 1.5g of powdered calcium oxide. Use the thermometer diagrams to record the results in the table.



initial temperature/°C

final temperature/°C


Table of results

Evnoriment	temperature/°C			
Experiment	initial	final	difference	
1				
2				
3				
4				

		[4]
(a)	What would be observed in Experiment 2?	
		[1]

[3]

(b) Draw a bar chart of the results of the experiments on the grid below.



experiment number

(c)	Wh	ich experiment produced	
	(i)	the smallest temperature change,	
			[1]
	(ii)	the largest temperature change?	
			[1]
(d)	Giv	e two reasons why the temperature changes are different in (c) .	
	1.		
	2.		
			[2]

(e)	In Experiment 1, how would you know which reactant is in excess? Explain your answer.
	rol
	[2]
(f)	Explain how the temperature changes would differ in the experiments if 100 cm ³ of hydrochloric acid were used.
	[2]
	[Total: 16]

5 Three different liquids **P**, **Q** and **R** were analysed. **Q** was an aqueous solution of sodium hydroxide.


The tests on the liquids and some of the observations are in the following table. Complete the observations in the table.

tests	observations	
(a) Test the pH of the liquids using indicator paper. Note the colour of the paper.	P colour red	
·	pH 1	
	Q colour	
	pH[2]	
	R colour orange	
	pH 5	
(b) (i) Add a 5 cm piece of magnesium to about 3 cm ³ of liquid P in a test–tube. Test the gas given off.	bubbles of gas lighted splint pops	
(ii) Repeat (b)(i) using liquids Q , and R . Do not test for any gases.	Q[2]	

For Examiner's Use

	tests	observations	
	(c) To about 2 cm ³ of liquid P add 1 spatula measure of sodium carbonate. Test the gas given off.	[3]	
	(d) By using a teat pipette add aqueous silver nitrate to about 1 cm³ of liquid P .	white precipitate	
	(e) By using a teat pipette add liquid Q to about 1 cm ³ of aqueous iron(II) sulphate.	[2]	
(f)	Name the gas given off in test (b)(i).	[1]	
(g)	Name the gas given off in test (c).	[1]	
(h)	Identify liquid P .	[1]	
(i)	What conclusions can you draw about liqu		
		[2]	
		[Total: 14]	

6 Magnesium reacts with dilute sulphuric acid to form hydrogen gas. The speed of the reaction was investigated using the apparatus below.




In an experiment 50 cm³ of dilute sulphuric acid was added to a large piece of magnesium. A student measured the total volume of gas produced at 2 minute intervals.

Use the measuring cylinder diagrams to complete the table

time/minutes	measuring cylinder diagram	total volume of collected/cm³
0	5 	
2	10 15 20	
4	—25 = = 30 = = = 35	
6	25 = 30 = 35	
8	35 40 45	
10	40 = 45 = 50	
12	40 = 45 = 50	

(a) Plot the student's results on the grid. Use the points to draw a smooth line graph.



[3]

(b) (i) At which time does the result appear to be inaccurate?

[1]

(ii) Use the graph to deduce what the correct volume should be at this time.


[1]

[Total:8]

For Examiner's Use

7 Diesel is a liquid fuel obtained from crude oil. Biodiesel is a fuel made from oil obtained from the seeds of plants such as sunflowers.

Using the apparatus below plan an experiment to investigate which of these two fuels produces more energy.



••••••	 	
		[6]
	 	 [6]

[Total: 6]

BLANK PAGE

BLANK PAGE

16

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.