

Cambridge IGCSE™

CHEMISTRY

0620/62 October/November 2023

Paper 6 Alternative to Practical MARK SCHEME Maximum Mark: 40

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the October/November 2023 series for most Cambridge IGCSE, Cambridge International A and AS Level components, and some Cambridge O Level components.

Generic Marking Principles

These general marking principles must be applied by all examiners when marking candidate answers. They should be applied alongside the specific content of the mark scheme or generic level descriptors for a question. Each question paper and mark scheme will also comply with these marking principles.

GENERIC MARKING PRINCIPLE 1:

Marks must be awarded in line with:

- the specific content of the mark scheme or the generic level descriptors for the question ٠
- the specific skills defined in the mark scheme or in the generic level descriptors for the question .
- the standard of response required by a candidate as exemplified by the standardisation scripts.

GENERIC MARKING PRINCIPLE 2:

Marks awarded are always whole marks (not half marks, or other fractions).

GENERIC MARKING PRINCIPLE 3:

Marks must be awarded **positively**:

- marks are awarded for correct/valid answers, as defined in the mark scheme. However, credit is given for valid answers which go beyond ٠ the scope of the syllabus and mark scheme, referring to your Team Leader as appropriate
- marks are awarded when candidates clearly demonstrate what they know and can do ٠
- marks are not deducted for errors •
- marks are not deducted for omissions .
- answers should only be judged on the quality of spelling, punctuation and grammar when these features are specifically assessed by the • question as indicated by the mark scheme. The meaning, however, should be unambiguous.

GENERIC MARKING PRINCIPLE 4:

Rules must be applied consistently, e.g. in situations where candidates have not followed instructions or in the application of generic level descriptors.

GENERIC MARKING PRINCIPLE 5:

Marks should be awarded using the full range of marks defined in the mark scheme for the question (however; the use of the full mark range may be limited according to the quality of the candidate responses seen).

GENERIC MARKING PRINCIPLE 6:

Marks awarded are based solely on the requirements as defined in the mark scheme. Marks should not be awarded with grade thresholds or grade descriptors in mind.

Science-Specific Marking Principles

- 1 Examiners should consider the context and scientific use of any keywords when awarding marks. Although keywords may be present, marks should not be awarded if the keywords are used incorrectly.
- 2 The examiner should not choose between contradictory statements given in the same question part, and credit should not be awarded for any correct statement that is contradicted within the same question part. Wrong science that is irrelevant to the question should be ignored.
- 3 Although spellings do not have to be correct, spellings of syllabus terms must allow for clear and unambiguous separation from other syllabus terms with which they may be confused (e.g. ethane / ethene, glucagon / glycogen, refraction / reflection).
- 4 The error carried forward (ecf) principle should be applied, where appropriate. If an incorrect answer is subsequently used in a scientifically correct way, the candidate should be awarded these subsequent marking points. Further guidance will be included in the mark scheme where necessary and any exceptions to this general principle will be noted.

5 <u>'List rule' guidance</u>

For questions that require *n* responses (e.g. State **two** reasons ...):

- The response should be read as continuous prose, even when numbered answer spaces are provided.
- Any response marked *ignore* in the mark scheme should not count towards **n**.
- Incorrect responses should not be awarded credit but will still count towards *n*.
- Read the entire response to check for any responses that contradict those that would otherwise be credited. Credit should **not** be awarded for any responses that are contradicted within the rest of the response. Where two responses contradict one another, this should be treated as a single incorrect response.
- Non-contradictory responses after the first *n* responses may be ignored even if they include incorrect science.

6 <u>Calculation specific guidance</u>

Correct answers to calculations should be given full credit even if there is no working or incorrect working, **unless** the question states 'show your working'.

For questions in which the number of significant figures required is not stated, credit should be awarded for correct answers when rounded by the examiner to the number of significant figures given in the mark scheme. This may not apply to measured values.

For answers given in standard form (e.g. $a \times 10^n$) in which the convention of restricting the value of the coefficient (a) to a value between 1 and 10 is not followed, credit may still be awarded if the answer can be converted to the answer given in the mark scheme.

Unless a separate mark is given for a unit, a missing or incorrect unit will normally mean that the final calculation mark is not awarded. Exceptions to this general principle will be noted in the mark scheme.

7 <u>Guidance for chemical equations</u>

Multiples / fractions of coefficients used in chemical equations are acceptable unless stated otherwise in the mark scheme.

State symbols given in an equation should be ignored unless asked for in the question or stated otherwise in the mark scheme.

0620/62

			FUI	BLISHED			
Examples of ho State three rease	w to apply the list ru ons [3]	le					Ţ]
Α	1 Correct	✓		F	1 Correct	✓	
~	2 Correct	√	2	(4 responses)	2 Correct	✓	2
	3 Wrong	×	2		3 Correct CON (of 3.)	× (discount 3)	
В	1 Correct,	✓, ✓		G	1 Correct	✓	
	Correct		3	(5 responses)	2 Correct	\checkmark	
(4 responses)	2 Correct	✓	、		3 Correct	✓	3
	3 Wrong	ignore			Correct CON (of 4.)	ignore ignore	
С	1 Correct	✓					
(4 responses)	2 Correct,	✓,	2	H	1 Correct	✓	
,	Wrong	×	2	(4 responses)	2 Correct	×	2
	3 Correct	ignore			3 CON (of 2.) Correct	(discount 2) ✓	
D	1 Correct	✓					
(4 responses)	2 Correct,	×,		1	1 Correct	✓	
(CON (of 2.)	(discount 2)	2	(4 responses)	2 Correct	×	2
	3 Correct	\checkmark			3 Correct CON (of 2.)	√ (discount 2)	
Е	1 Correct	✓				1	
(4 responses)	2 Correct	\checkmark	3				
	3 Correct, Wrong	~					

Question	Answer	Marks
1(a)	pestle and mortar	1
1(b)	so that all the calcium carbonate reacts / dissolves	1
1(c)	A (filter) funnel	2
	B (conical) flask	
1(d)(i)	hydrochloric acid / calcium chloride	1
1(d)(ii)	M1 dry	2
	M2 determine the mass	

Question	Answer	
2(a)	M1 all volumes of aqueous sodium hydroxide completed correctly (9.0, 8.0, 7.0, 4.0, 3.0, 2.0)	
	M2 and M3 all six temperatures completed correctly (26.0; 28.0; 30.0; 30.5; 29.0; 27.5)	
	M4 all temperatures completed to same number of dp and all volumes of aqueous sodium hydroxide shown to 1 dp	
2(b)	M1 appropriate vertical scale so that points and extrapolation to maximum extend over halfway up scale	5
	M2 and M3 all points plotted correctly	
	M4 two suitable straight lines	
	M5 straight lines extended so that they cross / meet	
2(c)(i)	M1 working shown on graph from where lines cross	3
	M2 volume of dilute hydrochloric acid correct for their graph	
	M3 volume of aqueous sodium hydroxide correct based on their recorded volume of hydrochloric acid	

Question	Answer	Marks
2(c)(ii)	7	1
2(c)(iii)	dilute hydrochloric acid and (as volume) less / low(er) / small(er) than sodium hydroxide (to be neutral / reach maximum temperature)	1
2(d)	M1 pH: no change	3
	M2 temperature: higher / greater / bigger	
	M3 explanation: polystyrene is a better insulator than glass	
2(e)	(volumetric) pipettes not available in those sizes / (volumetric) pipettes do not measure variable volumes	1

Question	Answer	
3(a)(i)	M1 zinc/Zn ²⁺	2
	M2 aluminium / Al ³⁺	
3(a)(ii)	M1 add excess / more ammonia	2
	M2 for zinc ions: precipitate redissolves / colourless solution OR for aluminium ions: the precipitate remains / stays white / unchanged	
3(b)	bromide / Br	1
3(c)	(light / pale / apple) green (flame colour)	1
3(d)	(red) litmus turns blue	1
3(e)	no change / (remains) colourless / no reaction	1
3(f)	white precipitate	1

Question	Answer	Marks
4	any 6 from:	6
	add excess solute, find mass not dissolved	
	MP1 known / stated volume of water	
	MP2 add known / stated mass of sodium sulfate (in excess)	
	MP3 water and solute combined in a suitable named container (test-tube / boiling tube / beaker / flask / polystyrene cup)	
	MP4 stir / mix / shake	
	MP5 filter	
	MP6 dry and weigh residue (ignore washing)	
	MP7 repeat at (an)other temperature(s)	
	OR	
	add solute gradually, stop when no more dissolves	
	MP1 known / stated volume of water	
	MP2 add stated measure of sodium sulfate	
	MP3 water and solute combined in a suitable named container (test-tube / boiling tube / beaker / flask / polystyrene cup)	
	MP4 stir / mix / shake	
	MP5 if all dissolved, add more sodium sulfate	
	MP6 count number of measures / mass added / mass left from initial supply MP7 repeat at (an)other temperature(s)	
	OR	
	add excess solute, find mass that dissolves	
	MP1 known / stated volume of water	
	MP2 add excess of sodium sulfate	
	MP3 water and solute combined in a suitable named container (test-tube / boiling tube / beaker / flask / polystyrene cup)	
	MP4 stir / mix / shake	
	MP5 filter	
	MP6 heat solution to evaporate all water and weigh residue MP7 repeat at (an)other temperature(s)	

Question	Answer	Marks
4	OR	
	fixed mass solute add solvent until all dissolves MP1 add known volumes / fixed quantities of water gradually to MP2 known / stated <u>mass</u> of sodium sulfate MP3 water and solute combined in a suitable named container (test-tube / boiling tube / beaker / flask / polystyrene cup) MP4 stir / mix / shake MP5 if not all dissolved, add more water MP6 measure total volume of water added MP7 repeat at (an)other temperature(s)	
	max 6	