

Cambridge O Level

MATHEMATICS (SYLLABUS D)

Paper 2

MARK SCHEME

Maximum Mark: 100

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the October/November 2021 series for most Cambridge IGCSE™, Cambridge International A and AS Level components and some Cambridge O Level components.

Generic Marking Principles

These general marking principles must be applied by all examiners when marking candidate answers. They should be applied alongside the specific content of the mark scheme or generic level descriptors for a question. Each question paper and mark scheme will also comply with these marking principles.

GENERIC MARKING PRINCIPLE 1:

Marks must be awarded in line with:

- the specific content of the mark scheme or the generic level descriptors for the question
- the specific skills defined in the mark scheme or in the generic level descriptors for the question
- the standard of response required by a candidate as exemplified by the standardisation scripts.

GENERIC MARKING PRINCIPLE 2:

Marks awarded are always whole marks (not half marks, or other fractions).

GENERIC MARKING PRINCIPLE 3:

Marks must be awarded **positively**:

- marks are awarded for correct/valid answers, as defined in the mark scheme. However, credit is given for valid answers which go beyond the scope of the syllabus and mark scheme, referring to your Team Leader as appropriate
- marks are awarded when candidates clearly demonstrate what they know and can do
- marks are not deducted for errors
- marks are not deducted for omissions
- answers should only be judged on the quality of spelling, punctuation and grammar when these features are specifically assessed by the question as indicated by the mark scheme. The meaning, however, should be unambiguous.

GENERIC MARKING PRINCIPLE 4:

Rules must be applied consistently, e.g. in situations where candidates have not followed instructions or in the application of generic level descriptors.

GENERIC MARKING PRINCIPLE 5:

Marks should be awarded using the full range of marks defined in the mark scheme for the question (however; the use of the full mark range may be limited according to the quality of the candidate responses seen).

GENERIC MARKING PRINCIPLE 6:

Marks awarded are based solely on the requirements as defined in the mark scheme. Marks should not be awarded with grade thresholds or grade descriptors in mind.

© UCLES 2021 Page 2 of 8

Maths-Specific Marking Principles			
1	Unless a particular method has been specified in the question, full marks may be awarded for any correct method. However, if a calculation is required then no marks will be awarded for a scale drawing.		
2	Unless specified in the question, answers may be given as fractions, decimals or in standard form. Ignore superfluous zeros, provided that the degree of accuracy is not affected.		
3	Allow alternative conventions for notation if used consistently throughout the paper, e.g. commas being used as decimal points.		
4	Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored (isw).		
5	Where a candidate has misread a number in the question and used that value consistently throughout, provided that number does not alter the difficulty or the method required, award all marks earned and deduct just 1 mark for the misread.		
6	Recovery within working is allowed, e.g. a notation error in the working where the following line of working makes the candidate's intent clear.		

Abbreviations

cao correct answer only

dep dependent

FT follow through after error isw ignore subsequent working

oe or equivalent SC Special Case

nfww not from wrong working

soi seen or implied

© UCLES 2021 Page 3 of 8

Question	Answer	Marks	Partial Marks
1(a)(i)	1980	2	M1 for $2250 - \frac{12}{100} \times 2250$ oe or B1 for 270
			After 0 scored, SC1 for answer 3960
1(a)(ii)	14 : 31 final answer	2	M1 for 700 : 1550 oe
- (1)(1)		_	After 0 scored, SC1 for answer 31 : 14
1(b)	77.65 cao	3	M2 for $350 - \frac{19500}{71.6}$ oe OR M1 for 350×71.6 soi M1 for $\frac{their 25060 - 19500}{71.6}$ oe
1(c)(i)	1.333×10^7 final answer	1	
1(c)(ii)	749 or 7.49×10^2 cao	2	M1 for $\frac{4.44 \times 10^{10}}{5.93 \times 10^7}$ oe
1(c)(iii)	739 000 or 7.39 × 10 ⁵	2	M1 for $\frac{(100+23.5)}{100}x = 9.13 \times 10^5$ soi
2(a)(i)	16 to 20	1	
2(a)(ii)	240	2	M1 for $\frac{90}{54}$ [×144] or $\frac{144}{54}$ [×90] or $54x = 90 \times 144$
2(b)(i)	Correct histogram	3	B1 for 3 or more rectangles on correct bases B1 for 3 or more correct frequency densities soi
2(b)(ii)	28.8	2	M1 for $\frac{30+42}{250}[\times 100]$ oe or for $\frac{k}{250} \times 100$, where $42 < k < 102$ but $k \neq 75$
3(a)	$-5.5 \text{ or } -5\frac{1}{2} \text{ or } -\frac{11}{2}$	1	
3(b)	Correct smooth curve	3	B2FT for 6 or 7 points correctly plotted or B1FT for 4 or 5 points correctly plotted
3(c)	Line $y = 3$ only intersects the graph once oe	2	M1 for $\frac{x^3}{2} - 3x - 1 = 3$ soi or $y = 3$ soi

© UCLES 2021 Page 4 of 8

Question	Answer	Marks	Partial Marks
3(d)(i)	Ruled line through $(1, 1)$ and $(-2, -1)$	1	
3(d)(ii)	$\frac{2}{3}$ nfww	2	M1 for gradient = $\frac{1+1}{1+2}$ oe
3(d)(iii)	FT reading three <i>x</i> -values where <i>their L</i> intersects <i>their</i> curve	2	B1FT for two correct
4(a)(i)	20 24	1	Both correct
4(a)(ii)	4n + 4 oe final answer	2	B1 for $4n + k$ oe seen
4(a)(iii)	36	2	M1 for <i>their</i> $(4p + 4) = 150$ soi
4(b)(i)	44	2	M1 for $\frac{26-2}{4}$ or difference = [-]6
4(b)(ii)	50-6n oe final answer	2	B1 for $-6n + k$ oe seen
5(a)	4c + 3e = 85 oe $2c + 5e = 67$ oe	B1	
	Correct method to eliminate one variable	M1	FT their equations
	[Card =] 16 [Envelope =] 7 final answer	A2	A1 for either $c = 16$ or $e = 7$ If A0 scored, SC1FT for a pair of positive values that satisfy either equation or for correct answers with no working
5(b)	(x+5)(x-5) nfww final answer	1	
5(c)	$\frac{5r}{r-2}$ or $\frac{-5r}{2-r}$ final answer	3	B1 for $rt - 5r = 2t$ or $\frac{rt - 5r}{2} = t$
			M1 for isolation of terms in <i>t</i> M1 for factorising and completing to <i>t</i> = Maximum 2 marks if final answer not correct
5(d)	$\frac{5x+19}{(x-5)(2x+1)} \text{ or } \frac{5x+19}{2x^2-9x-5}$	3	B1 for $4(2x+1)-3(x-5)$ oe isw
	final answer		B1 for denominator $(x-5)(2x+1)$ oe isw

© UCLES 2021 Page 5 of 8

Question	Answer	Marks	Partial Marks
6(a)(i)a	$\frac{1}{8}$ oe	1	
6(a)(i)b	$\frac{5}{8}$ oe	1	
6(a)(ii)	$\frac{9}{64}$ oe	2	M1 for $\frac{3}{8} \times \frac{3}{8}$
6(b)	$\frac{13}{40}$ oe	3	M2 for $\frac{7}{16} \times \frac{6}{15} + \frac{6}{16} \times \frac{5}{15} + \frac{3}{16} \times \frac{2}{15}$ oe or M1 for $\frac{7}{16} \times \frac{6}{15}$ or $\frac{6}{16} \times \frac{5}{15}$ or $\frac{3}{16} \times \frac{2}{15}$ After 0 scored, SC1 for answer $\frac{47}{128}$
7(a)(i)	(-1, 4.5)	1	
7(a)(ii)	(-1, 13)	1	
7(a)(iii)	7.21[1]	2	M1 for $(-4)^2 + 6^2$ oe
7(b)(i)	$\mathbf{b} - \mathbf{a}$	1	
7(b)(ii)	$\frac{1}{4}\mathbf{a} + \frac{1}{4}\mathbf{b} \text{or} \frac{1}{4}(\mathbf{a} + \mathbf{b})$	3	M1 for correct vector route along the lines of the diagram B1 for $\overrightarrow{BC} = \frac{\mathbf{a}}{2}$ soi or for $\overrightarrow{NB} = \frac{1}{4} their(\mathbf{b} - \mathbf{a})$ soi or $\overrightarrow{NA} = \frac{3}{4} their(\mathbf{a} - \mathbf{b})$ soi
8(a)	$\frac{3\times110}{\pi\times3.5^2} \text{ oe}$	M2	M1 for $\frac{1}{3} \times \pi \times 3.5^2 \times h = 110$ oe
	= 8.573 to 8.574	A1	
8(b)	9.26 or 9.256 to 9.262	2	M1 for $3.5^2 + 8.57^2$

© UCLES 2021 Page 6 of 8

Question	Answer	Marks	Partial Marks
8(c)	135.7 to 136.1 nfww	4	M3 for $\frac{360 \times \pi \times 7}{2 \times \pi \times their 9.26}$ oe or M2 for $\frac{x}{360} \times 2 \times \pi \times their 9.26 = \pi \times 7$ oe or M1 for $\frac{x}{360} \times 2 \times \pi \times their 9.26$ seen or $\frac{\pi \times 7}{360}$ oe seen Alternative method: M3 for $\frac{360 \times \pi \times 3.5 \times their 9.26}{\pi \times (their 9.26)^2}$ oe or M2 for $\frac{x}{360} \times \pi \times (their 9.26)^2 = \pi \times 3.5 \times their 9.26$ oe or M1 for $\frac{x}{360} \times \pi \times (their 9.26)^2$ seen or $\pi \times 3.5 \times their 9.26$ seen
8(d)	8.01	2	M1 for $\sqrt[3]{\frac{165}{110}}$ oe or $\sqrt[3]{\frac{110}{165}}$ oe or $\left(\frac{7}{x}\right)^3 = \frac{110}{165}$ oe
9(a)	[W =] x + 5 $[L =] 2(x + 5) oe final answers$	2	B1 for $[W =]x + 5$ or B1FT for $[L =]2 \times their$ algebraicW
9(b)	$(x+5)\times 2(x+5) + 2(x\times(x+5))$ $+2(x\times 2(x+5))$ oe	M2	FT their algebraic expressions in x for length and width B1FT for two different areas seen e.g. two of $(x + 5) \times 2(x + 5)$, $x(x + 5)$, $x \times 2(x + 5)$ or $2((x + 5) \times 2(x + 5) + x(x + 5) + x \times 2(x + 5))$
	$2x^2 + 20x + 50 + 2x^2 + 10x + 4x^2 + 20x = 210$	M1	Set equal to 210 and expansion of brackets. Must have three different areas from width and length of form $ax + b$, a and $b \ne 0$
	Correct simplification to $4x^2 + 25x - 80 = 0$	A1	
9(c)	$\frac{-25 \pm \sqrt{25^2 - 4 \times 4 \times -80}}{2 \times 4} \text{ oe or }$ $\frac{-25}{8} \pm \sqrt{\left(\frac{25}{8}\right)^2 - \frac{-80}{4}}$	B2	B1 for $\sqrt{25^2 - 4 \times 4 \times -80}$ oe or $\frac{-25 \pm []}{2 \times 4}$ oe or $\left(x + \frac{25}{8}\right)^2$
	2.33 and -8.58	B1	

© UCLES 2021 Page 7 of 8

Question	Answer	Marks	Partial Marks
9(d)	250 or 251 or 250.3 to 250.5	2	M1 for their 2.33 × (their 2.33 + 5) × 2(their 2.33 + 5)
9(e)	6.5 nfww	3	B1 for 255 and 261.5 seen M1 for <i>their</i> 261.5 – <i>their</i> 255
10(a)	207.2 to 207.3 nfww	4	B1 for $\angle BAD = 55$ soi nfww M2 for sin $[] = \frac{290\sin(their55)}{350}$ or M1 for $\frac{350}{\sin(their55)} = \frac{290}{\sin[]}$ oe
10(b)	17.6 to 17.8 nfww	5	M1 for $CD = 350 \sin (70 - their 42.7)$ oe or $350 \sin (their (a) - 180)$ oe AND M3 for $\tan = \frac{290 \tan 10}{their CD}$ or M2 for [height of mast =] 290 tan 10 or M1 for $\tan 10 = \frac{[]}{290}$

© UCLES 2021 Page 8 of 8