

Cambridge O Level

MATHEMATICS (SYLLABUS D)

Paper 2 MARK SCHEME Maximum Mark: 100 4024/21 October/November 2023

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the October/November 2023 series for most Cambridge IGCSE, Cambridge International A and AS Level components, and some Cambridge O Level components.

Generic Marking Principles

These general marking principles must be applied by all examiners when marking candidate answers. They should be applied alongside the specific content of the mark scheme or generic level descriptors for a question. Each question paper and mark scheme will also comply with these marking principles.

GENERIC MARKING PRINCIPLE 1:

Marks must be awarded in line with:

- the specific content of the mark scheme or the generic level descriptors for the question
- the specific skills defined in the mark scheme or in the generic level descriptors for the question
- the standard of response required by a candidate as exemplified by the standardisation scripts.

GENERIC MARKING PRINCIPLE 2:

Marks awarded are always whole marks (not half marks, or other fractions).

GENERIC MARKING PRINCIPLE 3:

Marks must be awarded **positively**:

- marks are awarded for correct/valid answers, as defined in the mark scheme. However, credit is given for valid answers which go beyond the scope of the syllabus and mark scheme, referring to your Team Leader as appropriate
- marks are awarded when candidates clearly demonstrate what they know and can do
- marks are not deducted for errors
- marks are not deducted for omissions
- answers should only be judged on the quality of spelling, punctuation and grammar when these features are specifically assessed by the question as indicated by the mark scheme. The meaning, however, should be unambiguous.

GENERIC MARKING PRINCIPLE 4:

Rules must be applied consistently, e.g. in situations where candidates have not followed instructions or in the application of generic level descriptors.

GENERIC MARKING PRINCIPLE 5:

Marks should be awarded using the full range of marks defined in the mark scheme for the question (however; the use of the full mark range may be limited according to the quality of the candidate responses seen).

GENERIC MARKING PRINCIPLE 6:

Marks awarded are based solely on the requirements as defined in the mark scheme. Marks should not be awarded with grade thresholds or grade descriptors in mind.

Mathematics Specific Marking Principles

- 1 Unless a particular method has been specified in the question, full marks may be awarded for any correct method. However, if a calculation is required then no marks will be awarded for a scale drawing.
- 2 Unless specified in the question, non-integer answers may be given as fractions, decimals or in standard form. Ignore superfluous zeros, provided that the degree of accuracy is not affected.
- 3 Allow alternative conventions for notation if used consistently throughout the paper, e.g. commas being used as decimal points.
- 4 Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored (isw).
- 5 Where a candidate has misread a number or sign in the question and used that value consistently throughout, provided that number does not alter the difficulty or the method required, award all marks earned and deduct just 1 A or B mark for the misread.
- 6 Recovery within working is allowed, e.g. a notation error in the working where the following line of working makes the candidate's intent clear.

Abbreviations

cao	correct answer only
dep	dependent
FT	follow through after error
isw	ignore subsequent working
oe	or equivalent
SC	Special Case
nfww	not from wrong working
soi	seen or implied
	-

Question	Answer	Marks	Partial Marks
1(a)	28 028	2	B1 for 8372 seen or M1 for $36400 - \frac{23}{100} \times 36400$ oe
1(b)	2670	2	M1 for $\frac{890}{5} [\times k]$ oe where $k \ge 1$
1(c)(i)	3.5	2	M1 for $\frac{702800 - 678202}{702800} [\times 100]$ oe or $\frac{678202}{702800} \times 100$ oe
1(c)(ii)	627 500	2	M1 for $\frac{(100+12)}{100}x = 702800$ soi
1(d)(i)	1.148×10^6 cao	1	
1(d)(ii)	Population densities 1100 or 1101.[] 2030 or 2026.[] 1320 or 1315.[] 510 or 509.6 seen	M2	M1 for one correct population density seen
	Bahrain	B1	
2(a)(i)	<i>x</i> = 2	1	
2(a)(ii)	Correct reflection (5, 2), (5, 4), (6, 4)	1	
2(b)	Rotation 90° anticlockwise oe (0, 0) oe	3	B1 for each
3(a)(i)	$\frac{1}{4}$ oe	1	
3(a)(ii)	$(0, -\frac{5}{4})$ oe	1	

Cambridge O Level – Mark Schewww.dynamicpap/ersecoden 2023 PUBLISHED

Question	Answer	Marks	Partial Marks
3(b)(i)	6.71 or 6.708	2	M1 for $(42)^2 + (5 - 8)^2$ oe
3(b)(ii)	$[y=]-\frac{1}{2}x+7$ oe final answer	3	M1 for $\frac{8-5}{-2-4}$ oe
			M1 for attempt to find c by substitution of (4, 5) or (-2, 8) or (1, 6.5) into $y = (their m)x + c$ or use of $y - b = (their m)(x - a)$
4(a)(i)	40	1	
4(a)(ii)	7n-2 oe final answer	2	B1 for $7n + j$ oe seen, j may be 0
4(a)(iii)	50	3	M1 for correct equation for sum of terms FT use of <i>their</i> $7n-2$
			M1 for correct rearrangement to $ak = b$, dependent on equation with two terms in <i>k</i> and a constant Not $k + k + 1 = 703$ oe
4(b)	58	5	B4 for $a = 4$ and $b = -2$
			OR
			B2 for $1^2 + a + b = 3 + a + b = 3$ or better and $3^2 + 3a + b = 19 + 3a + b = 19$ or better or B1 for one correct
			M1 for correct method to eliminate one variable from <i>their</i> simultaneous equations
			M1 for 6^2 + <i>their a</i> × 6 + <i>their b</i>
			Alternative method
			B3 for second term = 10
			OR
			B1 for second difference = 2
			B2 for first pair of differences 7 and 9 or B1 for sum of first pair of differences = 16
			AND
			B1 for next first differences 11, 13, 15
5(a)(i)	46	1	
5(a)(ii)	[0]8 15	1	

Cambridge O Level – Mark Scheweww.dynamicphp@rscoden2023 PUBLISHED

Question	Answer	Marks	Partial Marks
5(b)(i)	55 to 57	1	
5(b)(ii)	30 nfww	2	B1 for 54 or 24 seen
5(b)(iii)	17.5	3	B1 for 132 or 28 M1 for $\frac{160 - their \ 132}{160} [\times 100]$ or $\frac{their \ 28}{160} [\times 100]$ or $\frac{their \ 132}{160} \times 100$
5(c)	50.3 or 50.34	3	B1 for 36.5 and 43.5 seen M1 for $\frac{their \ 36.5}{their \ 43.5} \times 60$ oe
6(a)	$\frac{30}{x}$	1	
6(b)	Correct unexpanded expression for shaded area e.g. $[y=]30-(x-2)\left(\frac{30}{x}-2\right)$ or $[y=]2(x-2)+2\times2+2\left(\frac{30}{x}-2\right)$ or $[y=]2x+2\left(\frac{30}{x}-2\right)$ or $[y=]2(x-2)+2\left(\frac{30}{x}\right)$ Correct expansion of brackets and rearrangement to $y=2x+\frac{60}{x}-4$	M2 A1	B1 for $(x-2)$ or $\left(their \frac{30}{x} - 2\right)$ seen $A = \begin{bmatrix} E \\ F \\ F \end{bmatrix} = \begin{bmatrix} D \\ G \\ C \end{bmatrix}$ A0 if any errors or omissions
6(c)	28.3	1	
6(d)	Correct smooth curve	3	B2 FT for 7 or 8 points correctly plotted or B1 FT for 5 or 6 points correctly plotted
6(e)	FT reading the two values from <i>their</i> graph at $y = 24$	2	B1 FT for one correct B1 FT for second value = $30 \div their$ first value or for <i>their</i> second value at $y = 24$
7(a)	5a-3b final answer	2	B1 for answer $5a + kb$ or $ka - 3b$ $k \neq 0$ or for $5a - 3b$ seen
7(b)	11x+1 final answer	2	B1 for answer $11x + k$ or $kx + 1$ $k \neq 0$ or M1 for $6x - 9 + 5x + 10$ seen

Cambridge O Level – Mark Schewww.dynamicpap/ersecoden 2023 PUBLISHED

Question	Answer	Marks	Partial Marks
7(c)	$\frac{-(-2)\pm\sqrt{(-2)^2-4\times6\times-9}}{2\times6} \text{ or }$ $\frac{-(-2)}{2\times6}\pm\sqrt{\left(\frac{-2}{2\times6}\right)^2-\left(\frac{-9}{6}\right)}$	B2	B1 for $\sqrt{(-2)^2 - 4 \times 6 \times -9}$ oe or for $\frac{-(-2) \pm \sqrt{their \text{ discriminant}}}{2 \times 6}$ or for $\left(x + \frac{-2}{2 \times 6}\right)^2$
	-1.07 and 1.40	B1	
7(d)(i)	$\frac{xy}{8}$ final answer	1	
7(d)(ii)	$\frac{38-x}{(2x+1)(x-5)} \text{ or } \frac{38-x}{2x^2-9x-5}$ final answer	3	B1 for $3(2x+1)-7(x-5)$ oe isw B1 for denominator $(2x+1)(x-5)$ oe isw
8(a)(i)	1	1	
8(a)(ii)	2	1	
8(a)(iii)	1.95	2	$\frac{\mathbf{M1} \text{ for}}{\left(\begin{bmatrix} 0 \times 7 \end{bmatrix} + 1 \times 11 + 2 \times 9 + 3 \times 5 + 4 \times 6 + 5 \times 2 \right)}{40} \text{ oe}$
8(a)(iv)	$\frac{1}{5}$ cao	2	M1 for $\frac{6+2}{40}$ oe
8(b)(i)	$\frac{21}{50}$ oe	2	M1 for $\frac{7}{10} \times \frac{3}{10} [\times 2]$ oe
8(b)(ii)	$\frac{7}{40}$ oe	3	M2 for $\frac{3}{10} \times \frac{2}{9} \times \frac{7}{8} \times 3$ oe or M1 for $\frac{3}{10} \times \frac{2}{9} \times \frac{7}{8}$ oe If 0 scored, SC1 for answer $\frac{189}{1000}$ oe
9(a)(i)	688	B1	
	cm ³	B1	
9(a)(ii)	$\frac{34.4}{\frac{1}{2} \times 6 \times \sin 55}$	M2	M1 for $\frac{1}{2} \times 6 \times AC \times \sin 55 = 34.4$
	13.998[] or 14.00	A1	

Cambridge O Level – Mark Scheweww.dynamicpap/ersecoden 2023 PUBLISHED

Question	Answer	Marks	Partial Marks
9(a)(iii)	702 or 701.6 to 701.8	5	B2 for $[BC^2 =]135.4$ to 135.6 soi or M2 for $\sqrt{6^2 + 14^2 - 2 \times 6 \times 14\cos 55}$ or M1 for $6^2 + 14^2 - 2 \times 6 \times 14\cos 55$ AND M2 for $2 \times 34.4 + 6 \times 20 + 14 \times 20 + their BC \times 20$ oe or M1 for sum of areas of 4 faces from the M2 calculation
9(b)	9.41 or 9.407	4	M1 for area of base $=\frac{98}{8}$ soi M2 for <i>their</i> 12.25 + <i>their</i> 12.25 + 8 ² oe or M1 for <i>their</i> 12.25 + <i>their</i> 12.25 oe or <i>their</i> 12.25 + 8 ² oe
10(a)	AO = DO [equal] radii BO = CO [equal] radii $\angle AOB = \angle DOC$ [vertically] opposite [Congruent] SAS	3	B2 for two pairs of equal sides/angles with correct reasons or B1 for one pair of equal sides/angles with correct reason or for two appropriate pairs with no or incorrect reasons Angles described using convention of 3 letters

Cambridge O Level – Mark Schewww.dynamicpap/ersecoden 2023 PUBLISHED

Question	Answer	Marks	Partial Marks
10(b)	4.39 or 4.40 or 4.392 to 4.398		M2 for $[AOB =]2 \times \sin^{-1}\left(\frac{4.5}{5}\right)$ oe or $\cos[] = \frac{5^2 + 5^2 - 9^2}{2 \times 5 \times 5}$ oe or M1 for $\sin[] = \frac{4.5}{5}$ oe or $9^2 = 5^2 + 5^2 - 2 \times 5 \times 5 \cos[]$ oe AND M2 for $10\pi - \left(9 + 9 + 2 \times \frac{their AOD}{360} \times 10\pi\right)$ oe or M1 for $[2 \times] \frac{their AOD}{360} \times 10\pi$ oe