

UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS General Certificate of Education Ordinary Level

CANDIDATE NAME			
 CENTRE NUMBER		CANDIDATE NUMBER	
ADDITIONAL MATHEMATICS			4037/22
Paper 2			May/June 2013
			2 hours
Candidates answer on the Question Paper.			
No Additional Materials are required.			

# **READ THESE INSTRUCTIONS FIRST**

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use a pencil for any diagrams or graphs.

Do not use staples, paper clips, highlighters, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

Answer all the questions.

Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question. The use of an electronic calculator is expected, where appropriate. You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together. The number of marks is given in brackets [] at the end of each question or part question. The total number of marks for this paper is 80.

This document consists of **15** printed pages and **1** blank page.



### Mathematical Formulae

#### 1. ALGEBRA

Quadratic Equation

For the equation  $ax^2 + bx + c = 0$ ,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \cdot$$

Binomial Theorem

$$(a+b)^n = a^n + \binom{n}{1}a^{n-1}b + \binom{n}{2}a^{n-2}b^2 + \dots + \binom{n}{r}a^{n-r}b^r + \dots + b^n,$$
  
where *n* is a positive integer and  $\binom{n}{r} = \frac{n!}{(n-r)!r!}$ .

## 2. TRIGONOMETRY


Identities

$$\sin^2 A + \cos^2 A = 1$$
$$\sec^2 A = 1 + \tan^2 A$$
$$\csc^2 A = 1 + \cot^2 A$$


Formulae for  $\triangle ABC$ 

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$
$$a^2 = b^2 + c^2 - 2bc \cos A$$
$$\Delta = \frac{1}{2} bc \sin A$$


Examiner's Use



Variables x and y are such that when  $\sqrt{y}$  is plotted against  $x^2$  a straight line graph passing through the points (1, 3) and (4, 18) is obtained. Express y in terms of x. [4]



(b) Express 
$$\frac{y \times (4x^3)^2}{\sqrt{8y^3}}$$
 in the form  $2^a \times x^b \times y^c$ , where *a*, *b* and *c* are constants. [3]



The diagram shows the graph of a function y = f(x). On the same axes sketch the graph of  $y = f^{-1}(x)$ . [2]

3

- 4 The position vectors of the points A and B, relative to an origin O, are 4i - 21j and 22i - 30jrespectively. The point C lies on AB such that  $A\hat{B} = 3A\hat{C}$ . Examiner's Use
  - (i) Find the position vector of C relative to O.

(ii) Find the unit vector in the direction  $\overrightarrow{OC}$ .


[2]

For

[4]

For Examiner's Use


### 5 Calculators must not be used in this question.



The diagram shows a trapezium *ABCD* in which AD = 7 cm and  $AB = (4 + \sqrt{5})\text{ cm}$ . *AX* is perpendicular to *DC* with DX = 2 cm and XC = x cm. Given that the area of trapezium *ABCD* is  $15(\sqrt{5} + 2)\text{ cm}^2$ , obtain an expression for x in the form  $a + b\sqrt{5}$ , where a and b are integers. [6]

Examiner's Use






The shaded region in the diagram is a segment of a circle with centre O and radius r cm.

Angle  $AOB = \frac{\pi}{3}$  radians.


(i) Show that the perimeter of the segment is  $r\left(\frac{3+\pi}{3}\right)$  cm. [2]

(ii) Given that the perimeter of the segment is 26 cm, find the value of r and the area of the segment. [5]



# www.dynamicpapers.com

[2]



10

(ii) Given that angle  $ABC = 90^\circ$ , find the equation of *BC*.

(iii) Given that the length of AC is 10 units, find the coordinates of each of the two possible positions of point C. [4]

For Examiner's Use

Use

9 (a) The graph of  $y = k(3^x) + c$  passes through the points (0, 14) and (-2, 6). Find the value of k and of *c*. [3] Examiner's

(b) The variables x and y are connected by the equation  $y = e^x + 25 - 24e^{-x}$ .

- (i) Find the value of y when x = 4. [1]
- (ii) Find the value of  $e^x$  when y = 20 and hence find the corresponding value of x. [4]

(a) The function f is defined, for  $0^{\circ} \le x \le 360^{\circ}$ , by  $f(x) = 1 + 3\cos 2x$ . 10 For Examiner's Use (i) Sketch the graph of y = f(x) on the axes below. [4] y 360° x 0 180° 270° 90° -5State the amplitude of f. **(ii)** [1] State the period of f. [1] (iii) (b) Given that  $\cos x = p$ , where  $270^\circ < x < 360^\circ$ , find  $\csc x$  in terms of p. [3]

11 A curve has equation 
$$y = 3x + \frac{1}{(x-4)^3}$$
.  
(i) Find  $\frac{dy}{dx}$  and  $\frac{d^2y}{dx^2}$ .  
[4]

(ii) Show that the coordinates of the stationary points of the curve are (5, 16) and (3, 8). [2]

(iii) Determine the nature of each of these stationary points.

[2]

Examiner's

Use

[2]

(iv) Find 
$$\int \left( 3x + \frac{1}{(x-4)^3} \right) dx$$
.

(v) Hence find the area of the region enclosed by the curve, the line x = 5, the x-axis and the line x = 6. [2]

**BLANK PAGE** 

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.