www.dynamicpapers.com

Cambridge Assessment International Education

Cambridge O Level

CANDIDATE NAME									
CENTRE NUMBER						CANDIDATE NUMBER				

4855957787 5

ADDITIONAL MATHEMATICS **4037/14**

Paper 1 **May/June 2021**

2 hours

You must answer on the question paper.

No additional materials are needed.

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 80.
- The number of marks for each question or part question is shown in brackets [ ].

This document has **16** pages.

© UCLES 2021

www.dynamicpapers.com

Mathematical Formulae

1. ALGEBRA

Quadratic Equation

For the equation $ax^2 + bx + c = 0$,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Binomial Theorem

$$(a+b)^n = a^n + \binom{n}{1}a^{n-1}b + \binom{n}{2}a^{n-2}b^2 + \ldots + \binom{n}{r}a^{n-r}b^r + \ldots + b^n$$

where n is a positive integer and $\binom{n}{r} = \frac{n!}{(n-r)!r!}$

Arithmetic series $u_n = a + (n-1)d$

$$S_n = \frac{1}{2}n(a+l) = \frac{1}{2}n\{2a + (n-1)d\}$$

Geometric series $u_n = ar^{n-1}$

$$S_n = \frac{a(1-r^n)}{1-r} \quad (r \neq 1)$$

$$S_\infty = \frac{a}{1-r} \quad (|r| < 1)$$

2. TRIGONOMETRY

Identities

$$\sin^2 A + \cos^2 A = 1$$
$$\sec^2 A = 1 + \tan^2 A$$
$$\operatorname{cosec}^2 A = 1 + \cot^2 A$$

Formulae for ΔABC

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$


$$a^2 = b^2 + c^2 - 2bc\cos A$$

$$\Delta = \frac{1}{2}bc\sin A$$

© UCLES 2021


www.dynamicpapers.com

1 **(a)** On the axes below, sketch the graph of $y = 6\cos 2x - 1$ for $0° \leqslant x \leqslant 360°$.



[3]

(b) The graph of $y = a + b\sin c\theta$ for $-180° \leqslant \theta \leqslant 180°$ is shown below.




Write down the value of each of the constants a, b and c. [2]

$a =$ ..... $b =$ ..... $c =$ .....

www.dynamicpapers.com

2 **(a)** On the axes below, sketch the graphs of $y = |x-3|$ and $y = \left|\frac{2}{5}x\right|$, giving the coordinates of the points where the graphs meet the axes. [3]



(b) Solve the equation $\left|\frac{2}{5}x\right| = |x-3|$. [2]

© UCLES 2021

www.dynamicpapers.com

3 **(a)** Find the first 3 terms in the expansion, in ascending powers of x, of $(a-3x)^{10}$, where a is a constant. [3]

(b) Given that a is positive and that the three terms found in **part (a)** can also be written as $p+qx+\frac{405}{256}x^2$, find the value of each of the constants a, p and q. [3]

www.dynamicpapers.com


4 **(a)** Find $\frac{\mathrm{d}}{\mathrm{d}x}(2x+1)^{\frac{5}{2}}$. [2]

(b) Hence find $\int (2x+1)^{\frac{3}{2}}\,\mathrm{d}x$. [2]

© UCLES 2021

www.dynamicpapers.com

(c)



The diagram shows the graph of the curve $y = (2x+1)^{\frac{3}{2}}$ for $x \geqslant 0$. The shaded region enclosed by the curve, the axes and the line $x = a$ is equal to 48.4 square units. Find the value of a, showing all your working. [3]

www.dynamicpapers.com

5 **(a)** A 5-digit number is to be formed from the digits 2, 5, 6, 7 and 9. Each digit may only be used once.

(i) Find the number of different 5-digit numbers that can be formed. [1]

(ii) Find the percentage of these numbers that are odd. [2]

(b) 12 people are placed at random in 3 groups of 4 people each. Find the number of ways that this can be done. [3]

© UCLES 2021

www.dynamicpapers.com

6 **(a)** Solve the simultaneous equations

$$\log_a(x+y) = 0,$$
$$\log_a(x+1) = 2\log_a y.$$ [4]

(b) Given that $\log_p q^2 \times \log_q p^3 = A$, find the value of the constant A. [3]

www.dynamicpapers.com

7 A curve is such that $\frac{d^2y}{dx^2} = 8\sin 2x$. The curve has a gradient of 6 at the point $\left(\frac{\pi}{2}, 4\pi\right)$.

Find the equation of the curve. [8]

© UCLES 2021

www.dynamicpapers.com


8 The polynomial $\mathrm{p}(x)$ is $ax^3 + bx^2 + 7x + 1$, where a and b are integers. It is given that $2x + 1$ is a factor of $\mathrm{p}(x)$ and that when $\mathrm{p}(x)$ is divided by $x - 3$ there is a remainder of 175.

(a) Find the value of a and of b. [5]

(b) Using your values of a and b from **part (a)**, find the remainder when $\mathrm{p}'(x)$ is divided by $x - 1$. [3]

www.dynamicpapers.com

9 In this question all lengths are in centimetres.
Do not use a calculator in this question.



The diagram shows the trapezium $ABCD$, where $AB = 5\sqrt{3}-3$, $DC = 5\sqrt{3}-1$ and $AD = \sqrt{3}+2$.

(a) Find the area of $ABCD$, giving your answer in the form $a+b\sqrt{3}$, where a and b are integers. [3]

© UCLES 2021


www.dynamicpapers.com

(b) Given that angle $BCD = \theta$ radians, find the value of $\cot\theta$ in the form $c + d\sqrt{3}$, where c and d are integers. [3]

(c) Using your answer to **part (b)**, find the value of $\text{cosec}^2\theta$ in the form $e + f\sqrt{3}$, where e and f are integers. [2]

www.dynamicpapers.com

10



The diagram shows the figure $ABCDEFG$, where $ABFG$ and $BCDE$ are rectangles of length x cm and width $\frac{x}{3}$ cm. The sector BFE of the circle, centre B, radius x cm, has an angle of θ radians. It is given that the area of BFE is $2\,\text{cm}^2$.

(a) Show that the perimeter, P cm, of the figure $ABCDEFG$ is given by $P = \frac{10x}{3} + \frac{4}{x}$. [5]

www.dynamicpapers.com

(b) Given that x can vary, find the minimum value of P in the form $q\sqrt{30}$, where q is a rational number. [4]

(c) Verify that P is a minimum. [1]

Question 11 is printed on the next page.

www.dynamicpapers.com

11 The tangent at the point where $x = 1$ on the curve $y = 6x\ln(x^2 + 1)$ intersects the y-axis at the point P. This tangent also intersects the line $x = 2$ at the point Q. A line through P, parallel to the x-axis, meets the line $x = 2$ at the point R. Find the exact area of triangle PQR. [10]

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.

© UCLES 2021