#### UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS

General Certificate of Education O Level

#### MARK SCHEME for the June 2004 question papers

	5054 PHYSICS
5054/01	Paper 1 (Multiple Choice), maximum mark 40
5054/02	Paper 2 (Theory), maximum mark 75
5054/03	Paper 3 (Practical Test), maximum mark 30
5054/04	Paper 4 (Alternative to Practical), maximum mark 30

These mark schemes are published as an aid to teachers and students, to indicate the requirements of the examination. They show the basis on which Examiners were initially instructed to award marks. They do not indicate the details of the discussions that took place at an Examiners' meeting before marking began. Any substantial changes to the mark scheme that arose from these discussions will be recorded in the published Report on the Examination.

All Examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes must be read in conjunction with the question papers and the Report on the Examination.

CIE will not enter into discussion or correspondence in connection with these mark schemes.

CIE is publishing the mark schemes for the June 2004 question papers for most IGCSE and GCE Advanced Level syllabuses.



GCE O Level

MARK SCHEME

MAXIMUM MARK: 40

### SYLLABUS/COMPONENT: 5054/01

PHYSICS Paper 1 (Multiple Choice)



#### www.dynamicpapers.com

Page 1	Mark Scheme	Syllabus	Paper
	PHYSICS – JUNE 2004	5054	1

Question Number	Key	Question Number	Key
1	Α	21	Α
2	Α	22	D
3	С	23	С
4	D	24	Α
5	С	25	С
6	D	26	В
7	С	27	В
8	Α	28	D
9	D	29	В
10	Α	30	Α
11	С	31	В
12	В	32	D
13	Α	33	D
14	Α	34	В
15	D	35	С
16	В	36	Α
17	В	37	Α
18	В	38	D
19	D	39	Α
20	С	40	В

TOTAL 40

GCE O Level

MARK SCHEME

# MAXIMUM MARK: 75

## SYLLABUS/COMPONENT: 5054/02

PHYSICS Paper 2 (Theory)



			www.dynamicpape	ers.com	
	Page	e 1	Mark Scheme Syllabus	Paper	
			PHYSICS – JUNE 2004 5054	2	
Se	ectior	ηA			
1	(a)	(i)	weight / gravity / gravitational (force)		B1
		(ii)	air / wind resistance or drag or friction / upthrust		B1
	(b)	(i) (ii)	e.g. resistance opposes gravity or decreases acc.	C1 C1 C1	B1 B2
		(iii)	air resistance = weight / no resultant / net / overall force / downw	wards	
			force balances upwards force	Total	B1 <b>[6]</b>
2	(a) (b)	(i) (ii)	radiation no molecules or medium (to vibrate, conduct, convect) / vacuum <b>hot air</b> rises (hot) air expands / density decreases		B1 B1 B1 B1
	(c)		fiberglass or air is a bad conductor/ insulator / lags / reduces heat fiberglass traps air or prevents convection (ignore radiation statements)	flow	B1 B1
				Total	[6]
3	(a) (b)		rise in temperature / hot / heated road / bridge / rail / metal expands or gap reduces no buckling / deformation / breaking / cracking / twisting / tilting any other problem + solution e.g. concrete cracks – leave a gap, telephone wires sag – put them high / tight hot water cracks glass – use thin glass / car engines seize up – cool them water freezes in pipes – lag them or use antifreeze / tyres burst – let air out pipes bend – use flexible joints / dashboard deforms – car in shade wrong readings on measuring cylinder – use correct temp.		B1 B1 B1
				Total	[4]
	(-)				[-1]
4	(a)		distance traveled per unit time <b>or</b> in one second / distance ÷ time <b>or</b> rate of change of distance		B1
	(b)		s = d/t in any algebraic or numerical form any doubling of distance or final time		C1 C1
			0.48 s (allow 0.24s 2/3 accept 0.5s)		A1
	(c)		60/0.48 (5) 123.75 accept 120, 123, 124 (ecf (b))		C1 A1
				Total	[6]
5	(a)	(i)	magnetic (field) of current / coil / recording head		
		(ii)	or head is magnetized / an electromagnet magnetism / magnetic field or current or poles on head reverses /	,	B1
		(iii)	changes direction (accept "due to alternating current") each direction / one cycle longer (on tape)		B1 B1
	(b)	(ii) (i) (ii)	need to keep record / tape stored or played iron, steel etc	Total	B1 B1
				rotai	[5]

			www.dynamicpapers.co	m
	Page	e 2	Mark Scheme Syllabus Paper	
			PHYSICS – JUNE 2004 5054 2	
6	(a)	(i) (ii)	voltage past maximum or 3V / off scale / outside range reading less accurate or sensitive / not far up scale or smaller deflection	B1
	(b)	(i)	V = I R in any algebraic format 4/12 0.33 A (accept 1/3 A)	B1 C1 A1
		(ii)	(i) * 30 or (i) * 18 + 4 or 30*4/12 9.9 - 10 V (e.c.f (i), e.g. if (i) = 0.3, 0.3*30 = 9V or 0.3*18+4 = 9.4 V) only 1 unit error in this question	C1 A1
			Tota	I [7]
7	(a)	(i)	filament is hot / heated (by current from 6V supply) / thermionic emission	B1
		(ii)	anode is positive / anode attracts electrons / electrons attracted to + (electric) field from anode to cathode	B1
		(iii)	otherwise electrons stopped / deflected / slowed down / Collide (with air atoms) (accept no opposition to movement, to reach screen, to avoid air resistance)	B1
	(b)		up and down vertical <b>or</b> side to side movement ( <b>not</b> on both axes) electrons deflected by electric field <b>or</b> attracted to + or repelled by –	B1
			or plates are charged (e.g. plates are +ve and –ve)	B1 I [5]
8	(a) (b)		radon (gas) cancer / mutation / <b>cell</b> damage <b>or</b> death	B1
	()		radiation sickness or adds to readings (accept count with no source)	B1
	(c) (d) (e)		(outer) space / stars / Sun (not sunlight) number of protons and neutrons (not no. nucleons) 84 216 (values reversed B1) Tota	B1 B1 B2 I [6]

]
I
(
I
[
ſ
N
A
/ E
E
 (
Č
4
(
(
( ( ( /

Total [15]

			www.dy	namicpap	pers.com	
I	Page 4		Mark Scheme	Syllabus	Paper	
			PHYSICS – JUNE 2004	5054	2	
11	(a)	(i)	P.E. decreases (A to B or C to D or downhill or	r initially)		B1
			K.E. gained (P.E. $\rightarrow$ K.E2)			B1
			K.E. to P.E. <b>change</b> must be clear and from B to	C or uphill		B1
		(ii)	mgh algebraic form seen			C1
			500*10*30			C1
		/····	150 000 J			A1
		(iii)	conservation of energy cited <b>or</b> clear that loss of I	P.E.		<u> </u>
			has become K.E. 500*10*20 or 500*10*10 or 50 000 seen			C1 C1
			100 000 J (allow g=9			A1
				5.0)		
	(b)	(i)	velocity involves direction <b>or</b> is a vector (speed do direction (of carriage) changes / carriage turns	Des not) (accept on diag	ram)	B1
		(ii)	force towards centre (of curve) / inwards	(accept centrip	,	B1
	(-)					01
	(c)		F = ma in any algebraic form or 3000 = 500a 3000/500 6(.0) m/s <sup>2</sup>			C1 C1 A1
					Total	[15]
			Total fe	or paper :		[75]

GCE O Level

MARK SCHEME

MAXIMUM MARK: 30

## SYLLABUS/COMPONENT: 5054/03

PHYSICS Paper 3 (Practical Test)



Pa	ige 1	Mark Scheme	Syllabus	Paper
1 4		PHYSICS – JUNE 2004	5054	3
L	I	11110100 - 0011L 2004	5054	5
1.	(a), (b) & (c)	Repeat measurements taken for either $t_1$ or $t_2$ .		B1
		Correct $T_1$ in the range 1.40 s to 1.60 s to 0.01	1 s	B1
		Correct $T_2$ within $\pm 0.1$ s of $T_1$		<b>B</b> 1
	(d)	Comment on Either reaction time – however expressed Or range of values		B1
	(e) Or	Sensible conclusion based on their results e.g Time for one oscillation is independent of the (if periods are the same within the limits of ur Time for one oscillation increases / decreases in mass. (Allow direct or inverse proportion)	e mass. ncertainty)	2
		(provided their results show this)		B1
			Tota	al [5]
2.	(a)	Power supply, ammeter and switch in series we between A and B, voltmeter in parallel with p		B1
	(b), (c) & (d)	<i>I</i> values in region of 0.3 A and 0.45 A with unleast once and at least one current to 0.01 A. (Allow Centre variation)	nit seen at	B1
		Both V values in the region of 4.5 V with unit once and at least one voltage to 0.1 V. (Allow Centre variation)		B1
		<i>R</i> values in the region of 15 $\Omega$ and 10 $\Omega$ with least once.	unit seen at	B1
	(e)	Resistance increases as diameter decreases. (Allow resistance is inversely proportional to	diameter or a	B1 urea)
			Tota	1 [5]

Total [5]

VANANAL dy	ynamicpapers.com
<b>vvvvv</b>	

		www.dynamicp	papers.con
P	Page 2	Mark Scheme Syllab	us Paper
		PHYSICS – JUNE 2004 5054	3
3.	(a) and	(b) Sensible temperatures with unit seen at least once.	B1
		At least one reading attempted to better than 1 $^\circ\text{C}$	B1
		$V_{\rm F}$ numerically to (1.0 to 3.0) x temperature drop and correct calculation of $V_{\rm I}$ with unit seen at least or	nce. B1
		$m_{\rm I}$ numerically equal to $V_{\rm I}$ .	DI
	(c) and	(d) Sensible values for all the thermal energy changes wi unit seen at least once.	th M1
	(e)	Energy gained greater than energy lost as cold water thermal energy from beaker / surroundings	gains A1
			Total [5]
4. <u>In</u>	nitial read	ings.	
	(b)	$x 0.60 \pm 0.05$ m with unit.	B1
	(c)	$y 0.20 \pm 0.05$ m with unit.	B1
		(Penalise missing unit once only) x and y recorded to 0.001 m or better.	B1
<u>T</u>	<u>[able</u> ]		
	(d)	Table with units for $d$ , $D$ and $1/D$ .	B1

At least one reading with D greater than or equal to 1.00 m. B1

At least one reading with D less than or equal to 0.70 m **B**1

Correct calculation of  $(d/D)^2$  and 1/D to at least 2 s.f. **B**1

Five good values judged according to the table below. B1

D	Range of $(d/D)^2$	1 / D
0.65	0.06 - 0.10	1.54
0.70	0.12-0.16	1.43
0.75	0.18-0.22	1.33
0.80	0.23 - 0.27	1.25
0.85	0.27 - 0.31	1.18
0.90	0.31 - 0.35	1.11
0.95	0.35 - 0.39	1.05
1.00	0.38 - 0.42	1.00

Page 3	Mark Scheme	Syllabus	Paper
	PHYSICS – JUNE 2004	5054	3
raph.			
(e)	Axes labelled with unit and correct orientation		B1
	Suitable scale y axis 1 cm = $0.02 / 0.025$ x axis 1 cm = $0.1$ or $0.05$ m <sup>-1</sup>		M1
	Two points plotted correctly – check the two p from the line.	oints furth	est A1
	Best fit fine line and finely plotted points.		B1
Calculations.			
(f) and (g)	Large triangle.		B1
	Correct calculation of $S$ and $f$ (ignore sign)		B1
	Value of $f$ in range 0.130 m to 0.170 m with un	nit.	B1
			Total

GCE O Level

MARK SCHEME

MAXIMUM MARK: 30

## SYLLABUS/COMPONENT: 5054/04

PHYSICS (Alternative to Practical)



Page 1	Mark Scheme	Syllabus	Paper
	PHYSICS – JUNE 2004	5054	4

#### Question 1

<ul> <li>(a) Uses two rays from X and Y (clear <u>intention</u> to touch hole edges) One X and one Y ray "touch" an edge of the hole and meet screen Any one X and one Y are neat lines (rule and sharp "pencil") allow apparent "refraction" or "diffraction" at hole One correct X and the corresponding Y labeled on screen Arrows on rays; no broken lines penalty -1 (max).</li> </ul>			
		(b) XY in range 54 to 56 mm (unit required), accept in cm	B1
		Tota	ıl [5]
Question 2			
<ul> <li>(a) 4 items correct, 3mks; 3 items = 2mks; 2 items = 1mk. Accept historical symb Accept any other component provided that the function of the circuit is not compromised.</li> <li>Penalise -1 (max) :- short circuit (e.g. line behind component, unless signs of of rubber) or any compromised circuit function.</li> </ul>			
(b) Correct polarities, +ve signs for correct terminals of cell and ammeter (re diode).	B1		
<ul> <li>(c) No current / I = 0, (do not accept "nothing"), accept very small "reverse" curre</li> <li>/ lamp does not light.</li> </ul>	nt B1		
(d) One from: limit current / prevent overheating / current indicator / provides resistance	B1		
Question 3	ıl [6]		
(a) Any method <u>based</u> on rule reading at 25°C – rule reading at top of			
thermometer bulb. NB / required. Mark text or diagram or <u>Fig 3.1</u> Bula as close as passible to thermometer (on diagram < 1 cm) (	B1		
Rule as close as possible to thermometer (on diagram < 1 cm) / uses fiducial aid	B1 B1		
With the eye/line of sight perpendicular to the rule/end of mercury thread			
<ul> <li>(b) (i) I<sub>0</sub> = 5.6 - 5.8 (cm), I<sub>100</sub> = 22.6 - 22.8 (cm) ignore unit</li> <li>(ii) ΔI / 100, clear, correct arithmetic ecf, 2 or 3 dcp, ignore unit, accept any correct ΔI / Δθ from graph.</li> <li>(iii) linearly, or (I - I<sub>0</sub>) ∝ θ accept/line has a constant/uniform m, note that "directly proportional" automatically looses the mark.</li> </ul>			
		▼ - 4 -	1 101

Total [6]

VANANA/ dy	ynamicpapers.com
	VNANICOADEIS COM
** ** ** . M	

Page 2	Mark Scheme	Syllab	us Pape	r
	PHYSICS – JUNE 2004	5054	4	

#### Question 4

(a) (i) V initial = a volume between 40cm <sup>3</sup> and 60cm <sup>3</sup> : (allow use of beaker) must be able to displace 40cm <sup>3</sup> / prevents overflowing /	
exceeding cm <sup>3</sup> limit	B1
<ul> <li>(ii) {V<sub>max</sub> - V<sub>initial</sub>} / change in volume is found / change in volume obtained is</li> <li>= V<sub>metal</sub> / any related answer that has an association of measurement of volume.</li> </ul>	B1
(iii) Any good point e.g. tap cylinder to release air / how avoiding parallax / water at 20°C / careful pouring / avoid splashing / use set square / repeat	
average / reading the position of the bottom of the meniscus. (b) Scale calibration of cylinder is correct at 20°C / liquid needs to be at 20°C	B1 B1
(c) Water (on the metal would be) included in the (repeat) volume of the metal; or something that means the same, not just erroneous.	B1
Total	[6]
Question 5	
(a) Axes correct, scale that cannot be x2 / is not "awkward" and with units Correct plotting, nearest ½ small square, check first point and obvious	B1
plot errors.	B1
Line judgement re plots (line does not go through all correctly plotted points,	D1
so accept smooth line through 5 points i.e., one point not on the line) Neat smooth thin line	B1 B1
<b>(b)</b> Mark cands diagram <u>or Fig 5.1:</u>	
(i) Object displace downwards OR screen displaced downwards Any ray from the top of object through the lens to meet screen.	B1
Be generous re art and accuracy of position,	B1 B1
(ii) put centres in line	וס
Tota	I [7]

Paper Total 30

www.dynamicpapers.com