



UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS  
General Certificate of Education Ordinary Level

CANDIDATE NAME

CENTRE NUMBER 

--	--	--	--	--

CANDIDATE NUMBER 

--	--	--	--



**CHEMISTRY** **5070/32**  
Paper 3 Practical Test **October/November 2013**  
**1 hour 30 minutes**

Candidates answer on the Question Paper.  
Additional Materials: As listed in the Confidential Instructions

**READ THESE INSTRUCTIONS FIRST**

Write your Centre number, candidate number and name on all the work you hand in.  
Write in dark blue or black ink.  
You may use a soft pencil for any diagrams, graphs or rough work.  
Do not use staples, paper clips, highlighters, glue or correction fluid.  
**DO NOT WRITE IN ANY BARCODES.**

Answer **all** questions.  
Electronic calculators may be used.  
Qualitative Analysis Notes are printed on page 8.  
You should show the essential steps in any calculations and record experimental results in the spaces provided on the Question Paper.

At the end of the examination, fasten all your work securely together.  
The number of marks is given in brackets [ ] at the end of each question or part question.

For Examiner's Use	
1	
2	
<b>Total</b>	

This document consists of **7** printed pages and **1** blank page.

- 1 **P** is an aqueous solution which contains a mixture of sodium carbonate,  $\text{Na}_2\text{CO}_3$ , and sodium hydroxide,  $\text{NaOH}$ .

The concentration of sodium carbonate in **P** is  $0.0200 \text{ mol/dm}^3$ .

You are to determine by titration the volume of dilute sulfuric acid, **Q**, needed to neutralise a volume of **P**, and then calculate the concentration of sodium hydroxide present.

**Q** is  $0.100 \text{ mol/dm}^3$  sulfuric acid,  $\text{H}_2\text{SO}_4$ .

- (a) Put **Q** into the burette.

Pipette a  $25.0 \text{ cm}^3$  (or  $20.0 \text{ cm}^3$ ) portion of **P** into a flask and titrate with **Q**, using the indicator provided.

Record your results in the table, repeating the titration as many times as you consider necessary to achieve consistent results.

## Results

### *Burette readings*

titration number	1	2	
final reading / $\text{cm}^3$			
initial reading / $\text{cm}^3$			
volume of <b>Q</b> used / $\text{cm}^3$			
best titration results (✓)			

## Summary

Tick (✓) the best titration results.

Using these results, the average volume of **Q** required was .....  $\text{cm}^3$ .

Volume of **P** used was .....  $\text{cm}^3$ .

[12]

- (b) **Q** is 0.100 mol/dm<sup>3</sup> sulfuric acid, H<sub>2</sub>SO<sub>4</sub>.  
Calculate the number of moles of sulfuric acid present in your average volume of **Q**.

For  
Examiner's  
Use

moles of sulfuric acid present ..... [1]

- (c) The concentration of sodium carbonate, Na<sub>2</sub>CO<sub>3</sub>, in **P** is 0.0200 mol/dm<sup>3</sup>.  
Calculate the number of moles of sodium carbonate present in your volume of **P**.

moles of sodium carbonate present ..... [1]

- (d) Using your answer to (c), deduce the number of moles of sulfuric acid which react with the sodium carbonate present in your volume of **P**.




moles of sulfuric acid which react with the sodium carbonate ..... [1]

- (e) Using your answers to (b) and (d), calculate the number of moles of sulfuric acid which react with the sodium hydroxide in your volume of **P**.

moles of sulfuric acid which react with the sodium hydroxide ..... [1]

- (f) Using your answer to (e), calculate the concentration, in mol/dm<sup>3</sup>, of sodium hydroxide in **P**.



concentration of sodium hydroxide in **P** ..... mol/dm<sup>3</sup> [1]

[Total: 17]

2 You are provided with solution **R** and solid **S**.

Carry out the following tests and record your observations in the table.  
You should test and name any gas evolved.

For  
Examiner's  
Use

test no.	test	observations
1	To 2 cm depth of <b>R</b> in a test-tube, add a few drops of aqueous silver nitrate.  Keep this mixture for use in tests <b>2</b> and <b>3</b> .	
2	Transfer about half of the mixture from test <b>1</b> to a test-tube and add dilute nitric acid.	
3	To the remainder of the mixture from test <b>1</b> , add aqueous ammonia.	
4	To 2 cm depth of <b>R</b> in a test-tube, add a small amount of solid calcium carbonate.	
5	To 2 cm depth of <b>R</b> in a test-tube, add a piece of magnesium ribbon.	
6	To 1 cm depth of acidified potassium manganate(VII) solution in a test-tube, add a small amount of solid <b>S</b> .	

test no.	test	observations
7	Dissolve a small amount of solid <b>S</b> in 3 cm depth of distilled water in a test-tube.  Keep this solution for use in tests <b>8</b> and <b>9</b> .	
8	<p><b>(a)</b> To 1 cm depth of aqueous barium nitrate in a test-tube, add a few drops of the solution of <b>S</b> from test <b>7</b>.</p> <p><b>(b)</b> To the mixture from <b>(a)</b>, add <b>R</b>.</p>	
9	<p><b>(a)</b> To 1 cm depth of silver nitrate in a boiling tube, add the solution of <b>S</b> from test <b>7</b> until a change occurs.</p> <p><b>(b)</b> To the mixture from <b>(a)</b>, add the remainder of the solution of <b>S</b>.</p> <p><b>(c)</b> Warm the mixture from <b>(b)</b> until it just begins to boil and then leave the mixture to stand.</p>	
10	<p><b>(a)</b> To 1 cm depth of aqueous iron(III) chloride in a boiling tube, add a small amount of solid <b>S</b> and mix well.</p> <p><b>(b)</b> To the mixture from <b>(a)</b>, add an equal volume of <b>R</b> and then warm the mixture for about 10 seconds.</p> <p><b>(c)</b> To the mixture from <b>(b)</b>, add aqueous sodium hydroxide.</p>	

For  
Examiner's  
Use

[20]

Question 2 continues on page 6.

**Conclusions**

Identify a cation and an anion present in **R**.

A cation present in **R** is .....

An anion present in **R** is .....

In tests **6** and **10**, **S** is acting as .....

[3]

[Total: 23]

*For  
Examiner's  
Use*

**BLANK PAGE**

---

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

### QUALITATIVE ANALYSIS NOTES

#### Tests for anions

<i>anion</i>	<i>test</i>	<i>test result</i>
carbonate ( $\text{CO}_3^{2-}$ )	add dilute acid	effervescence, carbon dioxide produced
chloride ( $\text{Cl}^-$ ) [in solution]	acidify with dilute nitric acid, then add aqueous silver nitrate	white ppt.
iodide ( $\text{I}^-$ ) [in solution]	acidify with dilute nitric acid, then add aqueous silver nitrate	yellow ppt.
nitrate ( $\text{NO}_3^-$ ) [in solution]	add aqueous sodium hydroxide then add aluminium foil; warm carefully	ammonia produced
sulfate ( $\text{SO}_4^{2-}$ ) [in solution]	acidify with dilute nitric acid, then add aqueous barium nitrate	white ppt.

#### Tests for aqueous cations

<i>cation</i>	<i>effect of aqueous sodium hydroxide</i>	<i>effect of aqueous ammonia</i>
aluminium ( $\text{Al}^{3+}$ )	white ppt., soluble in excess giving a colourless solution	white ppt., insoluble in excess
ammonium ( $\text{NH}_4^+$ )	ammonia produced on warming	–
calcium ( $\text{Ca}^{2+}$ )	white ppt., insoluble in excess	no ppt., or very slight white ppt.
copper(II) ( $\text{Cu}^{2+}$ )	light blue ppt., insoluble in excess	light blue ppt., soluble in excess giving a dark blue solution
iron(II) ( $\text{Fe}^{2+}$ )	green ppt., insoluble in excess	green ppt., insoluble in excess
iron(III) ( $\text{Fe}^{3+}$ )	red-brown ppt., insoluble in excess	red-brown ppt., insoluble in excess
zinc ( $\text{Zn}^{2+}$ )	white ppt., soluble in excess giving a colourless solution	white ppt., soluble in excess giving a colourless solution

#### Tests for gases

<i>gas</i>	<i>test and test result</i>
ammonia ( $\text{NH}_3$ )	turns damp litmus paper blue
carbon dioxide ( $\text{CO}_2$ )	turns limewater milky
chlorine ( $\text{Cl}_2$ )	bleaches damp litmus paper
hydrogen ( $\text{H}_2$ )	'pops' with a lighted splint
oxygen ( $\text{O}_2$ )	relights a glowing splint