



Cambridge Assessment International Education
Cambridge Ordinary Level

CHEMISTRY

5070/32

Paper 3 Practical Test

October/November 2017

MARK SCHEME

Maximum Mark: 40

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the October/November 2017 series for most Cambridge IGCSE[®], Cambridge International A and AS Level components and some Cambridge O Level components.

Question	Answer	Marks
1(a)	<p>Titration</p> <p>Measurements (1) Both readings i.e. initial and final are present for each titration, readings are recorded to 1dp, no reading is in excess of 50.0 and no initial reading is given as 50.0</p> <p>Titres (1) All the titres are calculated correctly i.e. no subtraction errors</p> <p>Accuracy (6) For the two best titres give: 3 marks for a titre within 0.2cm³ of the Supervisor's value 2 marks for a titre within 0.3cm³ of the Supervisor's value 1 mark for a titre within 0.4cm³ of the Supervisor's value</p> <p>Concordance (3) Give 3 marks if all the ticked values are within 0.2cm³ Give 2 marks if all the ticked values are within 0.3cm³ Give 1 marks if all the ticked values are within 0.4cm³</p> <p>Average (1) Give 1 mark if the candidate calculates a correct average of selected titres</p>	12
1(b)	<p>Pipette volume = 25cm³ and assuming average volume of P used = 20.4cm³ Concentration of sulfuric acid in P in mol/dm³ = $(25.0 \times 0.586) / (20.4 \times 2)$ (1) = 0.359 (1)</p>	2
1(c)	<p>Moles of sulfuric acid in 10cm³ of concentrated acid = (b) / 2 = 0.359 / 2 = 0.180</p>	1
1(d)	<p>Concentration of concentrated sulfuric acid in mol/dm³ = (c) × 100 = 0.180 × 100 = 18.0</p>	1
1(e)	<p>Mass of sulfuric acid in 1dm³ of concentrated sulfuric acid in g = (d) × 98 = 18.0 × 98 = 1760</p>	1

Question	Answer	Marks
<p>General points R is aluminium chloride S is copper(II) chloride For gases: to gain credit for the name of the gas produced, the test must be at least partially correct. Solutions: colourless is not equivalent to clear and clear is not equivalent to colourless No credit is given for conclusions based upon incorrect observations.</p>		
<p>2 R (test 1)</p>	<p>(a) white ppt (1) (b) (ppt) dissolves / soluble (in excess) (1) colourless solution (1)</p>	<p>21</p>
<p>2 R (test 2)</p>	<p>(a) white ppt (1) (b) (ppt) remains / insoluble (in excess) (1) (c) no reaction (1)</p>	
<p>2 R (test 3)</p>	<p>(a) no reaction (1) (b) no reaction (1) (c) white ppt (1)</p>	
<p>2 S (test 1)</p>	<p>(a) blue ppt (1) (b) (ppt) remains / insoluble (in excess) (1)</p>	
<p>2 S (test 2)</p>	<p>(a) blue ppt (1) (b) (ppt) dissolves / soluble (in excess) (1) dark blue solution (1) (c) bubbles (1) gas relights a glowing splint (1) oxygen (1) (liquid) turns darker (1)</p>	
<p>2 S (test 3)</p>	<p>(a) no reaction (1) (b) no reaction (1) (c) white ppt (1)</p>	

Question	Answer	Marks
Conclusions	<p>R is aluminium chloride / $AlCl_3 / Al^{3+} Cl^-$ (1) Evidence: Test 1 white ppt which dissolves in excess, Test 2 white ppt which does not dissolve in excess and Test 3 correct in (a),(b) and (c)</p> <p>S is copper(II) chloride / $CuCl_2 / Cu^{2+} Cl^-$ (1) Evidence: Test 1 blue ppt which does not dissolve in excess, Test 2 blue ppt which dissolves in excess and Test 3 correct in (a), (b) and (c)</p>	2