

Cambridge O Level

CHEMISTRY 5070/11

Paper 1 Multiple Choice

October/November 2022

1 hour

You must answer on the multiple choice answer sheet.

You will need: Multiple choice answer sheet

Soft clean eraser

Soft pencil (type B or HB is recommended)


INSTRUCTIONS

There are forty questions on this paper. Answer all questions.

- For each question there are four possible answers **A**, **B**, **C** and **D**. Choose the **one** you consider correct and record your choice in soft pencil on the multiple choice answer sheet.
- Follow the instructions on the multiple choice answer sheet.
- Write in soft pencil.
- Write your name, centre number and candidate number on the multiple choice answer sheet in the spaces provided unless this has been done for you.
- Do not use correction fluid.
- Do not write on any bar codes.
- You may use a calculator.

INFORMATION


- The total mark for this paper is 40.
- Each correct answer will score one mark.
- Any rough working should be done on this question paper.
- The Periodic Table is printed in the question paper.



1 Which row shows the most appropriate apparatus for the measurement given?

	quantity	apparatus
Α	25.0 cm ³ of solution	measuring cylinder
В	32.7 cm ³ of solution	pipette
С	75 cm ³ of gas	gas syringe
D	80 cm ³ of solution	burette

2 A liquid, X, is distilled from a mixture using the apparatus shown.



During the distillation, the thermometer reads from 157 °C to 160 °C.

Which information about liquid X is correct?

- **A** The liquids in X may or may not be miscible (mix with each other).
- **B** X is a mixture that can be separated by distillation.
- **C** X must contain two liquids with boiling points 157 °C and 160 °C.
- **D** X must have been obtained by the fractional distillation of petroleum (crude oil).


3 An aqueous solution contains a salt, Y.

Addition of an aqueous solution X results in a precipitate being formed that redissolves when more X is added.


What could solution X and salt Y be?

	solution X	salt Y
Α	HC <i>l</i> (aq)	$AgNO_3$
В	H₂SO₄(aq)	Ba(NO ₃) ₂
С	NaOH(aq)	CuSO₄
D	NaOH(aq)	ZnSO₄

- Which gas diffuses the fastest at the same pressure?
 - nitrogen at 25°C
 - В nitrogen at 50 °C
 - C oxygen at 25°C
 - oxygen at 50°C
- 5 The diagram shows the outer shell electrons of the atoms of two elements, Q and R.







key

x = an electron

= a nucleus

The sulfate of Q is insoluble.

Element R is gaseous at room temperature and pressure.

Which row could be correct?

	proton number of Q	relative atomic mass of R
Α	12	35.5
В	12	80
С	56	80
D	56	35.5


- 6 Which statement about iodine atoms and iodide ions is correct?
 - **A** They are both isotopes of iodine.
 - **B** They undergo the same chemical reactions.
 - **C** They have the same number of protons.
 - **D** They have the same physical properties.
- 7 The element chlorine has two isotopes, ${}^{35}_{17}Cl$ and ${}^{37}_{17}Cl$.

In the Periodic Table, chlorine is shown as $^{35.5}_{17}$ Cl.

Which row shows the correct percentage of each isotope in a sample of naturally occurring chlorine?

	percentage of $^{35}_{17}\text{C}l$	percentage of $^{37}_{17}\text{C}l$
Α	25	75
В	40	60
С	50	50
D	75	25

- 8 What is the nucleon number of the isotope of uranium, $^{235}_{92}$ U?
 - **A** 92
- **B** 143
- **C** 235
- **D** 327
- **9** Silicon dioxide has a giant structure. Each silicon atom is joined to four oxygen atoms by covalent bonds. Part of the structure is shown.



Which property would silicon dioxide be expected to have?

- A a good conductor of electricity
- B a high melting point
- C reacts with hydrochloric acid
- **D** soluble in water

10 Elements X and Y react to form compound XY. Element Y has more electrons in its outer shell than element X. Compound XY conducts electricity in the molten state.

Which row correctly states the electron change that occurs during the reaction and the type of bonding in compound XY?

	electron change during formation of compound XY	type of bonding in compound XY		
Α	X donates electrons to Y	ionic		
В	X shares electrons with Y	covalent		
С	Y donates electrons to X	covalent		
D	Y shares electrons with X	ionic		

11 Which compound has the most single bonds in one mole

A CH₃CH₃

B CH₃CH₂OH

C CH₃CO₂H

D CH₃CHCH₂

12 The formula of ammonium metavanadate is NH₄VO₃. It consists of NH₄ ions and VO₃ ions.

What are the charges on these ions?

	NH₄ ion	VO ₃ ion
Α	1+	1–
В	2+	2–
С	3+	4–
D	4+	3–

13	Which mass of	oxygen gas	combines with exact	lv 16 a c	of sulfu	r to f	form sul	fur dioxide.	SO	,?
----	---------------	------------	---------------------	-----------	----------	--------	----------	--------------	----	----

A 4 q

B 8g

C 16g

D 32 g

14 The atomic number of ruthenium is 44. One of the oxides of ruthenium is a black solid, X. 5.79 g of X contains 1.39 g of oxygen.

What is the empirical formula of X?

 \mathbf{A} Ru₂O

B RuO

 \mathbf{C} RuO₂

D RuO₄

15 250 cm³ of 1.0 mol/dm³ hydrochloric acid reacts with an excess of solid sodium carbonate. The equation is shown.

$$2 HC\mathit{l}(aq) + Na_2CO_3(s) \rightarrow 2 NaC\mathit{l}(aq) + CO_2(g) + H_2O(I)$$

What is the volume of carbon dioxide produced when measured at room temperature and pressure?

- **A** $3.0\,\text{dm}^3$
- **B** $6.0 \, \text{dm}^3$ **C** $12 \, \text{dm}^3$
- **D** 24 dm³

16 When excess aqueous barium chloride is added to 25.0 cm³ of 1.00 mol/dm³ sodium sulfate, a white precipitate of barium sulfate is formed.

$$Na_2SO_4(aq) + BaCl_2(aq) \rightarrow BaSO_4(s) + 2NaCl(aq)$$

The precipitate is filtered off, washed, dried and weighed. 5.36 g barium sulfate is obtained.

What is the percentage yield of barium sulfate?

[M_r: Na₂SO₄, 142; BaCl₂, 208; BaSO₄, 233; NaCl, 58.5]

- **A** 2.3%
- **B** 27%
- **C** 92%
- **D** 97%

17 Aqueous copper(II) sulfate is electrolysed using inert electrodes.

Which statement is correct?

- Copper is collected at the anode.
- В Hydrogen is collected at the cathode.
- C Oxygen is collected at the anode.
- **D** Sulfur is collected at the cathode.
- **18** Concentrated aqueous sodium chloride is electrolysed using inert electrodes.

Which equation shows the reaction that occurs at the anode?

A
$$2Cl^- \rightarrow Cl_2 + 2e^-$$

B
$$2Cl^- + 2e^- \rightarrow Cl_2$$

$$\mathbf{C} \quad 2\mathbf{H}^{+} + 2\mathbf{e}^{-} \rightarrow \mathbf{H}_{2}$$

D
$$4OH^{-} \rightarrow O_{2} + 2H_{2}O + 4e^{-}$$

19 Which pair of equations correctly represents the reactions taking place at the anode and at the cathode during the electrolysis of molten silver bromide?

	anode	cathode
Α	$2Br^- \rightarrow Br_2 + 2e^-$	$Ag^{2+} + 2e^{-} \rightarrow Ag$
В	$Br^{2-} \rightarrow Br_2 + 2e^-$	$Ag^+ + e^- \rightarrow Ag$
С	$2Br^- \rightarrow Br_2 + 2e^-$	$Ag^+ + e^- \rightarrow Ag$
D	$Ag^+ + e^- \rightarrow Ag$	$2Br^- \rightarrow Br_2 + 2e^-$

- 20 Which two processes are both endothermic?
 - A combustion and cracking
 - **B** combustion and fermentation
 - C cracking and photosynthesis
 - **D** respiration and photosynthesis
- 21 A sample of sulfuric acid is added to 10 g of zinc granules. A reaction occurs and a gas is produced. The rate of the reaction is increased if a small amount of copper is added. The copper is unchanged after the reaction.

Which statement about this reaction is correct?

- **A** Copper acts as a biological catalyst in this reaction.
- **B** Copper lowers the activation energy of this reaction.
- **C** The rate of the reaction is greater as the particle size of the zinc is greater.
- **D** The rate of the reaction is greater if the pressure is increased.


22 The rate of reaction between magnesium and dilute hydrochloric acid is investigated. The equation is shown.

$$Mg(s) + 2HCl(aq) \rightarrow MgCl_2(aq) + H_2(g)$$

A known mass of magnesium is added to an excess of dilute hydrochloric acid.

The concentration of the hydrochloric acid and the volume of hydrogen produced is measured at regular time intervals.

Which pair of graphs correctly shows the experimental results?



23 In which reaction is the underlined substance reduced?

A CaCO₃ + 2HC
$$l \rightarrow$$
 CaC l_2 + H₂O + CO₂

B
$$Cu^{2+} + Zn \rightarrow Cu + Zn^{2+}$$

C Fe₂O₃ + 3CO
$$\rightarrow$$
 2Fe + 3CO₂

$$\textbf{D} \quad \mathsf{H}_2\mathsf{SO}_4 \, + \, \underline{\mathsf{Zn}} \, \to \, \mathsf{ZnSO}_4 \, + \, \mathsf{H}_2$$

- 24 Which change involves reduction?
 - A calcium carbonate to calcium oxide
 - **B** copper to brass
 - **C** ethene to poly(ethene)
 - **D** sand to silicon
- **25** Thiosulfate ions, $S_2O_3^{2-}$, react with iodine, I_2 , in aqueous solution.

$$2S_2O_3^{2-}(aq) + I_2(aq) \rightarrow 2I^{-}(aq) + S_4O_6^{2-}(aq)$$

In this reaction, the $S_2O_3^{2-}$ ions1..... electrons and are2.....

Which words correctly complete gaps 1 and 2?

	1	2		
Α	gain	oxidised		
В	gain	reduced		
С	lose	oxidised		
D	lose	reduced		

26 The equation shows a reaction in the Contact process.

$$2SO_2(g) + O_2(g) \rightleftharpoons 2SO_3(g)$$
 $\Delta H = -196 \text{ kJ/mol}$

Which change would move the position of equilibrium to the left?

- A adding more O₂
- **B** increasing the pressure
- **C** increasing the temperature
- D removing SO₃ from the reacting mixture

27 Some medicines contain the magnesium salt of a fatty acid. The fatty acids are members of the homologous series of carboxylic acids and contain one carboxylic acid group.

What is the carboxylic acid functional group and how many moles of a fatty acid react with one mole of magnesium?

	functional group	moles of fatty acid		
Α	–CO₂H	1		
В	–CO₂H	2		
С	–CH₂OH	1		
D	–CH₂OH	2		

28 Sodium sulfate, Na₂SO₄, and sodium hydrogensulfate, NaHSO₄, can both be prepared using aqueous sodium hydroxide and sulfuric acid.

$$2NaOH(aq) + H_2SO_4(aq) \rightarrow Na_2SO_4(aq) + 2H_2O(I)$$

 $NaOH(aq) + H_2SO_4(aq) \rightarrow NaHSO_4(aq) + H_2O(I)$

50 cm³ of 1 mol/dm³ sodium hydroxide is used each time.

Which row shows the correct volumes of 1 mol/dm³ sulfuric acid required to prepare a sample of sodium sulfate and a sample of sodium hydrogensulfate?

	volume of sulfuric acid required to make sodium sulfate /cm³	volume of sulfuric acid required to make sodium hydrogensulfate /cm³
Α	25	50
В	25	12.5
С	50	12.5
D	50	25

- **29** Which substance reacts with dilute sulfuric acid in the preparation of a pure sample of lead(II) sulfate?
 - **A** aqueous lead(II) nitrate
 - **B** lead foil
 - **C** powdered lead(II) carbonate
 - **D** powdered lead(II) oxide

- **30** Which statement about sulfuric acid is correct?
 - **A** In the manufacture of sulfuric acid, iron is used as the catalyst in the Contact process.
 - **B** Sulfuric acid is used in some batteries.
 - **C** Sulfuric acid is used as a fertiliser.
 - **D** Sulfuric acid is used as a food preservative.
- **31** Two statements are given.
 - statement 1 Going down Group I and Group VII, the melting point of the elements increases.
 - statement 2 Chlorine can displace iodine from aqueous potassium iodide but cannot displace bromine from aqueous potassium bromide.

Which statements are correct?

- A both statement 1 and statement 2
- B statement 1 only
- C statement 2 only
- **D** neither statement 1 nor statement 2
- **32** A reversible reaction involves a solid reacting with hydrogen.

Which of the metals, aluminium and iron, would catalyse the reaction and what is their effect on the position of equilibrium?

	act as a catalyst	position of equilibrium
Α	both aluminium and iron	moves to the right
В	both aluminium and iron	no change
С	iron only	moves to the right
D	iron only	no change

33 Iron is obtained in the blast furnace from the ore haematite.

Which statement is correct?

- **A** Calcium carbonate is used to remove acidic impurities.
- **B** Coke is reduced to carbon dioxide.
- **C** Haematite is oxidised by carbon monoxide.
- **D** Haematite undergoes thermal decomposition.

- **34** Three statements about the carbon cycle are given.
 - 1 The carbon cycle regulates the amount of carbon dioxide in the atmosphere.
 - 2 During photosynthesis, carbon dioxide is produced.
 - 3 Combustion of hydrocarbons requires oxygen.

Which statements are correct?

- **A** 1 and 2 only **B** 1 and 3 only **C** 2 and 3 only **D** 1, 2 and 3
- **35** Carbon is used in the purification of the water supply.

What is the reason for this?

- A to remove mud and other insoluble solids
- B to remove nitrates caused by the excessive use of fertiliser
- C to remove tastes and odours
- **D** to sterilise the water by removing harmful bacteria
- **36** Octane is an alkane with eight carbon atoms per molecule.

What is the molecular formula of octane, and how does its boiling point compare with that of butane?

	molecular formula of octane	boiling point of octane
Α	C ₈ H ₁₆	higher than butane
В	C ₈ H ₁₆	lower than butane
С	C ₈ H ₁₈	lower than butane
D	C ₈ H ₁₈	higher than butane

- 37 Which equation for the reaction between propane and chlorine is correct?
 - A $C_3H_6 + Cl_2 \rightarrow C_3H_6Cl_2$
 - **B** $C_3H_8 + Cl_2 \rightarrow C_3H_6Cl_2 + H_2$
 - C $C_3H_8 + Cl_2 \rightarrow CH_3Cl + C_2H_5Cl$
 - $D \quad C_3H_8 + Cl_2 \rightarrow C_3H_7Cl + HCl$

38 Propanoic acid reacts with calcium carbonate. The products of this reaction are calcium propanoate, carbon dioxide and water.

What is the equation for this reaction?

- A $2C_2H_5COOH + Ca_2CO_3 \rightarrow 2C_2H_5COOCa + CO_2 + H_2O$
- **B** $2C_2H_5COOH + CaCO_3 \rightarrow (C_2H_5COO)_2Ca + CO_2 + H_2O$
- \mathbf{C} 2C₃H₇COOH + Ca₂CO₃ \rightarrow 2C₃H₇COOCa + CO₂ + H₂O
- **D** $2C_3H_7COOH + CaCO_3 \rightarrow (C_3H_7COO)_2Ca + CO_2 + H_2O$
- **39** The monomer, CH₃CH=CHCH₃, can be used to make an addition polymer.


This addition polymer has a chain of carbon atoms joined to each other by C–C single bonds.

Each of these carbon atoms is also bonded to at least one other atom or group of atoms. These are called side groups.

Which statement describes the carbon atoms in the polymer chain made from CH₃CH=CHCH₃?

- A Every carbon atom in the chain has one –CH₃ and one hydrogen atom as side groups.
- **B** Every carbon atom in the chain is joined to a CH₃–CH– side group.
- **C** Every carbon atom in the chain is joined to either two –CH₃ or to two hydrogen atoms as side groups.
- **D** Every carbon atom in the chain is joined to hydrogen atoms only as side groups.

40 Which row correctly shows the structure of a polymer and the monomers from which it is made?



15

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.

The Periodic Table of Elements

	\equiv	2	He	elium 4	10	Je Pe	neon 20	18	٩r	argon 40	36	궃	rypton 84	54	Xe	t31	98	Zh Kh	nodon –			
			_	خ																		
	₹				6	ш	fluorine 19	17	Cl	chlorine 35.5	35	Ā	bromine 80	53	Н	iodine 127	85	¥	astatine			
	>				80	0	oxygen 16	16	ഗ	sulfur 32	34	Se	selenium 79	52	<u>a</u>	tellurium 128	84	Ъ	moloud –	116	_	livemorium -
	>	-			7	z	nitrogen 14	15	۵	phosphorus 31	33	As	arsenic 75	51	Sp	antimony 122	83	:E	bismuth 209			
	≥				9	ပ	carbon 12	14	Si	silicon 28	32	Ge	germanium 73	20	Sn	tin 119	82	Pb	lead 207	114	Εl	flerovium -
	≡	-			2	В	boron 11	13	Ρl	aluminium 27	31	Ga	gallium 70	49	In	indium 115	81	l_l	thallium 204			
		-						I			30	Zu	zinc 65	48	ဗ	cadmium 112	80	Hg	mercury 201	112	S	copernicium
											29	J.	copper 64	47	Ag	silver 108	62	Αu	gold 197	111	Rg	roentgenium
dn	,										28	Z	nickel 59	46	Pd	palladium 106	78	귙	platinum 195	110	Ds	darmstadtium -
Group											27	ပိ	cobalt 59	45	뫈	rhodium 103	77	'n	iridium 192	109	¥	meitnerium
		-	I	hydrogen 1							26	Ьe	iron 56	44	Ru	ruthenium 101	9/	SO	osmium 190	108	Hs	hassium
					ı						25	Mn	manganese 55	43	ည	technetium -	75	Re	rhenium 186	107	Bh	bohrium
						loc	SS				24	ပ်	chromium 52	42	Mo	molybdenum 96	74	≥	tungsten 184	106	Sg	seaborgium -
				Key	atomic number	atomic symbo	name relative atomic mass				23	>	vanadium 51	41	g	niobium 93	73	<u>a</u>	tantalum 181	105	Ор	dubnium
					,,0	ato	rela				22	i=	titanium 48	40	Zr	zirconium 91	72	士	hafnium 178	104	꿆	rutherfordium -
								•			21	Sc	scandium 45	39	>	yttrium 89	57-71	lanthanoids		89–103	actinoids	
	=				4	Be	beryllium 9	12	Mg	magnesium 24	20	Ca	calcium 40	38	Š	strontium 88	56	Ba	barium 137	88	Ra	radium
	_				က	:=	lithium 7	1	Na	sodium 23	19	¥	potassium 39	37	Вb	rubidium 85	55	Cs	caesium 133	87	Ļ	francium

rı Lu	lutetium 175	103	۲	lawrencium	I
70 Yb	ytterbium 173	102	%	nobelium	I
e9 Tm	thulium 169	101	Md	mendelevium	I
® 可	erbium 167	100	Fm	ferminm	I
67 Ho	holmium 165	66	Es	einsteinium	I
66 Dy	dysprosium 163	86	Ç	californium	I
65 Tb	terbium 159	97	ă	berkelium	I
64 G d	gadolinium 157	96	Cm	curium	I
63 Eu	europium 152	92	Am	americium	I
Sm	samarium 150	94	Pu	plutonium	I
Pm	promethium —	93	dN	neptunium	I
9 P	neodymium 144	92	⊃	uranium	238
P	praseodymium 141	91	Ра	protactinium	231
58 Ce	cerium 140	06	T	thorium	232
57 La	lanthanum 139	88	Ac	actinium	I

lanthanoids

actinoids

The volume of one mole of any gas is $24\,dm^3$ at room temperature and pressure (r.t.p.).