

# Cambridge O Level

CANDIDATE NAME					
CENTRE NUMBER			CANDIDATE NUMBER		

CHEMISTRY 5070/31

Paper 3 Practical Test

October/November 2022

1 hour 30 minutes

You must answer on the question paper.

You will need: The materials and apparatus listed in the confidential instructions

#### **INSTRUCTIONS**

- Answer all questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do not write on any bar codes.
- You may use a calculator.
- You should show all your working and use appropriate units.

#### **INFORMATION**

- The total mark for this paper is 40.
- The number of marks for each question or part question is shown in brackets [].
- Notes for use in qualitative analysis are provided in the question paper.

For Examiner's Use		
1		
2		
Total		

This document has 8 pages. Any blank pages are indicated.

**1 P** is a sample of dilute sulfuric acid.

**P** is prepared by adding 5.0 cm<sup>3</sup> of concentrated sulfuric acid to distilled water and making the total volume of the solution up to 250 cm<sup>3</sup> with distilled water.

**Q** is 0.360 mol/dm<sup>3</sup> sodium carbonate.

(a) Put P into the burette.

Pipette 25.0 cm<sup>3</sup> of **Q** into a flask and titrate with **P** using methyl orange indicator.

Record your results in the table, repeating the titration as many times as you consider necessary to achieve consistent results.

#### Results

### Burette readings

titration number	1	2	
final reading/cm <sup>3</sup>			
initial reading/cm <sup>3</sup>			
volume of <b>P</b> used/cm <sup>3</sup>			
best titration results (✓)			

### **Summary**

Tick  $(\checkmark)$  the best titration results.

[12]

(b)	<b>Q</b> is 0.360 mol/dm <sup>3</sup> sodium carbonate.		
	The equation for the reaction is shown.		
	$Na_2CO_3 + H_2SO_4 \rightarrow Na_2SO_4 + H_2O + CO_2$		
	Use your result from (a) to calculate the concentration, in mol/dm³, of sulfuric acid in P.		
	Give your answer to three significant figures.		
	mol/dm <sup>3</sup> [2]		
(c)	${f P}$ is prepared by adding 5.0 cm $^3$ of concentrated sulfuric acid to distilled water and making the total volume of the solution up to 250 cm $^3$ with distilled water.		
	Use your answer from <b>(b)</b> to calculate the number of moles of sulfuric acid in 5.0 cm <sup>3</sup> of concentrated sulfuric acid.		
	mol [1]		
(4)	Use your answer from <b>(c)</b> to calculate the concentration, in mol/dm <sup>3</sup> , of concentrated		
(u)	sulfuric acid.		
	mol/dm <sup>3</sup> [1]		
(e)	Use your answer from (d) to calculate the mass, in g, of sulfuric acid, $\rm H_2SO_4$ , in $\rm 1dm^3$ of concentrated sulfuric acid.		
	[M <sub>r</sub> : H <sub>2</sub> SO <sub>4</sub> , 98]		
	g [1]		
	[Total: 17]		

- 2 You are provided with two solutions, **R** and **S**.
  - (a) Do the following tests on **R** and record your observations in the table.

Test and name any gas evolved.

test no.		test	observations
1	(i)	To 1 cm depth of <b>R</b> in a test-tube, add aqueous sodium hydroxide until a change is seen.  To the mixture from (i), add excess aqueous sodium hydroxide.	
2	(ii)	To 1 cm depth of <b>R</b> in a test-tube, add aqueous ammonia until a change is seen.  To the mixture from (i), add excess aqueous ammonia.  Put 1 cm depth of aqueous hydrogen peroxide in a boiling tube.  Add the mixture from (ii) to this boiling tube.	
3		To 1 cm depth of <b>R</b> in a test-tube, add an equal volume of dilute nitric acid.  Pour half of the mixture from (i) into a test-tube and add an equal volume of aqueous barium nitrate.  To the other half of the mixture from (i), add an equal volume of aqueous silver nitrate.	

[12]

## (b) Conclusion

A solid is used to prepare solution  ${\bf R}.$ 

The name of the solid is ......

[1]

(c) Do the following tests on **S** and record your observations in the table.

Test and name any gas evolved.

test no.		test	observations
1	(i)	To 1 cm depth of <b>S</b> in a test-tube, add aqueous sodium hydroxide until a change is seen.	
	(ii)	To the mixture from (i), add excess aqueous sodium hydroxide.	
2	(i)	To 1 cm depth of <b>S</b> in a test-tube, add aqueous ammonia until a change is seen.	
	(ii)	To the mixture from (i), add excess aqueous ammonia.	
	(iii)	Put 1 cm depth of aqueous hydrogen peroxide in a boiling tube. Add the mixture from (ii) to this boiling tube.	
3	(i)	To 1 cm depth of <b>S</b> in a test-tube, add an equal volume of dilute nitric acid.	
	(ii)	Pour half of the mixture from (i) into a test-tube and add an equal volume of aqueous barium nitrate.	
	(iii)	To the other half of the mixture from (i), add an equal volume of aqueous silver nitrate.	
		alusian	[9

(d) Conclusion

A solid is used to prepare solution **S**.

The name of the solid is ......

[1]

[Total: 23]

### **BLANK PAGE**

7

### **BLANK PAGE**

#### **QUALITATIVE ANALYSIS NOTES**

#### **Tests for anions**

anion	test	test result
carbonate (CO <sub>3</sub> <sup>2-</sup> )	add dilute acid	effervescence, carbon dioxide produced
chloride (C <i>l</i> <sup>-</sup> ) [in solution]	acidify with dilute nitric acid, then add aqueous silver nitrate	white ppt.
iodide (I <sup>-</sup> ) [in solution]	acidify with dilute nitric acid, then add aqueous silver nitrate	yellow ppt.
nitrate (NO <sub>3</sub> <sup>-</sup> ) [in solution]	add aqueous sodium hydroxide then add aluminium foil; warm carefully	ammonia produced
sulfate (SO <sub>4</sub> <sup>2-</sup> ) [in solution]	acidify with dilute nitric acid, then add aqueous barium nitrate	white ppt., insoluble in excess dilute nitric acid

### Tests for aqueous cations

cation	effect of aqueous sodium hydroxide	effect of aqueous ammonia
aluminium (Al <sup>3+</sup> )	white ppt., soluble in excess giving a colourless solution	white ppt., insoluble in excess
ammonium (NH <sub>4</sub> <sup>+</sup> )	ammonia produced on warming	-
calcium (Ca <sup>2+</sup> )	white ppt., insoluble in excess	no ppt.
chromium(III) (Cr <sup>3+</sup> )	green ppt., soluble in excess giving a green solution	green ppt., insoluble in excess
copper(II) (Cu <sup>2+</sup> )	light blue ppt., insoluble in excess	light blue ppt., soluble in excess giving a dark blue solution
iron(II) (Fe <sup>2+</sup> )	green ppt., insoluble in excess	green ppt., insoluble in excess
iron(III) (Fe <sup>3+</sup> )	red-brown ppt., insoluble in excess	red-brown ppt., insoluble in excess
zinc (Zn <sup>2+</sup> )	white ppt., soluble in excess giving a colourless solution	white ppt., soluble in excess giving a colourless solution

### **Tests for gases**

gas	test and test result
ammonia (NH <sub>3</sub> )	turns damp red litmus paper blue
carbon dioxide (CO <sub>2</sub> )	turns limewater milky
chlorine (Cl <sub>2</sub> )	bleaches damp litmus paper
hydrogen (H <sub>2</sub> )	'pops' with a lighted splint
oxygen (O <sub>2</sub> )	relights a glowing splint

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.