www.dynamicpapers.com

Cambridge International AS & A Level

Cambridge International Examinations
Cambridge International Advanced Subsidiary and Advanced Level

CANDIDATE NAME

CENTRE NUMBER

CANDIDATE NUMBER

5858120452

CHEMISTRY **9701/33**

Paper 3 Advanced Practical Skills 1 **February/March 2017**

2 hours

Candidates answer on the Question Paper.

Additional Materials: As listed in the Confidential Instructions

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in.
Give details of the practical session and laboratory where appropriate, in the boxes provided.
Write in dark blue or black pen.
You may use an HB pencil for any diagrams or graphs.
Do not use staples, paper clips, glue or correction fluid.
DO **NOT** WRITE IN ANY BARCODES.

Answer **all** questions.
Electronic calculators may be used.
You may lose marks if you do not show your working or if you do not use appropriate units.
Use of a Data Booklet is unnecessary.

Qualitative Analysis Notes are printed on pages 10 and 11.
A copy of the Periodic Table is printed on page 12.

At the end of the examination, fasten all your work securely together.
The number of marks is given in brackets [ ] at the end of each question or part question.

Session
Laboratory

For Examiner's Use	
1	
2	
3	
Total	

This document consists of **12** printed pages.

© UCLES 2017

CAMBRIDGE International Examinations

www.dynamicpapers.com

1 The concentration of hydrogen peroxide may be given in mol dm^{-3} or as 'volume strength'. You will determine the concentration of hydrogen peroxide in mol dm^{-3} and in 'volume strength' by a gas collection method.
Hydrogen peroxide decomposes to form water and oxygen. The reaction is much faster in the presence of a catalyst such as manganese(IV) oxide.

$$2H_2O_2(aq) \rightarrow 2H_2O(l) + O_2(g)$$

'Volume strength' is defined as the volume of oxygen in cm^3 produced from the decomposition of 1.0 cm^3 of hydrogen peroxide at room temperature and pressure. For example, 1.0 cm^3 of '100 volume' hydrogen peroxide will produce 100 cm^3 of oxygen.

FA 1 is a solution of hydrogen peroxide, H_2O_2.
FA 2 is manganese(IV) oxide, MnO_2.

(a) Method

Read the whole method before starting any practical work.

The diagram below may help you in setting up your apparatus.



- Fill the tub with water to a depth of about 5 cm.
- Fill the 250 cm^3 measuring cylinder **completely** with water. Hold a piece of paper towel firmly over the top, invert the measuring cylinder and place it in the water in the tub.
- Remove the paper towel and clamp the inverted measuring cylinder so that the open end is in the water just above the base of the tub.
- Rinse the 50 cm^3 measuring cylinder with a little **FA 1** then use it to transfer 150 cm^3 of **FA 1** into the reaction flask labelled **X**.
- Check that the bung fits tightly in the neck of flask **X**, clamp flask **X** and place the end of the delivery tube into the inverted 250 cm^3 measuring cylinder.
- Remove the bung from the neck of the flask. Tip **FA 2** into the hydrogen peroxide and replace the bung **immediately**. Remove the flask from the clamp and swirl it to mix the contents. Swirl the flask occasionally until no more gas is given off. Replace the flask in the clamp.
- Measure and record the final volume of gas in the measuring cylinder in the space below.

Keep FA 1 for use in Question 2.

Result

[2]

© UCLES 2017

www.dynamicpapers.com

(b) Calculations

Show your working and appropriate significant figures in the final answer to **each** step of your calculations.

(i) Use the information on page 2 to calculate the 'volume strength' of **FA 1**.

'volume strength' of **FA 1** = ...

(ii) Calculate the number of moles of oxygen collected in the measuring cylinder.
[Assume 1 mole of gas occupies 24.0 dm^3 under these conditions.]

moles of O_2 = ... mol

(iii) Using your answer to **(ii)** calculate the number of moles of hydrogen peroxide in the volume of **FA 1** added to flask **X**.

moles of H_2O_2 = ... mol

(iv) Calculate the concentration of hydrogen peroxide, **FA 1**, in mol dm^{-3}.

concentration of H_2O_2, **FA 1** = ... mol dm^{-3}
[4]

© UCLES 2017

[Turn over

www.dynamicpapers.com

(c) (i) A source of error in this experiment is that some oxygen escapes before the bung can be inserted.

Suggest a change to the practical procedure given in **(a)** to reduce this source of error. You may draw a diagram as part of your answer.

.....

.....

(ii) The error in reading a 50 cm^3 measuring cylinder is $\pm 0.5\,cm^3$.

Calculate the maximum percentage error in the volume of hydrogen peroxide added to flask **X** in **(a)**.

maximum percentage error in volume of H_2O_2 = ... %

(iii) Explain why the presence of 20 cm^3 of air in the 250 cm^3 measuring cylinder before the start of the experiment would decrease the accuracy of the results obtained in **(a)**.

.....

.....

.....

[4]

(d) If you repeated the method described using half the mass of **FA 2**, what volume of gas would you expect to collect? Explain your answer.

.....

..... [1]

[Total: 11]

© UCLES 2017

www.dynamicpapers.com

2 You will carry out a second experiment to determine the concentration of hydrogen peroxide, **FA 1**, in mol dm^{-3}, by titration with acidified aqueous potassium manganate(VII). The equation for the reaction is given below.

$$2MnO_4^-(aq) + 5H_2O_2(aq) + 6H^+(aq) \rightarrow 2Mn^{2+}(aq) + 8H_2O(l) + 5O_2(g)$$

FA 1 is a solution of hydrogen peroxide, H_2O_2.
FA 3 is 0.0300 mol dm^{-3} potassium manganate(VII), $KMnO_4$.
FA 4 is dilute sulfuric acid.

(a) Method

- Fill the burette with **FA 3**.
- Pipette 25.0 cm^3 of **FA 1** into a conical flask.
- Use the 25 cm^3 measuring cylinder to add approximately 20 cm^3 of **FA 4** to the conical flask.
- Perform a rough titration and record your burette readings in the space below.

The rough titre is ..... cm^3.

- Carry out as many accurate titrations as you think necessary to obtain consistent results.
- Make certain any recorded results show the precision of your practical work.
- Record, in a suitable form below, all of your burette readings and the volume of **FA 3** added in each accurate titration.

I	
II	
III	
IV	
V	
VI	
VII	

[7]

(b) From your accurate titration results, obtain a suitable value for the volume of **FA 3** to be used in your calculations. Show clearly how you obtained this value.

25.0 cm^3 of **FA 1** required ..... cm^3 of **FA 3**. [1]

© UCLES 2017

[Turn over

www.dynamicpapers.com

(c) Calculations

Show your working and appropriate significant figures in the final answer to **each** step of your calculations.

(i) Calculate the number of moles of manganate(VII) ions present in the volume of **FA 3** calculated in **(b)**.

moles of MnO_4^- = ... mol

(ii) Calculate the number of moles of hydrogen peroxide present in 25.0 cm^3 of **FA 1**.

moles of H_2O_2 = ... mol

(iii) Using your answer to **(ii)** calculate the concentration, in mol dm^{-3}, of hydrogen peroxide in **FA 1**.

concentration of H_2O_2 in **FA 1** = ... mol dm^{-3}
[4]

[Total: 12]

© UCLES 2017

www.dynamicpapers.com

3 Qualitative Analysis

At each stage of any test you are to record details of the following.

- colour changes seen
- the formation of any precipitate
- the solubility of such precipitates in an excess of the reagent added

Where gases are released they should be identified by a test, **described in the appropriate place in your observations**.

You should indicate clearly at what stage in a test a change occurs.
No additional tests for ions present should be attempted.

If any solution is warmed, a boiling tube MUST be used.

Rinse and reuse test-tubes and boiling tubes where possible.

Where reagents are selected for use in a test, the name or correct formula of the element or compound must be given.

www.dynamicpapers.com

(a) **FA 5**, **FA 6** and **FA 7** are solutions, **some** of which contain ions that are listed on pages 10 and 11.

test	*observations*		
	FA 5	**FA 6**	**FA 7**
(i) To a 0.5 cm depth of solution in a boiling tube add aqueous sodium hydroxide, then			
warm gently.			
Allow to cool, add a piece of aluminium foil and warm again.			
(ii) To a 1 cm depth of solution in a test-tube add two or three drops of acidified aqueous potassium manganate(VII). (Do **not** use **FA 3**.)			
If no reaction occurs, pour the mixture into a boiling tube and warm gently.			
(iii) To a 1 cm depth of solution in a test-tube add a 2 cm depth of '10 volume' hydrogen peroxide and leave to stand. (Do **not** use **FA 1**.)			
(iv) To a 1 cm depth of solution in a test-tube add a 1 cm depth of dilute hydrochloric acid, then			
add a 1 cm depth of aqueous barium chloride or aqueous barium nitrate.			

[11]

© UCLES 2017

www.dynamicpapers.com

(b) (i) Identify as many ions present in **FA 5**, **FA 6** and **FA 7** as possible from your observations. If an ion cannot be identified from the tests, write 'unknown' in the space.

	cation(s)	anion(s)
FA 5		
FA 6		
FA 7		

(ii) Describe another test you could carry out to confirm the identity of a cation you have identified in **(i)**. Record the reagent(s) and expected observation(s) in the space below. **Do not carry out this test.**

(iii) Write an ionic equation for the reaction that would occur in **(ii)**. Include state symbols.

.....

[6]

[Total:17]

© UCLES 2017

www.dynamicpapers.com

Qualitative Analysis Notes

1 Reactions of aqueous cations

ion	*reaction with*	
	NaOH(aq)	NH_3(aq)
aluminium, Al^{3+}(aq)	white ppt. soluble in excess	white ppt. insoluble in excess
ammonium, NH_4^+(aq)	no ppt. ammonia produced on heating	–
barium, Ba^{2+}(aq)	faint white ppt. is nearly always observed unless reagents are pure	no ppt.
calcium, Ca^{2+}(aq)	white ppt. with high [Ca^{2+}(aq)]	no ppt.
chromium(III), Cr^{3+}(aq)	grey-green ppt. soluble in excess	grey-green ppt. insoluble in excess
copper(II), Cu^{2+}(aq)	pale blue ppt. insoluble in excess	blue ppt. soluble in excess giving dark blue solution
iron(II), Fe^{2+}(aq)	green ppt. turning brown on contact with air insoluble in excess	green ppt. turning brown on contact with air insoluble in excess
iron(III), Fe^{3+}(aq)	red-brown ppt. insoluble in excess	red-brown ppt. insoluble in excess
magnesium, Mg^{2+}(aq)	white ppt. insoluble in excess	white ppt. insoluble in excess
manganese(II), Mn^{2+}(aq)	off-white ppt. rapidly turning brown on contact with air insoluble in excess	off-white ppt. rapidly turning brown on contact with air insoluble in excess
zinc, Zn^{2+}(aq)	white ppt. soluble in excess	white ppt. soluble in excess

© UCLES 2017

www.dynamicpapers.com

2 Reactions of anions

ion	*reaction*
carbonate, CO_3^{2-}	CO_2 liberated by dilute acids
chloride, Cl^-(aq)	gives white ppt. with Ag^+(aq) (soluble in NH_3(aq))
bromide, Br^-(aq)	gives cream ppt. with Ag^+(aq) (partially soluble in NH_3(aq))
iodide, I^-(aq)	gives yellow ppt. with Ag^+(aq) (insoluble in NH_3(aq))
nitrate, NO_3^-(aq)	NH_3 liberated on heating with OH^-(aq) and Al foil
nitrite, NO_2^-(aq)	NH_3 liberated on heating with OH^-(aq) and Al foil; NO liberated by dilute acids (colourless NO → (pale) brown NO_2 in air)
sulfate, SO_4^{2-}(aq)	gives white ppt. with Ba^{2+}(aq) (insoluble in excess dilute strong acids)
sulfite, SO_3^{2-}(aq)	gives white ppt. with Ba^{2+}(aq) (soluble in excess dilute strong acids)

3 Tests for gases

gas	*test and test result*
ammonia, NH_3	turns damp red litmus paper blue
carbon dioxide, CO_2	gives a white ppt. with limewater (ppt. dissolves with excess CO_2)
chlorine, Cl_2	bleaches damp litmus paper
hydrogen, H_2	‘pops’ with a lighted splint
oxygen, O_2	relights a glowing splint

© UCLES 2017

www.dynamicpapers.com

The Periodic Table of Elements

Group																	
1	2											13	14	15	16	17	18
							1 H hydrogen 1.0										2 He helium 4.0
3 Li lithium 6.9	4 Be beryllium 9.0											5 B boron 10.8	6 C carbon 12.0	7 N nitrogen 14.0	8 O oxygen 16.0	9 F fluorine 19.0	10 Ne neon 20.2
11 Na sodium 23.0	12 Mg magnesium 24.3											13 Al aluminium 27.0	14 Si silicon 28.1	15 P phosphorus 31.0	16 S sulfur 32.1	17 Cl chlorine 35.5	18 Ar argon 39.9
19 K potassium 39.1	20 Ca calcium 40.1	21 Sc scandium 45.0	22 Ti titanium 47.9	23 V vanadium 50.9	24 Cr chromium 52.0	25 Mn manganese 54.9	26 Fe iron 55.8	27 Co cobalt 58.9	28 Ni nickel 58.7	29 Cu copper 63.5	30 Zn zinc 65.4	31 Ga gallium 69.7	32 Ge germanium 72.6	33 As arsenic 74.9	34 Se selenium 79.0	35 Br bromine 79.9	36 Kr krypton 83.8
37 Rb rubidium 85.5	38 Sr strontium 87.6	39 Y yttrium 88.9	40 Zr zirconium 91.2	41 Nb niobium 92.9	42 Mo molybdenum 95.9	43 Tc technetium –	44 Ru ruthenium 101.1	45 Rh rhodium 102.9	46 Pd palladium 106.4	47 Ag silver 107.9	48 Cd cadmium 112.4	49 In indium 114.8	50 Sn tin 118.7	51 Sb antimony 121.8	52 Te tellurium 127.6	53 I iodine 126.9	54 Xe xenon 131.3
55 Cs caesium 132.9	56 Ba barium 137.3	57–71 lanthanoids	72 Hf hafnium 178.5	73 Ta tantalum 180.9	74 W tungsten 183.8	75 Re rhenium 186.2	76 Os osmium 190.2	77 Ir iridium 192.2	78 Pt platinum 195.1	79 Au gold 197.0	80 Hg mercury 200.6	81 Tl thallium 204.4	82 Pb lead 207.2	83 Bi bismuth 209.0	84 Po polonium –	85 At astatine –	86 Rn radon –
87 Fr francium –	88 Ra radium –	89–103 actinoids	104 Rf rutherfordium –	105 Db dubnium –	106 Sg seaborgium –	107 Bh bohrium –	108 Hs hassium –	109 Mt meitnerium –	110 Ds darmstadtium –	111 Rg roentgenium –	112 Cn copernicium –		114 Fl flerovium –		116 Lv livermorium –		

Key

atomic number
atomic symbol
name
relative atomic mass

lanthanoids	57 La lanthanum 138.9	58 Ce cerium 140.1	59 Pr praseodymium 140.9	60 Nd neodymium 144.4	61 Pm promethium –	62 Sm samarium 150.4	63 Eu europium 152.0	64 Gd gadolinium 157.3	65 Tb terbium 158.9	66 Dy dysprosium 162.5	67 Ho holmium 164.9	68 Er erbium 167.3	69 Tm thulium 168.9	70 Yb ytterbium 173.1	71 Lu lutetium 175.0
actinoids	89 Ac actinium –	90 Th thorium 232.0	91 Pa protactinium 231.0	92 U uranium 238.0	93 Np neptunium –	94 Pu plutonium –	95 Am americium –	96 Cm curium –	97 Bk berkelium –	98 Cf californium –	99 Es einsteinium –	100 Fm fermium –	101 Md mendelevium –	102 No nobelium –	103 Lr lawrencium –

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cie.org.uk after the live examination series.

© UCLES 2017