UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS General Certificate of Education

Advanced Subsidiary Level and Advanced Level

CHEMISTRY

Paper 2 Structured Questions AS Core



9701/02

May/June 2006

1 hour 15 minutes

Candidates answer on the Question Paper. Additional Materials: Data Booklet

Candidate Name							
Centre Number				Candidate Number			

READ THESE INSTRUCTIONS FIRST

Write your name, Centre number and candidate number on all the work you hand in.

Write in dark blue or black pen in the spaces provided on the Question Paper.

You may use a pencil for any diagrams, graphs, or rough working.

Do not use staples, paper clips, highlighters, glue or correction fluid.

Answer all questions.

You may lose marks if you do not show your working or if you do not use appropriate units.

A Data Booklet is provided.

You may use a calculator.

The number of marks is given in brackets [] at the end of each question or part question.

At the end of the examination, fasten all your work securely together.

DO NOT WRITE IN THE BARCODE.

DO NOT WRITE IN THE GREY AREAS BETWEEN THE PAGES.

For Examiner's Use		
1		
2		
3		
4		
5		
Total		

Answer **all** the questions in the spaces provided.

For
Examiner's
1100

		alt is a pale green crystalline solid which is soluble in water. Mohr's salt is a 'double ch contains
		two cations, one of which is Fe ²⁺ ,
		one anion which is SO_4^{2-} ,
		and water of crystallisation.
(a)	Sol	e identity of the second cation was determined by the following test. Id Mohr's salt was heated with solid sodium hydroxide and a colourless gas was lived. The gas readily dissolved in water giving an alkaline solution.
	(i)	What is the gas?
	(ii)	What is the formula of the second cation identified by this test?
	(iii)	In this test, a grey/green solid residue was also formed.
		Suggest a name or formula for this colid
		Suggest a name or formula for this solid.
		Suggest a fiame or formula for this solid. [3]
(b)	hyd	
(b)	hyd salt	[3] identity of the anion present in Mohr's salt was confirmed by adding dilute rochloric acid followed by aqueous barium chloride to an aqueous solution of Mohr's
(b)	hyd salt	[3] identity of the anion present in Mohr's salt was confirmed by adding dilute rochloric acid followed by aqueous barium chloride to an aqueous solution of Mohr's . A white precipitate was formed.
(b)	hyd salt Sug 	[3] identity of the anion present in Mohr's salt was confirmed by adding dilute rochloric acid followed by aqueous barium chloride to an aqueous solution of Mohr's . A white precipitate was formed.
	hyd salt Sug 	[3] identity of the anion present in Mohr's salt was confirmed by adding dilute rochloric acid followed by aqueous barium chloride to an aqueous solution of Mohr's. A white precipitate was formed. [1] agest the identity of the white precipitate. [1] and a double salt such as Mohr's salt is made, the two individual salts are mixed.
	Sug Wh	[3] identity of the anion present in Mohr's salt was confirmed by adding dilute rochloric acid followed by aqueous barium chloride to an aqueous solution of Mohr's. A white precipitate was formed. Igest the identity of the white precipitate. [1] en a double salt such as Mohr's salt is made, the two individual salts are mixed ether in a 1:1 molar ratio, dissolved in water and the solution crystallised. Give the formula of each of the two salts that would be mixed to make the double

(ii)	Calculate the relative formula mass of each of the salts present in Mohr's salt. salt 1	For Examiner's Use
	relative formula mass of salt 1salt 2	
(iii)	relative formula mass of salt 2 The crystals of the double salt contain water of crystallisation.	
	The relative formula mass of Mohr's salt is 392. Use your answers to (ii) to calculate the number of moles of water of crystallisation present in one mole of Mohr's salt.	
	[6]	
	[Total: 10]	

2 The unsaturated hydrocarbon ethyne (acetylene), C_2H_2 , is widely used in 'oxy-acetylene torches' for cutting and welding metals. In the torch, ethyne is burned in oxygen to produce a flame with a temperature of 3400 K.

For Examiner's Use

(a) Ethyne is a linear molecule with a triple bond, C=C, between the two carbon atoms.

Draw a 'dot-and-cross' diagram of an ethyne molecule.

[1]

(b) When used for cutting or welding, ethyne is transported in cylinders which contain the gas under pressure. A typical cylinder has a volume of 76 dm³ and contains ethyne gas at 1515 kPa pressure at a temperature of 25 °C.

Use the general gas equation, pV = nRT, to calculate the amount, in moles, of ethyne in this cylinder.

[2]

(c) In some countries, ethyne is manufactured from calcium carbide, CaC₂, which is produced by heating quicklime and coke together at 2300 K.

$$CaO + 3C \rightarrow CaC_2 + CO$$

When water is added to the CaC_2 , calcium hydroxide, $Ca(OH)_2$, and ethyne, C_2H_2 , are produced.

(i) Construct a balanced equation for the formation of ethyne from calcium carbide.

(ii) Use this equation and your answer to part (b) to calculate the mass of CaC₂ which will react with an excess of water to produce enough ethyne to fill 100 cylinders of the gas.

[3]

(d)	Use	equation for the complete combustion of ethyne is given below. e appropriate bond energy data from the <i>Data Booklet</i> to calculate a value for the nalpy change of combustion of ethyne.	For Examiner's Use
		$C_2H_2(g) + \frac{5}{2}O_2(g) \rightarrow 2CO_2(g) + H_2O(g)$	
		[3]	
		[O]	
(e)	The	value for the standard enthalpy change of combustion of ethyne is -1300 kJ mol ⁻¹ .	
	(i)	Define the term standard enthalpy change of combustion.	
	(ii)	Explain why your answer to (d) does not have the same value as the standard enthalpy change of combustion.	

[Total: 12]

[3]

3

This	s que	stion is about the e	elements of Group VII,	the halogens.	
(a)	Con	nplete the following	table.		
		halogen	colour	physical state at]
			00.00.	room temperature	
		chlorine			
		bromine			_
		iodine			[2]
(b)		•	c acid is added to sepa and magnesium iodide	arate solid samples of ma	gnesium chloride,
	(i)	Describe, in each	case, one observatio	n you would be able to ma	ake.
		MgCl ₂			
		MgBr ₂			
		MgI ₂			
	(ii)	Give an equation chloride.	for the reaction of co	ncentrated sulphuric acid	d with magnesium
					[4]
(c)	Whe	en dilute nitric acid	and aqueous silver niti	rate are added to a solutio	on of a magnesium
` ,	hali	de, Mg X_2 , a pale cr	eam precipitate is for		-
		eous ammonia.	ible ili concentiated a	iqueous ammonia but no	or soluble in dilute
	(i)	What is the identit	y of the precipitate?		
	(ii)	Give an equation concentrated aque		s, for the reaction of the	e precipitate with
					[3]

)		of glass rod is plunged into separate gas jars, one containing hydrogen chloride and containing hydrogen iodide.
	(i)	For each gas, state what you would observe, if anything, and write an equation for any reaction that takes place.
		HC1
		HI
	(ii)	Explain your answer to (i) in terms of enthalpy changes.
((iii)	What is the role of the hot glass rod in any reaction that occurs?
		[6]
		[Total: 15]

4

in o	mpounds containing the allyl group, $\mathrm{CH_2=CHCH_2-}$, have pungent smells and are found philons and garlic. It also hol, $\mathrm{CH_2=CHCH_2OH}$, is a colourless liquid which is soluble in water.
(a)	Allyl alcohol behaves as an alkene and as a primary alcohol.
	Give the structural formula of the organic compound formed when allyl alcohol is
	(i) reacted with Br ₂ ,
	(ii) heated under reflux with an acidified solution of $\operatorname{Cr_2O_7^{2-}}$ ions.
	[2]
(b)	When allyl alcohol is reacted with MnO_2 at room temperature, propenal, $\mathrm{CH}_2\mathrm{=}\mathrm{CHCHO}$ is formed.
	What type of reaction is this?
	[1]
(c)	Allyl alcohol may be converted into propanal, $\mathrm{CH_3CH_2CHO}$, by using a ruthenium(IV) catalyst in water.
	CH ₂ =CHCH ₂ OH ruthenium(IV) catalyst CH ₃ CH ₂ CHO
	The reactant and the product are isomers. What form of isomerism do they display?
	[1]

(d)	Allyl alcohol can be converted into propanal in two steps $\mbox{without}$ the use of a ruthenium(IV) catalyst.
	$CH_2 \!\!=\!\! CHCH_2OH \ \xrightarrow{step I} \ CH_3CH_2CH_2OH \ \xrightarrow{step II} \ CH_3CH_2CHO$
	What reagents and conditions would be used for each step?
	step I
	reagent(s)
	condition(s)
	step II
	reagent(s)
	condition(s)[4]
(e)	By considering your answers to (b) and (d) , suggest what is unusual about the single-step reaction in (c) .
	[1]
(f)	Suggest the structural formula of the organic compound formed when allyl alcohol is
	(i) reacted with cold, dilute MnO ₄ ions,
	(ii) heated under reflux with acidified MnO ₄ ions.
	[3]
	[Total: 12]
	[Total. 12]

5

	•	nic compound is unchanged.
(a)	Wh	at is meant by the term substitution reaction?
		[1]
(b)		e example of a substitution reaction is the formation of an alcohol from a ogenoalkane.
	(i)	Write a balanced equation for the formation of ethanol from bromoethane.
	<i>(</i> 11)	
	(ii)	State the conditions for this reaction.
		[2]
(c)		few organic reactions, the product contains one more carbon atom than the starting terial.
	(i)	Write the equation for a reaction in which the organic compound bromoethane, which contains two carbon atoms, is converted into an organic compound which contains three carbon atoms.
	(ii)	State the conditions for this reaction.
		[2]

[Total: 11]

as the only organic compound.

(d) Ethanol may be converted into propanoic acid in a three-stage process which uses ethanol

	$C_2H_5OH \xrightarrow{\text{step I}} \mathbf{K} \xrightarrow{\text{step II}} \mathbf{L} \xrightarrow{\text{step III}} C_2H_5CO_2H$	
(i)	Give the structural formulae of the intermediate compounds K and L .	
	Κ	
	L	
(ii)	State the reagent(s) used and give the essential condition(s) for step I and step III.	for
	step I	
	reagent(s)	
	condition(s)	
	step III	
	reagent(s)	
	condition(s)	[6]

BLANK PAGE