

# Cambridge International AS & A Level

CANDIDATE NAME					
CENTRE NUMBER			CANDIDATE NUMBER		

CHEMISTRY 9701/23

Paper 2 AS Level Structured Questions

May/June 2021

1 hour 15 minutes

You must answer on the question paper.

You will need: Data booklet

#### **INSTRUCTIONS**

- Answer all questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do not use an erasable pen or correction fluid.
- Do not write on any bar codes.
- You may use a calculator.
- You should show all your working, use appropriate units and use an appropriate number of significant figures.

#### **INFORMATION**

- The total mark for this paper is 60.
- The number of marks for each question or part question is shown in brackets [ ].

This document has 12 pages. Any blank pages are indicated.

## Answer **all** the questions in the spaces provided.

1

The	e ele	ments in Group 17 are known as the halogens.	
(a)	Bet	ween the molecules of Group 17 elements van der Waals' forces exist.	
	(i)	State the trend in the relative strength of van der Waals' forces down Group 17.	
		[	1]
	(ii)	State the physical state of each of the halogens under room conditions.	
		chlorine	
		bromine	
		iodine	
			1]
(b)	A s I <sub>2</sub> (a	olution of aqueous bromide ions, Br <sup>-</sup> (aq), is added to separate samples of ${\rm C}l_2({\rm aq})$ are aq).	ıC
		scribe what is observed in each reaction. Explain your answer in terms of the relative ctivity of these elements as oxidising agents.	'E
	obs	servation on addition to C $l_2$ (aq)	
	obs	servation on addition to $ m I_2(aq)$	
	exp	lanation	
		[-	3
(c)	Ble	ach is made by reacting $\mathrm{C}l_2$ with cold NaOH(aq).	
	Wri	te an equation for the reaction of $\operatorname{Cl}_2$ with cold NaOH.	
			11

(d)	Wh	en C $l$ O $^-$ (aq) is added to water, it behaves as a Brønsted-Lowry base.	
	(i)	Define the term Brønsted-Lowry base.	
			[1]
	(ii)	Write an ionic equation for the reaction between $ClO^-$ and $H_2O$ .	
			[1 <sup>-</sup>
			•
(e)	The	concentration of NaC $l$ O in bleach <b>S</b> is $x g dm^{-3}$ .	
	Na	ClO reacts with H <sub>2</sub> O <sub>2</sub> (aq) as shown.	
		$H_2O_2(aq) + NaClO(aq) \rightarrow H_2O(I) + NaCl(aq) + O_2(g)$	
	A 5	.00 cm <sup>3</sup> sample of <b>S</b> completely reacts with $H_2O_2(aq)$ . The volume of $O_2(g)$ produced 0 cm <sup>3</sup> under room conditions.	is
	Ass	ume that only the NaC $l$ O in <b>S</b> reacts with $H_2O_2(aq)$ .	
	Cal	culate x. Show your working.	
		<i>x</i> = g dr	n⁻ <sup>(</sup> [3]
(f)		lium chlorate(I), NaC $l$ O, oxidises dilute hydrochloric acid to form three products. The ducts which contain chlorine have chlorine species with oxidation number $-1$ or 0.	he
	No	other species changes its oxidation number during the reaction.	
	Use	this information to complete the ionic equation.	
		C $lO$ <sup>−</sup> +HC $l$ → +	[2]
		[Total:	
		[ · · · · · · · · · · · · · · · · · · ·	٠.

Met	than	ol, CH <sub>3</sub> OH, is soluble in water because it forms hydrogen bonds with water molecules.
(a)		w a fully labelled diagram to show how a hydrogen bond forms between a water molecule a methanol molecule.
		[3]
(b)	Me	thanol has a melting point of –97.6 °C and a boiling point of 64.7 °C.
		ample of pure liquid methanol is added to a flask and then sealed. The sealed flask is left several days at constant temperature. The vapour pressure is then measured as 17 kPa.
	(i)	Describe what is meant by the term vapour pressure of methanol.
		[2]
	(ii)	Explain why some of the liquid becomes a vapour.
		[1]
(	(iii)	Suggest and explain why the vapour pressure of water at room temperature is lower than the vapour pressure of methanol at room temperature. Refer to the correct intermolecular forces in your answer.
		[2]

(c) Methanol is made by reacting carbon monoxide with hydrogen.

$$CO(g) + 2H_2(g) \rightleftharpoons CH_3OH(g)$$

Carbon monoxide and hydrogen react at  $1.0 \times 10^7 \, \text{Pa}$  and  $200 \, ^{\circ}\text{C}$ . Eventually the reaction mixture reaches dynamic equilibrium.

The table shows the amounts of each species present in the mixture.

	CO(g)	H <sub>2</sub> (g)	CH <sub>3</sub> OH(g)
initial amount/mol	1.0	2.0	0
equilibrium amount/mol	0.030	0.060	0.97

(i)	Explain what is meant by dynamic equilibrium.
	[2]
(ii)	Calculate the partial pressure of methanol vapour at equilibrium under these conditions. Show your working.
	Pa [2]
(iii)	Write an expression for the equilibrium constant, $K_{\rm p}$ , for this reaction. State the units in your answer.
	$\mathcal{K}_{p}$ =
	units = [2]
	[Total: 14]

[4]

- 3 Separate samples of **R**, **S**, **T** and **U** are added to cold water. The identity of each sample is unknown. However, each sample is known to be pure and can only be one of Ba(OH)<sub>2</sub>, NaCl, P<sub>4</sub>O<sub>10</sub> or SiCl<sub>4</sub>.
  - (a) (i) Use the observations in the table to identify each sample as one of Ba(OH)<sub>2</sub>, NaCl, P<sub>4</sub>O<sub>10</sub> and SiCl<sub>4</sub>. Write your answers in the table.

	state at room temperature	observations on addition of sample to water	identity of sample
R	solid	alkaline, colourless solution is produced, some white solid remains	
s	solid	white solid disappears, solution is neutral	
т	liquid	misty fumes produced, white solid is made in vigorous reaction	
U	solid	acidic, colourless solution produced in vigorous reaction	

	(ii)	Identify the formula of the white solid made when sample <b>T</b> reacts with water.	
			[1]
	(iii)	Name the solution formed when sample <b>U</b> reacts with water.	
			[1]
(b)	Ма	gnesium oxide and aluminium oxide have properties typical of ceramic materials.	
	(i)	Name one physical property typical of ceramic materials.	
			[1]
	(ii)	Give the formula of another Period 3 oxide which behaves as a ceramic material.	
			[1]


empirical formula = .....

(c) Tungsten oxide,  $W_x O_y$ , is used to give colour to ceramic materials.

A sample of $W_xO_y$ contains 79.29% tungsten by mass.
Calculate the empirical formula of W <sub>x</sub> O <sub>y</sub> .
Show your working.

[Total: 11]

**4** (a) 1,3-dichloropropan-2-ol can be made by reacting **M**.



(i) Give the systematic name of M.

F 4 7
111

(ii) Name the functional group present in **M** that changes during this reaction.


(iii) State a suitable reagent for this reaction.

**(b)** Separate samples of 1,3-dichloropropan-2-ol and 3-chloropropane-1,2-diol are heated with excess acidified  $\text{Cr}_2\text{O}_7^{2-}$  until there is no further reaction.

In each reaction, a different organic product, **Q** or **R**, is made.

1,3-dichloropropan-2-ol

3-chloropropane-1,2-diol



**Q** and **R** are tested separately with 2,4-dinitrophenylhydrazine solution, 2,4-DNPH, and sodium carbonate solution,  $Na_2CO_3(aq)$ .

Complete the table to give any relevant observations.

If no reaction occurs, write 'no visible change'.

reagent	observation with <b>Q</b>	observation with <b>R</b>
2,4-DNPH		
Na <sub>2</sub> CO <sub>3</sub> (aq)		

[4]

(c) Citric acid can be made from M in a four-step reaction.

Complete the table for each step of the reaction sequence to identify:

- the reagents and conditions required
- the type of reaction.

step	reagent and conditions	type of reaction
1		
2	dilute sulfuric acid	
3		
4	dilute sulfuric acid	

[5]

[Total: 12]

5 Compound **X** contains the same functional groups as citric acid.


The table describes some of the similarities and differences between citric acid and compound X.

	citric acid	X
chiral centre	no	yes
reaction with Na	fizzing	fizzing
reaction with H <sup>+</sup> /Cr <sub>2</sub> O <sub>7</sub> <sup>2-</sup>	remains orange	orange to green

(a) Complete the equation to show the reaction of excess sodium with citric acid. Show the skeletal structure of the product.

[3]

(b) (i) Use the information in the table to deduce the skeletal formula of **X**, C<sub>3</sub>H<sub>6</sub>O<sub>3</sub>. Draw the skeletal formula of **X** in the box. Label the chiral centre of compound **X** with an asterisk (\*).



[2]

(ii) Explain why compound **X** reacts with acidified Cr<sub>2</sub>O<sub>7</sub><sup>2-</sup> but citric acid does not.

(c)	Compound X is one of a pair of stereoisomers.
	Stereoisomerism occurs when a molecule has at least one of two key features.
	State the two key features that give rise to stereoisomerism.
	1
	2
	[2]
(d)	A structural isomer of compound <b>X</b> does not fizz when added to sodium.
	Explain what is meant by structural isomer.
	[1]
	[Total: 10]

12

### **BLANK PAGE**

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.