

CAMBRIDGE
INTERNATIONAL EXAMINATIONS

NOVEMBER 2002

GCE Advanced Level

MARK SCHEME

MAXIMUM MARK : 40

SYLLABUS/COMPONENT :9701 /6


**CHEMISTRY
(OPTIONS (A2))**



UNIVERSITY of CAMBRIDGE
Local Examinations Syndicate

Biochemistry

1. (a) (i)




(2 x 1)

(ii) Chiral / anomeric / optically active centre is created since rotation is possible at C₁

(1)
[3]

(b) Hydrogen bonding

(1)



(1)
[2]


(c) (i) glucose + ATP => glucose-6-phosphate + ADP

(1)

an enzyme / hexokinase / glucokinase is needed

(1)

(ii)




(1)

(iii) Glucose-6-phosphate is a competitive/reversible inhibitor

(1)

It fits into the active site on the enzyme/similar shape to glucose

(1)
[5]



Axes labelled (1)
 Graph (1)
 V_{max} and $V_{max} / 2$ indicated (1)

(ii) Check lines on sketch above for $V_{max}/2$ to give K_m as [S] (1)

(iii) The value of K_m shows the efficiency / effectiveness of the enzyme / affinity of the enzyme for the substrate / strength of the enzyme-substrate bond (1)

Small values for K_m indicate very efficient systems (1)
 [6]

(b) (i) Correct line on sketch (1)
 Competes for active sites on the enzyme (1)

(ii) Correct line on sketch (1)
 Increase the efficiency of the enzyme (1)
 [4]

Page 3	Mark Scheme	www.dynamicpapers.com	Syllabus	Paper
	A Level Examinations – November 2002	9701	6	

Environmental Chemistry

3. (a) Increased use of fertilisers (1)
- Leaching / runoff of soluble compounds such as nitrates (1)
- This increases the growth of algae (1)
- When these die and decay they use up dissolved oxygen / eutrophication (1)
- [4]
- (b) Water from the Baltic is less dense due to lower salinity **and** higher temperatures (**both** required) (1)
- (c) Nutrient levels are greatest in the North Sea water in which the algae grow (1)
- The 'jump' layer is not as mobile as the surface waters / little or no mixing (1)
- [2]
- (d) The algal decomposition mainly affects the deeper waters reducing the oxygen content (1)
- Oxygen loss is less significant at the surface (1)
- The smaller the cod populations, the fewer herrings are eaten (1)
- Cod are found at greater depths where the oxygen loss is greatest (1)
- [max 2]
- (e) This shows severely reducing conditions / a large oxygen loss. (1)

Page 4	Mark Scheme	Syllabus	Paper
	A Level Examinations – November 2002	9701	6

4. (a) Lack of flammability / inertness to combustion

Suitable volatility / easily liquefied

Lack of reactivity towards other chemicals present

Non-toxic

[max 2]



$\text{Cl}\cdot$ is recycled, and can thus destroy many ozone molecules (1)

[4]

- (c) (i) It breaks down more easily (1)

- (ii) CFC-11 must have a shorter residence time than CFC-12 (1)

CFC-12 must have a very long residence time (> 100 years) (1)

[max 2]

- (d) HCFCs are more readily destroyed in the troposphere (1)

The C—H bond is more readily attacked, and this promotes the breakdown of the molecule (1)

Polarisation of the C-H bond (1)

[max 2]

Page 5	Mark Scheme	www.dynamicpapers.com	
	A Level Examinations – November 2002	9701	6

Phase Equilibria

5. (a) As the molecules gain energy (1)
the forces between them become much weaker (1)
The magnitude of the change is proportional to ΔH_{vap} (1)
[max 2]
- (b) (i) H_2O has a high b.p. due to hydrogen bonding (1)
Diagram of water showing 2 H-bonds per molecule (1)
 H_2S to H_2Te have similar intermolecular dipole-dipole forces / van der Waals' (1)
- (ii) $\text{H}_2\text{O} : \frac{40.7}{373} = 0.109$ $\text{H}_2\text{S} : \frac{18.7}{213} = 0.088$
 $\text{H}_2\text{Se} : \frac{19.3}{243} = 0.079$ $\text{H}_2\text{Te} : \frac{23.2}{268} = 0.087$
Four values (1)
For similar bonding, b.p. and ΔH_{vap} are proportional (1)
Water has a higher ratio due to different / stronger hydrogen bonding (1)
[max 5]
- (c) (i) $P = P_A \times X_A$
The vapour pressure exerted by a gas is proportional to its mole fraction (1)
- (ii) Law holds only for similar intermolecular forces / H_2S and H_2Se both have van der Waals' forces (1)
 H_2O and H_2S have different forces (1)
[3]

Page 6	Mark Scheme	Syllabus	Paper
	A Level Examinations – November 2002	9701	6

6. (a) (i) Partition coefficient = $\frac{[X]_{\text{solvent 1}}}{[X]_{\text{solvent 2}}}$ (1)

(ii) $K = \frac{1.0 \times 10^{-2}}{4.0 \times 10^{-3}} = 2.5$ (1)

(iii) Let x mol of iodine be dissolved by the solvent

Then $(4.0 \times 10^{-4} - x)$ mol I_2 remain in 100 cm^3 water (1)

And x mol I_2 are present in 50 cm^3 of solvent (1)

$$2.5 = \frac{[\text{Concn in solvent}]}{[\text{Concn in water}]} = \frac{20x}{10(4.0 \times 10^{-4} - x)}$$
 (1)

This gives $20x = 25(4.0 \times 10^{-4} - x)$

$$45x = 10^{-2}$$

$$x = 2.2 \times 10^{-4} \text{ mol}$$
 (1)

Hence the concn of I_2 in the solvent is $20 \times 2.2 \times 10^{-4} \text{ mol dm}^{-3}$

or $4.4 \times 10^{-3} \text{ mol dm}^{-3}$ (1)
[max 6]

(b) (i) The solubility of a gas in a liquid is proportional to the (partial) pressure of the gas (1)

(ii) Solubility of $N_2 = 0.79 \times 23.6 = 18.6 \text{ cm}^3 \text{ dm}^{-3}$ (1)


Solubility of $O_2 = 0.20 \times 48.9 = 9.8 \text{ cm}^3 \text{ dm}^{-3}$ (1)

(iii) % N_2 is $\frac{18.6}{18.6 + 9.8} = \frac{18.6}{28.4} = 65.5\%$

And hence % $O_2 = 34.5\%$ (1)
[4]

Spectroscopy

7. (a)



-1 for every line over 3

(3 x 1)
[3]



(2 x 1)
[2]

-1 for every incorrect over 2


(c) (i) Diphenylmethanone will absorb at lower energy (longer wavelength) (1)

(ii) Energy levels are closer together (1)

hence less energy is required for transitions (1)
(allow longer chromophore / greater delocalisation / conjugation)

[3]

(d)




[2]

8. (a) (i) A suspension of an organic solid (1)
in a hydrocarbon oil / Nujol (1)
- (ii) Ethanol shows a strong IR absorption due to -OH (1)
It absorbs water which would attack the NaCl plates (1)
[max 3]

(b) Q $\text{-C}\equiv\text{N}$ R $\text{C}=\text{O}$ S -C-O-

(3 correct 2 marks
2 correct 1 mark)



Or ester isomers of the above, NOT -OH / -NH_2 containing isomers (1)
[3]

- (c) (i) Peak is at $M - 15$, hence CH_3 has been lost (1)
- (ii) T to U is a loss of 30, suggests loss of CH_2O or $\text{-CH}_2\text{NH}_2$ (1)
- (iii) Ratio of $M : M+1$ gives $n = \frac{0.11 \times 100}{2.5 \times 1.1}$ or 4 carbons (1)
- If K is saturated, it will contain 8 hydrogens ($\text{C}_4\text{H}_8\text{O}_x$) (1)
- This leaves a mass of 32 for the oxygen
- Hence K is $\text{C}_4\text{H}_8\text{O}_2$ (1)
[max 4]

Transition Elements

9. (a) Labelled diagram is acceptable

Impure copper anode, pure copper cathode (1)

Copper is transferred to the cathode (or equations) (1)

$\text{CuSO}_4(\text{aq})$ is the electrolyte (1)

Silver settles as the metal in the anode sludge (1)

Because E° is more +ve than Cu^{2+} (1)

Ni / Zn goes into solution as M^{2+} (1)

Because their E° is more negative than Cu^{2+} (1)
[max6]

(b) (i) Brass, with zinc; bronze, with tin etc (1)

(ii) moles of $\text{S}_2\text{O}_3^{2-} = 0.1 \times 20/1000 = 2 \times 10^{-3} \text{ mol}$ (1)

moles of $\text{I}_2 = 1 \times 10^{-3}$
moles of $\text{Cu}^{2+} = 2 \times 10^{-3}$ (1)

Mass of copper = $63.5 \times 2 \times 10^{-3} \text{ g}$
= 0.127 g

Hence % copper = 50.8% (1)
[4]

10. (a) $[\text{Ar}]3d^4$ [1]

(b) Mn(II) colourless OR pale pink

Mn(III) red

Mn(VI) green

Mn(VII) purple

4 correct, 3 marks
3 correct, 2 marks etc

[3]

(c) (i) From Data Book : $4\text{MnO}_4^{2-} - 4e^- \Rightarrow 4\text{MnO}_4^-$ $E^\circ = 0.56\text{V}$ (1)

$5\text{MnO}_4^{2-} + 8\text{H}^+ \Rightarrow \text{Mn}^{2+} + 4\text{MnO}_4^- + 4\text{H}_2\text{O}$ (1)

$E^\circ_{\text{cell}} = +1.74 - 0.56 = +1.18\text{V}$ (1)

(ii) Oxidation no = +5 (1)

$8\text{H}^+ + 3\text{MnO}_4^{3-} \Rightarrow 2\text{MnO}_2 + \text{MnO}_4^- + 4\text{H}_2\text{O}$
(1 for correct formulae, 1 for balancing)(2)

[6]