|             | Cambridge<br>International<br>AS & A Level  | Cambridge International Examinations<br>Cambridge International Advanced Subsidiary and Advar                           | nced Level  |          |  |  |  |  |  |  |  |  |  |
|-------------|---|---|-------------|----------|--|--|--|--|--|--|--|--|--|
|             | CANDIDATE<br>NAME   |   |             |          |  |  |  |  |  |  |  |  |  |
|             | CENTRE<br>NUMBER  | CANDIE  | NDIDATE BER |          |  |  |  |  |  |  |  |  |  |
| * 0 1 6 3 7 | CHEMISTRY<br>Paper 3 Advan  | 9701/31<br>October/November 2016  |             |          |  |  |  |  |  |  |  |  |  |
| 07882       | 2 hours         Candidates answer on the Question Paper.         Additional Materials:       As listed in the Confidential Instructions   |   |             |          |  |  |  |  |  |  |  |  |  |
| * 🚃         | READ THESE INSTRUCTIONS FIRST   |   |             |          |  |  |  |  |  |  |  |  |  |
|             | <ul> <li>Write your Centre number, candidate number and name on all the work you hand in.</li> <li>Give details of the practical session and laboratory where appropriate, in the boxes provided.</li> <li>Write in dark blue or black pen.</li> <li>You may use an HB pencil for any diagrams or graphs.</li> <li>Do not use staples, paper clips, glue or correction fluid.</li> <li>DO NOT WRITE IN ANY BARCODES.</li> <li>Answer all questions.</li> <li>Electronic calculators may be used.</li> <li>You may lose marks if you do not show your working or if you do not use appropriate units.</li> </ul> |   |             |          |  |  |  |  |  |  |  |  |  |
|             |   | ooklet is unnecessary.  | Sessio      | on       |  |  |  |  |  |  |  |  |  |
|             |   | ysis Notes are printed on pages 10 and 11.<br>eriodic Table is printed on page 12.                                      |             |          |  |  |  |  |  |  |  |  |  |
|             |   | e examination, fasten all your work securely together.<br>marks is given in brackets [ ] at the end of each question or | Laboratory  |          |  |  |  |  |  |  |  |  |  |
|             |   |   | For Examine | er's Use |  |  |  |  |  |  |  |  |  |
|             |   |   | 1           |          |  |  |  |  |  |  |  |  |  |
|             |   |   | 2           |          |  |  |  |  |  |  |  |  |  |
|             |   |   | 3           |          |  |  |  |  |  |  |  |  |  |
|             |   |   | Total       |          |  |  |  |  |  |  |  |  |  |
|             |   | This document consists of <b>12</b> printed pages.  |             |          |  |  |  |  |  |  |  |  |  |

1 In **Questions 1** and **2** you will determine the percentage purity of industrial grade calcium carbonate, CaCO<sub>3</sub>, by two different methods.


In the first method you will collect and measure the volume of gas given off in the reaction between a known mass of industrial grade calcium carbonate, in the form of small marble chips, and a known amount of dilute hydrochloric acid. The acid will be in excess. The impurities in the calcium carbonate will not react with the acid.

$$CaCO_3(s) + 2HCl(aq) \rightarrow CaCl_2(aq) + H_2O(I) + CO_2(g)$$

**FA 1** is industrial grade calcium carbonate,  $CaCO_3$ , in the form of small marble chips. **FA 2** is 2.00 mol dm<sup>-3</sup> hydrochloric acid, HC*l*.

#### (a) Method

#### **Read through the whole method before starting any practical work.** The diagram below may help you in setting up your apparatus.



- Fill the tub with water to a depth of about 5 cm.
- Fill the 250 cm<sup>3</sup> measuring cylinder **completely** with water. Hold a piece of paper towel firmly over the top, invert the measuring cylinder and place it in the water in the tub.
- Remove the paper towel and clamp the inverted measuring cylinder so the open end is in the water just above the base of the tub.
- Pipette 25.0 cm<sup>3</sup> of **FA 2** into the reaction flask labelled **X**.
- Check that the bung fits tightly in the neck of flask **X**, clamp flask **X** and place the end of the delivery tube into the inverted 250 cm<sup>3</sup> measuring cylinder.
- Weigh the container with **FA 1** and record the mass in the space on page 3.
- Remove the bung from the neck of the flask. Tip **FA 1** into the acid and replace the bung **immediately**. Remove the flask from the clamp and swirl it to mix the contents. Swirl the flask occasionally until no more gas is evolved. Replace the flask in the clamp.
- Reweigh the container and any residue of **FA 1** and record the mass in the space on page 3.
- Calculate and record in the space on page 3 the mass of FA 1 used.
- When no more gas is given off, measure and record the final volume of gas in the measuring cylinder in the space on page 3.

### Keep the contents of flask X for use in Question 2.

#### Results

#### (b) Calculations

Show your working and appropriate significant figures in the final answer to **each** step of your calculations.

3

(i) Calculate the number of moles of carbon dioxide gas collected in the measuring cylinder. (Assume that 1 mole of gas occupies 24.0 dm<sup>3</sup> under these conditions.)

moles of  $CO_2$  = ..... mol

(ii) Use your answer to (i) and the Periodic Table on page 12 to calculate the mass of pure calcium carbonate in the sample of industrial grade calcium carbonate, **FA 1**.

mass of  $CaCO_3$  = ..... g

(iii) Use your answer to (ii) and the mass of marble chips used in (a) to calculate a value for the percentage purity of the sample of industrial grade calcium carbonate, **FA 1**.

percentage purity of **FA 1** = .....% [4]

(c) Not all the carbon dioxide given off in the reaction is collected in the measuring cylinder.

Suggest a change to the method which would lead to an increase in the volume of carbon dioxide collected.

.....[1]

[Total: 7]

2 You will determine the amount of hydrochloric acid remaining in flask **X** after the reaction with the marble chips in **Question 1**. You will do this by titration with sodium hydroxide of known concentration.

 $NaOH(aq) + HCl(aq) \rightarrow NaCl(aq) + H_2O(I)$ 

The impurities in the calcium carbonate will not react with the alkali.

**FA 3** is 0.140 mol dm<sup>-3</sup> sodium hydroxide, NaOH. bromophenol blue indicator

#### (a) Method

- Transfer **all** the contents of flask **X** into the 250 cm<sup>3</sup> volumetric flask.
- Rinse flask **X** with distilled water and add the washings to the volumetric flask. Add distilled water up to the mark.
- Stopper the volumetric flask and mix the contents thoroughly. Label this solution FA 4.
- Rinse the pipette then use it to transfer 25.0 cm<sup>3</sup> of **FA 4** into a conical flask.
- Add about 10 drops of bromophenol blue indicator.
- Fill the burette with **FA 3**.
- Perform a rough titration and record your burette readings in the space below.

The rough titre is ...... cm<sup>3</sup>.

- Carry out as many accurate titrations as you think necessary to obtain consistent results.
- Record, in a suitable form below, all of your burette readings and the volume of **FA 3** added in each accurate titration.
- Make certain any recorded results show the precision of your practical work.



(b) From your accurate titration results, obtain a suitable value for the volume of **FA 3** to be used in your calculations. Show clearly how you obtained this value.

25.0 cm<sup>3</sup> of **FA 4** required ..... cm<sup>3</sup> of **FA 3**. [1]

### (c) Calculations

Show your working and appropriate significant figures in the final answer to **each** step of your calculations.

(i) Calculate the number of moles of sodium hydroxide, NaOH, present in the volume of **FA 3** you calculated in (b).

moles of NaOH = ..... mol

(ii) Use your answer to (i) and the equation on page 4 to determine the number of moles of hydrochloric acid, HC*l*, present in the 25.0 cm<sup>3</sup> of **FA 4** pipetted in (a).

moles of HCl = ..... mol

(iii) Use your answer to (ii) to calculate the number of moles of hydrochloric acid, HC*l*, remaining in flask **X** after the reaction in **1(a)**.

moles of HC*l* remaining = ..... mol

(iv) Use the relevant information on page 2 to calculate the number of moles of hydrochloric acid, HC*l*, pipetted into flask **X** in **1(a)**.

moles of HCl pipetted into flask **X** = ..... mol

(v) Use your answers to (iii) and (iv) to calculate the number of moles of hydrochloric acid, HC*l*, which reacted with the marble chips in flask **X**.

moles of HCl which reacted in flask **X** = ..... mol

(vi) Use your answer to (v), the equation in Question 1 and the Periodic Table on page 12 to calculate the mass of pure calcium carbonate, CaCO<sub>3</sub>, in the sample of industrial grade calcium carbonate, FA 1.

(vii) Use your answer to (vi) and the mass of marble chips recorded in 1(a) to calculate the percentage purity of FA 1.

percentage purity of **FA 1** = .....% [5]

(d) You have carried out two different methods to find the percentage purity of industrial grade calcium carbonate.

A source of error in **Question 1** is that some carbon dioxide escapes before the bung can be inserted.

How would this affect the percentage purity of **FA 1** calculated in the two questions? Explain your answers.

**Question 1** 

Question 2

.....

[3]

[Total: 16]

#### 3 Qualitative Analysis

At each stage of any test you are to record details of the following.

- colour changes seen
- the formation of any precipitate
- the solubility of such precipitates in an excess of the reagent added

Where gases are released they should be identified by a test, **described in the appropriate place in your observations**.

You should indicate clearly at what stage in a test a change occurs. No additional tests for ions present should be attempted.

#### If any solution is warmed, a boiling tube MUST be used.

Rinse and reuse test-tubes and boiling tubes where possible.

Where reagents are selected for use in a test, the name or correct formula of the element or compound must be given.

(a) **FA 5** and **FA 6** are solids each containing one cation and one anion. Carry out the following tests and record your observations in the table below.

| test  | observations |      |  |  |  |  |  |  |
|---|--------------|------|--|--|--|--|--|--|
| 1851  | FA 5         | FA 6 |  |  |  |  |  |  |
| <ul> <li>(i) Place a spatula<br/>measure of solid in a<br/>hard-glass test-tube<br/>and heat gently at<br/>first, then</li> </ul>   |              |      |  |  |  |  |  |  |
| heat strongly until no<br>further change takes<br>place.  |              |      |  |  |  |  |  |  |
| Leave the tube to<br>cool completely then<br>add a 2 cm depth of<br>dilute sulfuric acid to<br>the solid residue.<br>Shake the contents<br>of the tube then<br>leave it to stand. |              |      |  |  |  |  |  |  |

|       | 10 c1  | obser                             | vations           |  |  |  |  |  |
|-------|--|-----------------------------------|-------------------|--|--|--|--|--|
|       | test   | FA 5                              | FA 6              |  |  |  |  |  |
| (ii)  | Place a spatula<br>measure of solid in a<br>boiling tube and add<br>a 2 cm depth of dilute<br>sulfuric acid. |                                   |                   |  |  |  |  |  |
|       | Keep the s   | olutions formed in (ii) for tests | s (iii) and (iv). |  |  |  |  |  |
| (iii) | To a 1 cm depth of<br>solution from (ii)<br>in a test-tube, add<br>aqueous sodium<br>hydroxide.              |                                   |                   |  |  |  |  |  |
| (iv)  | To a 1 cm depth of<br>solution from (ii)<br>in a test-tube, add<br>aqueous ammonia.                          |                                   |                   |  |  |  |  |  |

(v) Identify as many ions as you can from your observations. Write 'unknown' where you have not been able to identify an ion.

| FA 5: cation | anion |
|--------------|-------|
| FA 6: cation | anion |

(vi) Write an equation, including state symbols, for the reaction between **FA 6** and dilute sulfuric acid.

[12]

- (b) **FA 7** is a solution containing one anion from those listed on page 11. The anion is either a halide or contains nitrogen.
  - (i) You are to select suitable reagents to determine the identity of this anion. Record these in a suitable form below.

(ii) Use these reagents to carry out tests to identify the anion in FA 7.

Record your observations and conclusions in the space below.

## **Qualitative Analysis Notes**

## *Key: [ppt. = precipitate]*

# 1 Reactions of aqueous cations

| ion                                 | react  | ion with   |
|-------------------------------------|--|--|
| 1011                                | NaOH(aq)   | NH <sub>3</sub> (aq)   |
| aluminium,<br>Al <sup>3+</sup> (aq) | white ppt.<br>soluble in excess  | white ppt.<br>insoluble in excess  |
| ammonium,<br>NH₄⁺(aq)               | no ppt.<br>ammonia produced on heating   | _  |
| barium,<br>Ba²⁺(aq)                 | no ppt. (if reagents are pure)   | no ppt.  |
| calcium,<br>Ca²+(aq)                | white ppt. with high [Ca²+(aq)]  | no ppt.  |
| chromium(III),<br>Cr³⁺(aq)          | grey-green ppt.<br>soluble in excess   | grey-green ppt.<br>insoluble in excess   |
| copper(II),<br>Cu²+(aq)             | pale blue ppt.<br>insoluble in excess  | blue ppt. soluble in excess giving dark blue solution                              |
| iron(II),<br>Fe²+(aq)               | green ppt. turning brown on contact<br>with air<br>insoluble in excess             | green ppt. turning brown on contact<br>with air<br>insoluble in excess             |
| iron(III),<br>Fe³+(aq)              | red-brown ppt.<br>insoluble in excess  | red-brown ppt.<br>insoluble in excess  |
| magnesium,<br>Mg²⁺(aq)              | white ppt.<br>insoluble in excess  | white ppt.<br>insoluble in excess  |
| manganese(II),<br>Mn²⁺(aq)          | off-white ppt. rapidly turning brown<br>on contact with air<br>insoluble in excess | off-white ppt. rapidly turning brown<br>on contact with air<br>insoluble in excess |
| zinc,<br>Zn²+(aq)                   | white ppt.<br>soluble in excess  | white ppt.<br>soluble in excess  |

## 2 Reactions of anions

| ion  | reaction  |
|--|---|
| carbonate,<br>CO <sub>3</sub> <sup>2–</sup>    | CO <sub>2</sub> liberated by dilute acids   |
| chloride,<br>C <i>l</i> ⁻(aq)                  | gives white ppt. with Ag⁺(aq) (soluble in NH₃(aq))  |
| bromide,<br>Br⁻(aq)                            | gives cream ppt. with Ag <sup>+</sup> (aq) (partially soluble in NH <sub>3</sub> (aq))  |
| iodide,<br>I⁻(aq)                              | gives yellow ppt. with Ag⁺(aq) (insoluble in NH₃(aq))   |
| nitrate,<br>NO₃⁻(aq)                           | NH <sub>3</sub> liberated on heating with OH⁻(aq) and A <i>l</i> foil   |
| nitrite,<br>NO₂⁻(aq)                           | $NH_3$ liberated on heating with OH <sup>-</sup> (aq) and A <i>l</i> foil;<br>NO liberated by dilute acids<br>(colourless NO $\rightarrow$ (pale) brown NO <sub>2</sub> in air) |
| sulfate,<br>SO <sub>4</sub> ²-(aq)             | gives white ppt. with Ba <sup>2+</sup> (aq) (insoluble in excess dilute strong acids)   |
| sulfite,<br>SO <sub>3</sub> <sup>2–</sup> (aq) | gives white ppt. with Ba <sup>2+</sup> (aq) (soluble in excess dilute strong acids)   |

# 3 Tests for gases

| gas                             | test and test result  |
|---------------------------------|---|
| ammonia, NH <sub>3</sub>        | turns damp red litmus paper blue  |
| carbon dioxide, CO <sub>2</sub> | gives a white ppt. with limewater (ppt. dissolves with excess CO <sub>2</sub> ) |
| chlorine, $Cl_2$                | bleaches damp litmus paper  |
| hydrogen, H <sub>2</sub>        | "pops" with a lighted splint  |
| oxygen, O <sub>2</sub>          | relights a glowing splint   |

|     |       |  |   |  |   |  |  |   |  |   |   |   |  |  |   |   |   |   |  |  |  |  | ]   |  |  |  |  |   |   |   |  |                  |         |
|-----|-------|--|---|--|---|--|--|---|--|---|---|---|--|--|---|---|---|---|--|--|--|--|---|--|--|--|--|---|---|---|--|------------------|---------|
|     | 18    | 2  | He  | helium<br>4.0  | 10  | Ne   | neon<br>20.2   | 18  | Ar   | argon<br>39.9   | 36  | ŗ   | krypton<br>83.8  | 5  | Xe  | xenon<br>131.3  | 86  | Rn  | radon<br>-   |  |  |  |   |  |  |  |  |   |   |   |  |                  |         |
|     | 17    |  |   |  | 6   | ш  | fluorine<br>19.0   | 17  | Cl   | chlorine<br>35.5  | 35  | В   | bromine<br>79.9  | 53   | Ι   | iodine<br>126.9   | 85  | At  | astatine<br>-  |  |  |  |   | 71   | Lu   | Iutetium<br>175.0  | 103  | Ļ   | lawrencium  |   |  |                  |         |
|     | 16    |  |   |  | 80  | 0  | oxygen<br>16.0   | 16  | S  | sulfur<br>32.1  | 34  | Se  | selenium<br>79.0   | 52   | Те  | tellurium<br>127.6  | 84  | Ро  | polonium<br>–  | 116  | Ľ  | livermorium<br>-   |   | 70   | γb   | ytterbium<br>173.1   | 102  | No  | nobelium  |   |  |                  |         |
|     | 15    |  |   |  |   | 7  | z  | nitrogen<br>14.0  | 15   | ٩   | phosphorus<br>31.0  | 33  | As   | arsenic<br>74.9  | 51  | Sb  | antimony<br>121.8   | 83  | Bi   | bismuth<br>209.0   |  |  |   |  | 69   | Tm   | thulium<br>168.9   | 101   | рМ  | mendelevium   |  |                  |         |
|     | 14    |  |   |  |   | 9  | U  | carbon<br>12.0  | 14   | S.  | silicon<br>28.1   | 32  | Ge   | germanium<br>72.6  | 50  | Sn  | tin<br>118.7  | 82  | Рр   | lead<br>207.2  | 114  | Fl   | flerovium<br>–  |  | 68   | ш  | erbium<br>167.3  | 100   | Еm  | fermium   |  |                  |         |
|     | 13    | -  |   | 5  | В   | boron<br>10.8  | 13   | Al  | aluminium<br>27.0  | 31  | Ga  | gallium<br>69.7   | 49   | In   | indium<br>114.8   | 81  | 11  | thallium<br>204.4                                     |  |  |  |  | 67  | Ч  | holmium<br>164.9   | 66   | Es   | einsteinium   |   |   |  |                  |         |
|     |       |  |   |  |   |  |  |   |  |   |   | 12  | 30   | Zn   | zinc<br>65.4  | 48  | Сq  | cadmium<br>112.4                                      | 80   | Hg   | mercury<br>200.6   | 112  | ő   | copernicium<br>-   |  | 99   | Dy   | dysprosium<br>162.5   | 98  | Ç   | californium  |                  |         |
|     |       |  |   |  |   |  |  |   |  |   |   |   |  |  |   | 11  | 29  | Cu  | copper<br>63.5   | 47   | Ag   | silver<br>107.9  | 79  | Au   | gold<br>197.0  | 111  | Rg   | roentgenium<br>-  |   | 65  | Tb   | terbium<br>158.9 | 97      |
| dn  |       |  |   |  |   |  |  |   |  | 10  | 28  | ïZ  | nickel<br>58.7   | 46   | Pd  | palladium<br>106.4  | 78  | ħ   | platinum<br>195.1  | 110  | Ds   | darmstadtium<br>-  |   | 64   | Gd   | gadolinium<br>157.3  | 96   | CB  | curium  |   |  |                  |         |
| Gro |       |  |   |  |   |  |  |   |  | o   | 27  | ပိ  | cobalt<br>58.9   | 45   | Rh  | rhodium<br>102.9  | 77  | Ir  | iridium<br>192.2   | 109  | Mt   | meitnerium<br>-  |   | 63   | Eu   | europium<br>152.0  | 95   | Am  | americium   |   |  |                  |         |
|     |       | -  | т   | hydrogen<br>1.0  |   |  |  |   |  | 80  | 26  | Fe  | iron<br>55.8   | 44   | Ru  | ruthenium<br>101.1  | 76  | Os  | osmium<br>190.2  | 108  | Hs   | hassium<br>-   |   | 62   | Sm   | samarium<br>150.4  | 94   | Pu  | plutonium   |   |  |                  |         |
|     |       |  |   |  | 1   |  |  |   |  | 7   | 25  | Mn  |  |  |   |   |   | Re  | rhenium<br>186.2   | 107  | Bh   | bohrium<br>I   |   | 61   | Pm   | promethium<br>-  | 93   | dN  | neptunium   |   |  |                  |         |
|     |       |  |   |  |   | loc  | ISS  |   |  | 9   | 24  | ŗ   | chromium<br>52.0   | 42   | Mo  | molybdenum<br>95.9  | 74  | S   | tungsten<br>183.8  | 106  | Sg   | seaborgium<br>-  |   |  | ΡN   | neodymium<br>144.4   | 92   | ⊃   | uranium   |   |  |                  |         |
|     |       |  |   |  |   | Key  | atomic number  | mic syml  | name<br>tive atomic ma   |   | L   | 5   | 23   | >  |   |   |   |   |  | Ta   | tantalum<br>180.9  | 105  | Db  | dubnium<br>–   |  | 59   | P  | praseodymium<br>140.9   | 91  | Ра  | protactinium   |                  |         |
|     |       |  |   |  |   |  |  |   |  | ato   | rela  |   |  | 4  | 22  | F   | titanium<br>47.9  | 40  | Zr   | zirconium<br>91.2  | 72   | Ηf   | hafnium<br>178.5  | 104  | Ŗ  | rutherfordium<br>-   |  | 58  |   |   |  | Тh               | thorium |
|     |       |  |   |  |   |  |  | -   |  | ю   | 21  | Sc  | scandium<br>45.0   | 39   | ≻   | yttrium<br>88.9   | 57-71   | lanthanoids   |  | 89-103   | actinoids  |  |   | 57   | La   | lanthanum<br>138.9   | 89   | Ac  | actinium  |   |  |                  |         |
|     | 2     |  |   |  | 4   | Be   | beryllium<br>9.0   | 12  | Mg   | magnesium<br>24.3                                       | 20  | Ca  | calcium<br>40.1  | 38   | S   | strontium<br>87.6   | 56  | Ba  | barium<br>137.3  | 88   | Ra   | radium<br>-  |   |  | ids  |  |  |   |   |   |  |                  |         |
|     | 1     |  |   |  | e   |  | lithium<br>6.9   | 11  | Na   | sodium<br>23.0  | 19  | ¥   | potassium<br>39.1  | 37   | Rb  | rubidium<br>85.5  | 55  | Cs  | caesium<br>132.9   | 87   | ŗ  | francium<br>-  |   |  | lanthano   |  |  | actinoids   |   |   |  |                  |         |
|     | Group | Group     13     14     15     16     17 | Group     13     14     15     16       1     1     1     1 | 2     13     14     15     16     17       1     1     1     1     1     1 | Group     13     14     15     16       1     1     1     1     1       Key     Nydrogen<br>1.0     1.0     1.0     1.0 | Group<br>A definition of the second sec | Group<br>Group<br>A to be atomic symbol<br>B to be atomic symbol<br>B to be atomic symbol<br>C N 0 0<br>F atomic symbol<br>C N 0<br>C N | Group     13     14     15     16     17       Rey     Natrogen<br>1.0     13     14     15     16     17       Rey     Natrogen<br>1.0     10     13     14     15     16     17       Be     Admin number<br>1.0     atomic number<br>1.0     Be     7     8     9       Portiogen<br>1200     1200     1200     1400     600     7     8 | Group     13     14     15     16     17       Rey     Natiogen<br>hydrogen<br>beyllum     13     14     15     16     17       Rey     Natiogen<br>hydrogen<br>beyllum     10     13     14     15     16     17       12     12     13     14     15     16     17 | $ \begin{array}{c c c c c c c c c c c c c c c c c c c $ | Group21314151617213141516174101010101710berylum<br>90anonc number<br>105678912121401516171712121201401501401719012121201101214015019020131312567891027,028101310281013102810100002810 | Image: Second | $\begin{tabular}{ c c c c c c c c c c c c c c c c c c c$ | Group222 $\frac{4}{13}$ 444131314131414151516171113141415151617111213141516171718191111121314141515161717181919111112131415151617171819191919191919191119111112131415151617171819< | Group           2         Important           Group           Important           A Mey           Mey           Important           Important | Group         2       13       14       15       16       17         Key       13       14       15       16       17         Field       100mic number         900m       900m       90       9       9       10       11       17       17         12       atomic number       10       11       12       14       15       16       17         12       atomic number       10       11       12       14       15       16       17         12       atomic number       11       11       12       14       15       16       17         13       14       5       6       7       8       9       10       11       12       14       16       17         13       14       5       6       7       8       9       14       16       17         13       14       12       14       12       14       16       17         14       15       16       11       12       14       16       17         14       5       6       7       8       9< | GroupI II | Croup           I I I I I I I I I I I I I I I I I I I | Induction in the transmission of the transmission of the transmission of the transmission of transmissin of transmission of transmissing of transm | Image: constraint of the | I Carbon       I Carbon <th< td=""><td>Clock         Interpretation         <th colspa="&lt;/td"><td>Group         Idea         Idea</td><td>Interpretation in the second second</td><td>I CIOID         2       I CIOID         2       I CIOID         I CIOID         VIENTIAL         <th colsp<="" td=""><td>Interpretation         Group           2         Total         13         14         15           For an intermediation in the proper and the proper proproproper proper proper proper proper proper proper p</td><td>Gloup         Idea           3         A         13         14         15         16         17         3         4         15         16         17         16         17         16         17         16         17         16         17         16         17         16         17         16         17         16         17         16         17         16         17         16         17         16         17         16         17         16         17         16         17         16         17         16         17         17         16         17         17         17         17         17         17         17         17         17         17         17</td><td>Glop         Identified in the formation of the for</td><td>2         Clop         13         14         15         16         17           2         A         Key         Image: Simple in the simple inthe simple inthe simple in the simple in the simple inthe simple i</td></th></td></th></td></th<> | Clock         Interpretation         Interpretation <th colspa="&lt;/td"><td>Group         Idea         Idea</td><td>Interpretation in the second second</td><td>I CIOID         2       I CIOID         2       I CIOID         I CIOID         VIENTIAL         <th colsp<="" td=""><td>Interpretation         Group           2         Total         13         14         15           For an intermediation in the proper and the proper proproproper proper proper proper proper proper proper p</td><td>Gloup         Idea           3         A         13         14         15         16         17         3         4         15         16         17         16         17         16         17         16         17         16         17         16         17         16         17         16         17         16         17         16         17         16         17         16         17         16         17         16         17         16         17         16         17         16         17         16         17         17         16         17         17         17         17         17         17         17         17         17         17         17</td><td>Glop         Identified in the formation of the for</td><td>2         Clop         13         14         15         16         17           2         A         Key         Image: Simple in the simple inthe simple inthe simple in the simple in the simple inthe simple i</td></th></td></th> | <td>Group         Idea         Idea</td> <td>Interpretation in the second second</td> <td>I CIOID         2       I CIOID         2       I CIOID         I CIOID         VIENTIAL         <th colsp<="" td=""><td>Interpretation         Group           2         Total         13         14         15           For an intermediation in the proper and the proper proproproper proper proper proper proper proper proper p</td><td>Gloup         Idea           3         A         13         14         15         16         17         3         4         15         16         17         16         17         16         17         16         17         16         17         16         17         16         17         16         17         16         17         16         17         16         17         16         17         16         17         16         17         16         17         16         17         16         17         16         17         17         16         17         17         17         17         17         17         17         17         17         17         17</td><td>Glop         Identified in the formation of the for</td><td>2         Clop         13         14         15         16         17           2         A         Key         Image: Simple in the simple inthe simple inthe simple in the simple in the simple inthe simple i</td></th></td> | Group         Idea         Idea | Interpretation in the second | I CIOID         2       I CIOID         2       I CIOID         I CIOID         VIENTIAL         VIENTIAL <th colsp<="" td=""><td>Interpretation         Group           2         Total         13         14         15           For an intermediation in the proper and the proper proproproper proper proper proper proper proper proper p</td><td>Gloup         Idea           3         A         13         14         15         16         17         3         4         15         16         17         16         17         16         17         16         17         16         17         16         17         16         17         16         17         16         17         16         17         16         17         16         17         16         17         16         17         16         17         16         17         16         17         16         17         17         16         17         17         17         17         17         17         17         17         17         17         17</td><td>Glop         Identified in the formation of the for</td><td>2         Clop         13         14         15         16         17           2         A         Key         Image: Simple in the simple inthe simple inthe simple in the simple in the simple inthe simple i</td></th> | <td>Interpretation         Group           2         Total         13         14         15           For an intermediation in the proper and the proper proproproper proper proper proper proper proper proper p</td> <td>Gloup         Idea           3         A         13         14         15         16         17         3         4         15         16         17         16         17         16         17         16         17         16         17         16         17         16         17         16         17         16         17         16         17         16         17         16         17         16         17         16         17         16         17         16         17         16         17         16         17         17         16         17         17         17         17         17         17         17         17         17         17         17</td> <td>Glop         Identified in the formation of the for</td> <td>2         Clop         13         14         15         16         17           2         A         Key         Image: Simple in the simple inthe simple inthe simple in the simple in the simple inthe simple i</td> | Interpretation         Group           2         Total         13         14         15           For an intermediation in the proper and the proper proproproper proper proper proper proper proper proper p | Gloup         Idea           3         A         13         14         15         16         17         3         4         15         16         17         16         17         16         17         16         17         16         17         16         17         16         17         16         17         16         17         16         17         16         17         16         17         16         17         16         17         16         17         16         17         16         17         16         17         17         16         17         17         17         17         17         17         17         17         17         17         17 | Glop         Identified in the formation of the for | 2         Clop         13         14         15         16         17           2         A         Key         Image: Simple in the simple inthe simple inthe simple in the simple in the simple inthe simple i |                  |         |

1

i

1

i.

i.

i.

I

238.0

231.0

232.0

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cie.org.uk after the live examination series.