



Cambridge International AS & A Level

CANDIDATE
NAME

--

CENTRE
NUMBER

--	--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--

* 8 2 0 5 0 4 4 8 0 4 *



CHEMISTRY

9701/33

Paper 3 Advanced Practical Skills 1

October/November 2022

2 hours

You must answer on the question paper.

You will need: The materials and apparatus listed in the confidential instructions

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- You may use a calculator.
- You should show all your working and use appropriate units.

INFORMATION

- The total mark for this paper is 40.
- The number of marks for each question or part question is shown in brackets [].
- The Periodic Table is printed in the question paper.
- Important values, constants and standards are printed in the question paper.
- Notes for use in qualitative analysis are provided in the question paper.

Session
Laboratory

For Examiner's Use	
1	
2	
3	
Total	

This document has **16** pages. Any blank pages are indicated.

Quantitative analysis


Read through the whole method before starting any practical work. Where appropriate, prepare a table for your results in the space provided.

Show the precision of the apparatus you used in the data you record.

Show your working and appropriate significant figures in the final answer to **each** step of your calculations.

1 An iron compound can contain Fe²⁺ ions, Fe³⁺ ions or both ions.

In this experiment, you will determine the percentage by mass of iron in **FA 1**, an unknown compound. You will first prepare a solution of the compound and then carry out a titration using acidified potassium manganate(VII), KMnO₄.



The end-point of the titration is when all the Fe²⁺ ions have been oxidised and so unreacted potassium manganate(VII) causes the colour of the solution to become a permanent pale pink.

FA 1 is a sample of the unknown compound.

FA 2 is 0.0100 mol dm⁻³ potassium manganate(VII), KMnO₄.

FA 3 is dilute sulfuric acid, H₂SO₄.

(a) Method

Weighing the compound

- Weigh the sample of **FA 1** and its container.
- Tip the **FA 1** into the 250 cm³ beaker.
- Reweigh the container including any residual **FA 1**.
- Record both your readings clearly in the space below.
- Calculate the mass of **FA 1** transferred into the beaker. Record the mass.

Preparing the solution

- Add approximately 200 cm³ of distilled water to the beaker and stir until the **FA 1** has dissolved.
- Pour the contents carefully into the 250 cm³ volumetric flask.
- Rinse the contents of the beaker with a little distilled water and add these washings to the flask.
- Fill the flask to the line with distilled water and shake thoroughly.
- Label this solution **FA 4**.

Titration

- Fill the burette with **FA 2**.
- Pipette 25.0 cm³ of **FA 4** into a conical flask.
- Use the 25 cm³ measuring cylinder to add 15 cm³ of **FA 3** to the conical flask.
- Add **FA 2** from the burette until the solution in the flask turns a permanent pink.
- Carry out a rough titration and record your burette readings in the space below.

rough titre = cm³

- Carry out as many accurate titrations as you think are necessary to obtain consistent results.
- Make sure that your recorded results show the precision of your practical work.
- Record in a suitable format, in the space below, all your burette readings and the volume of **FA 2** added in each accurate titration.

I	
II	
III	
IV	
V	
VI	
VII	
VIII	

[8]

- (b) From your accurate titration results, calculate a suitable mean value to use in your calculations. Show clearly how you obtained this value.

25.0 cm³ of **FA 4** required cm³ of **FA 2**. [1]

(c) Calculations

(i) Give your answers to **(c)(ii)**, **(c)(iii)** and **(c)(iv)** to the appropriate number of significant figures. [1]

(ii) Calculate the amount, in mol, of manganate(VII) ions in the volume recorded in **(b)**.

amount of MnO_4^- = mol [1]

(iii) Calculate the amount, in mol, of iron(II) ions in the weighed sample of **FA 1**.

amount of Fe^{2+} = mol [1]

(iv) Calculate the percentage by mass of iron in **FA 1**. Show your working.

percentage by mass of iron = % [2]

(d) A student carries out the same experiment as in **(a)**. The student receives a sample of **FA 1** in a container with a lid. The student records the initial mass of the container with its lid and the sample of **FA 1**. Then the student records the mass of the container with the residue but forgets to replace the lid.

How would this error alter the student's answer to **(c)(iv)**?
Explain your answer.

.....
..... [1]

(e) State **two** assumptions that have been made in calculating the percentage by mass of iron in **FA 1** in **(c)(iv)**.

.....
.....
..... [2]

[Total: 17]

Question 2 starts on the next page.

- 2 In this experiment you will determine the percentage by mass of magnesium in a hydrated salt, $\text{MgX}\cdot 7\text{H}_2\text{O}$, where **X** represents the anion.

You will measure the loss of mass when a sample of the hydrated salt is heated to form the anhydrous salt.

FA 5 is a pure sample of $\text{MgX}\cdot 7\text{H}_2\text{O}$

(a) Method

- Weigh the crucible with its lid. Record the mass.
- Add all of **FA 5** to the crucible.
- Reweigh the crucible with its lid and **FA 5**. Record the mass.
- Support the crucible in the pipe-clay triangle on top of the tripod.
- Heat the crucible with the lid on **gently** for about 1 minute.
- Remove the lid and then heat strongly for a further 4 minutes.
- Replace the lid and allow the crucible to cool for at least 5 minutes.

While the crucible is cooling you may wish to begin work on Question 3.

- When the crucible is cool enough to handle, reweigh the crucible with its lid and its contents. Record the mass.
- Calculate and record the mass of **FA 5** used.
- Calculate and record the mass of water lost.

I	
II	
III	
IV	
V	

[5]

(b) Calculations

- (i) Calculate the amount, in mol, of water lost.

amount of H_2O lost = mol [1]

(ii) Calculate the percentage by mass of magnesium in **FA 5**. Show your working.

percentage by mass of Mg = % [2]

(c) Suggest **two** assumptions that must be made for this experiment to give an accurate value of the percentage of Mg.

.....
.....
.....
..... [2]

[Total: 10]

Qualitative analysis

For each test you should record **all** your observations in the spaces provided.

Examples of observations include:

- colour changes seen
- the formation of any precipitate and its solubility (where appropriate) in an excess of the reagent added
- the formation of any gas and its identification (where appropriate) by a suitable test.

You should record clearly at what stage in a test an observation is made.

Where no change is observed you should write 'no change'.

Where reagents are selected for use in a test, the name or correct formula of the element or compound must be given.

If any solution is warmed, a boiling tube must be used.

Rinse and reuse test-tubes and boiling tubes where possible.

No additional tests should be attempted.

3 (a) FA 6 is an acidified aqueous solution of a salt which contains two cations and one anion, all of which are listed in the Qualitative analysis notes.

- (i) Select a reagent or reagents for use in tests to identify the two cations.
Record your observations.

[2]

- (ii) The anion in **FA 6** is either the sulfate ion, SO_4^{2-} , or the sulfite ion, SO_3^{2-} .
Select a reagent or reagents for use in tests to identify the anion.
Record your observations.

[2]

- (iii) Identify the ions present in **FA 6** and give their formulae.

The cations present in **FA 6** are and

The anion present in **FA 6** is

[2]

- (b) Carry out the tests and record your observations.
For each test use a 1 cm depth of **FA 6** in a test-tube.

<i>test</i>	<i>observations</i>
Test 1 Add a 1 cm depth of aqueous sodium sulfite, then	
add a 1 cm depth of dilute sulfuric acid. Rinse out this test-tube thoroughly.	
Test 2 Add a 1 cm depth of aqueous potassium iodide, then	
add a few drops of starch solution.	

[2]

- (c) **FA 7**, **FA 8** and **FA 9** are aqueous solutions of sodium salts. Each solution contains one of the following ions: CO_3^{2-} , Cl^- , Br^- .

You will identify which solution contains which ion.

- (i) Carry out the tests and record your observations.
For each test use a 1 cm depth of **FA 7**, **FA 8** or **FA 9** in a test-tube.

test	observations		
	FA 7	FA 8	FA 9
Add a few drops of aqueous silver nitrate, then			
add aqueous ammonia.			

[3]

- (ii) If your results are insufficient to identify which anion is present in each solution, carry out a further test. Record your test and observations.

[1]

- (iii) Identify which solution contains which ion.

The CO_3^{2-} ion is present in

The Cl^- ion is present in

The Br^- ion is present in

[1]

[Total: 13]

BLANK PAGE

BLANK PAGE

BLANK PAGE

Qualitative analysis notes

1 Reactions of cations

cation	reaction with	
	NaOH(aq)	NH ₃ (aq)
aluminium, Al ³⁺ (aq)	white ppt. soluble in excess	white ppt. insoluble in excess
ammonium, NH ₄ ⁺ (aq)	no ppt. ammonia produced on warming	–
barium, Ba ²⁺ (aq)	faint white ppt. is observed unless [Ba ²⁺ (aq)] is very low	no ppt.
calcium, Ca ²⁺ (aq)	white ppt. unless [Ca ²⁺ (aq)] is very low	no ppt.
chromium(III), Cr ³⁺ (aq)	grey-green ppt. soluble in excess giving dark green solution	grey-green ppt. insoluble in excess
copper(II), Cu ²⁺ (aq)	pale blue ppt. insoluble in excess	pale blue ppt. soluble in excess giving dark blue solution
iron(II), Fe ²⁺ (aq)	green ppt. turning brown on contact with air insoluble in excess	green ppt. turning brown on contact with air insoluble in excess
iron(III), Fe ³⁺ (aq)	red-brown ppt. insoluble in excess	red-brown ppt. insoluble in excess
magnesium, Mg ²⁺ (aq)	white ppt. insoluble in excess	white ppt. insoluble in excess
manganese(II), Mn ²⁺ (aq)	off-white ppt. rapidly turning brown on contact with air insoluble in excess	off-white ppt. rapidly turning brown on contact with air insoluble in excess
zinc, Zn ²⁺ (aq)	white ppt. soluble in excess	white ppt. soluble in excess

2 Reactions of anions

anion	reaction
carbonate, CO ₃ ²⁻	CO ₂ liberated by dilute acids
chloride, Cl ⁻ (aq)	gives white ppt. with Ag ⁺ (aq) (soluble in NH ₃ (aq))
bromide, Br ⁻ (aq)	gives cream/off-white ppt. with Ag ⁺ (aq) (partially soluble in NH ₃ (aq))
iodide, I ⁻ (aq)	gives pale yellow ppt. with Ag ⁺ (aq) (insoluble in NH ₃ (aq))
nitrate, NO ₃ ⁻ (aq)	NH ₃ liberated on heating with OH ⁻ (aq) and Al foil
nitrite, NO ₂ ⁻ (aq)	NH ₃ liberated on heating with OH ⁻ (aq) and Al foil; decolourises acidified aqueous KMnO ₄
sulfate, SO ₄ ²⁻ (aq)	gives white ppt. with Ba ²⁺ (aq) (insoluble in excess dilute strong acids); gives white ppt. with high [Ca ²⁺ (aq)]
sulfite, SO ₃ ²⁻ (aq)	gives white ppt. with Ba ²⁺ (aq) (soluble in excess dilute strong acids); decolourises acidified aqueous KMnO ₄
thiosulfate, S ₂ O ₃ ²⁻ (aq)	gives off-white/pale yellow ppt. slowly with H ⁺

3 Tests for gases

gas	test and test result
ammonia, NH ₃	turns damp red litmus paper blue
carbon dioxide, CO ₂	gives a white ppt. with limewater
hydrogen, H ₂	'pops' with a lighted splint
oxygen, O ₂	relights a glowing splint

4 Tests for elements

element	test and test result
iodine, I ₂	gives blue-black colour on addition of starch solution

Important values, constants and standards

molar gas constant	$R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$
Faraday constant	$F = 9.65 \times 10^4 \text{ C mol}^{-1}$
Avogadro constant	$L = 6.022 \times 10^{23} \text{ mol}^{-1}$
electronic charge	$e = -1.60 \times 10^{-19} \text{ C}$
molar volume of gas	$V_m = 22.4 \text{ dm}^3 \text{ mol}^{-1}$ at s.t.p. (101 kPa and 273 K) $V_m = 24.0 \text{ dm}^3 \text{ mol}^{-1}$ at room conditions
ionic product of water	$K_w = 1.00 \times 10^{-14} \text{ mol}^2 \text{ dm}^{-6}$ (at 298 K (25 °C))
specific heat capacity of water	$c = 4.18 \text{ kJ kg}^{-1} \text{ K}^{-1}$ (4.18 J g ⁻¹ K ⁻¹)

The Periodic Table of Elements

		Group																																		
1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18																			
		<div style="display: flex; justify-content: space-between; align-items: center;"> <div style="border: 1px solid black; padding: 5px;">1 H hydrogen 1.0</div> <div style="border: 1px solid black; padding: 5px;"> Key atomic number atomic symbol name relative atomic mass </div> </div>																																		
3 Li lithium 6.9	4 Be beryllium 9.0	11 Na sodium 23.0	12 Mg magnesium 24.3	19 K potassium 39.1	20 Ca calcium 40.1	21 Sc scandium 45.0	22 Ti titanium 47.9	23 V vanadium 50.9	24 Cr chromium 52.0	25 Mn manganese 54.9	26 Fe iron 55.8	27 Co cobalt 58.9	28 Ni nickel 58.7	29 Cu copper 63.5	30 Zn zinc 65.4	31 Ga gallium 69.7	32 Ge germanium 72.6	33 As arsenic 74.9	34 Se selenium 79.0	35 Br bromine 79.9	36 Kr krypton 83.8															
37 Rb rubidium 85.5	38 Sr strontium 87.6	39 Y yttrium 88.9	40 Zr zirconium 91.2	41 Nb niobium 92.9	42 Mo molybdenum 95.9	43 Tc technetium —	44 Ru ruthenium 101.1	45 Rh rhodium 102.9	46 Pd palladium 106.4	47 Ag silver 107.9	48 Cd cadmium 112.4	49 In indium 114.8	50 Sn tin 118.7	51 Sb antimony 121.8	52 Te tellurium 127.6	53 I iodine 126.9	54 Xe xenon 131.3	55 Cs caesium 132.9	56 Ba barium 137.3	57 La lanthanum 138.9	58 Ce cerium 140.1	59 Pr praseodymium 140.9	60 Nd neodymium 144.4	61 Pm promethium —	62 Sm samarium 150.4	63 Eu europium 152.0	64 Gd gadolinium 157.3	65 Tb terbium 158.9	66 Dy dysprosium 162.5	67 Ho holmium 164.9	68 Er erbium 167.3	69 Tm thulium 168.9	70 Yb ytterbium 173.1	71 Lu lutetium 175.0		
87 Fr francium —	88 Ra radium —	89 Ac actinium —	89–103 actinoids —	72 Hf hafnium 178.5	73 Ta tantalum 180.9	74 W tungsten 183.8	75 Re rhenium 186.2	76 Os osmium 190.2	77 Ir iridium 192.2	78 Pt platinum 195.1	79 Au gold 197.0	80 Hg mercury 200.6	81 Tl thallium 204.4	82 Pb lead 207.2	83 Bi bismuth 209.0	84 Po polonium —	85 At astatine —	86 Rn radon —	87 Fr francium —	88 Ra radium —	89–103 actinoids —	104 Rf rutherfordium —	105 Db dubnium —	106 Sg seaborgium —	107 Bh bohrium —	108 Hs hassium —	109 Mt meitnerium —	110 Ds darmstadtium —	111 Rg roentgenium —	112 Cn copernicium —	113 Nh nihonium —	114 Fl flerovium —	115 Mc moscovium —	116 Lv livermorium —	117 Ts tennessine —	118 Og oganesson —

lanthanoids

actinoids

57 La lanthanum 138.9	58 Ce cerium 140.1	59 Pr praseodymium 140.9	60 Nd neodymium 144.4	61 Pm promethium —	62 Sm samarium 150.4	63 Eu europium 152.0	64 Gd gadolinium 157.3	65 Tb terbium 158.9	66 Dy dysprosium 162.5	67 Ho holmium 164.9	68 Er erbium 167.3	69 Tm thulium 168.9	70 Yb ytterbium 173.1	71 Lu lutetium 175.0
89 Ac actinium —	90 Th thorium 232.0	91 Pa protactinium 231.0	92 U uranium 238.0	93 Np neptunium —	94 Pu plutonium —	95 Am americium —	96 Cm curium —	97 Bk berkelium —	98 Cf californium —	99 Es einsteinium —	100 Fm fermium —	101 Md mendelevium —	102 No nobelium —	103 Lr lawrencium —