



Cambridge International AS & A Level

CANDIDATE
NAME

--

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--



MATHEMATICS

9709/42

Paper 4 Mechanics

February/March 2022

1 hour 15 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.
- Where a numerical value for the acceleration due to gravity (g) is needed, use 10 m s^{-2} .

INFORMATION

- The total mark for this paper is 50.
- The number of marks for each question or part question is shown in brackets [].

This document has **16** pages. Any blank pages are indicated.

BLANK PAGE

1 A crane is used to raise a block of mass 600 kg vertically upwards at a constant speed through a height of 15 m. There is a resistance to the motion of the block, which the crane does 10 000 J of work to overcome.

(a) Find the total work done by the crane. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Given that the average power exerted by the crane is 12.5 kW, find the total time for which the block is in motion. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

2 A particle P is projected vertically upwards from horizontal ground with speed $u \text{ m s}^{-1}$. P reaches a maximum height of 20 m above the ground.

(a) Find the value of u . [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Find the total time for which P is at least 15 m above the ground. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

3 A car of mass m kg is towing a trailer of mass 300 kg down a straight hill inclined at 3° to the horizontal at a constant speed. There are resistance forces on the car and on the trailer, and the total work done against the resistance forces in a distance of 50 m is 40 000 J. The engine of the car is doing no work and the tow-bar is light and rigid.

(a) Find the value of m . [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

The resistance force on the trailer is 200 N.

(b) Find the tension in the tow-bar between the car and the trailer. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

4 The total mass of a cyclist and her bicycle is 70 kg. The cyclist is riding with constant power of 180 W up a straight hill inclined at an angle α to the horizontal, where $\sin \alpha = 0.05$. At an instant when the cyclist's speed is 6 m s^{-1} , her acceleration is -0.2 m s^{-2} . There is a constant resistance to motion of magnitude $F \text{ N}$.

(a) Find the value of F . [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....

.....

.....

5



Four coplanar forces act at a point. The magnitudes of the forces are 10 N , $F\text{ N}$, $G\text{ N}$ and $2F\text{ N}$. The directions of the forces are as shown in the diagram.

- (a) Given that the forces are in equilibrium, find the values of F and G . [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Given instead that $F = 3$, find the value of G for which the resultant of the forces is perpendicular to the 10 N force. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- 6** A cyclist starts from rest at a fixed point O and moves in a straight line, before coming to rest k seconds later. The acceleration of the cyclist at time t s after leaving O is $a \text{ m s}^{-2}$, where $a = 2t^{-\frac{1}{2}} - \frac{3}{5}t^{\frac{1}{2}}$ for $0 < t \leq k$.

- (a)** Find the value of k . [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (b)** Find the maximum speed of the cyclist. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (c) Find an expression for the displacement from O in terms of t . Hence find the total distance travelled by the cyclist from the time at which she reaches her maximum speed until she comes to rest. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

7 A bead, A , of mass 0.1 kg is threaded on a long straight rigid wire which is inclined at $\sin^{-1}\left(\frac{7}{25}\right)$ to the horizontal. A is released from rest and moves down the wire. The coefficient of friction between A and the wire is μ . When A has travelled 0.45 m down the wire, its speed is 0.6 m s^{-1} .

(a) Show that $\mu = 0.25$. [6]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

Another bead, B , of mass 0.5 kg is also threaded on the wire. At the point where A has travelled 0.45 m down the wire, it hits B which is instantaneously at rest on the wire. A is brought to instantaneous rest in the collision. The coefficient of friction between B and the wire is 0.275 .

(b) Find the time from when the collision occurs until A collides with B again. [6]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

Additional Page

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

BLANK PAGE

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.