

CAMBRIDGE INTERNATIONAL EXAMINATIONS
GCE Advanced Subsidiary Level

MARK SCHEME for the October/November 2012 series

9709 MATHEMATICS

9709/21

Paper 2, maximum raw mark 50

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the October/November 2012 series for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level components and some Ordinary Level components.

Page 2	Mark Scheme	Syllabus	Paper
	GCE AS LEVEL – October/November 2012	9709	21

Mark Scheme Notes

Marks are of the following three types:

M Method mark, awarded for a valid method applied to the problem. Method marks are not lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. Correct application of a formula without the formula being quoted obviously earns the M mark and in some cases an M mark can be implied from a correct answer.

A Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated method mark is earned (or implied).

B Mark for a correct result or statement independent of method marks.

- When a part of a question has two or more "method" steps, the M marks are generally independent unless the scheme specifically says otherwise; and similarly when there are several B marks allocated. The notation DM or DB (or dep*) is used to indicate that a particular M or B mark is dependent on an earlier M or B (asterisked) mark in the scheme. When two or more steps are run together by the candidate, the earlier marks are implied and full credit is given.
- The symbol ∇ implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A or B marks are given for correct work only. A and B marks are not given for fortuitously "correct" answers or results obtained from incorrect working.
- Note: B2 or A2 means that the candidate can earn 2 or 0.
B2/1/0 means that the candidate can earn anything from 0 to 2.

The marks indicated in the scheme may not be subdivided. If there is genuine doubt whether a candidate has earned a mark, allow the candidate the benefit of the doubt. Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored.

- Wrong or missing units in an answer should not lead to the loss of a mark unless the scheme specifically indicates otherwise.
- For a numerical answer, allow the A or B mark if a value is obtained which is correct to 3 s.f., or which would be correct to 3 s.f. if rounded (1 d.p. in the case of an angle). As stated above, an A or B mark is not given if a correct numerical answer arises fortuitously from incorrect working. For Mechanics questions, allow A or B marks for correct answers which arise from taking g equal to 9.8 or 9.81 instead of 10.

Page 3	Mark Scheme	Syllabus	Paper
	GCE AS LEVEL – October/November 2012	9709	21

The following abbreviations may be used in a mark scheme or used on the scripts:

AEF	Any Equivalent Form (of answer is equally acceptable)
AG	Answer Given on the question paper (so extra checking is needed to ensure that the detailed working leading to the result is valid)
BOD	Benefit of Doubt (allowed when the validity of a solution may not be absolutely clear)
CAO	Correct Answer Only (emphasising that no "follow through" from a previous error is allowed)
CWO	Correct Working Only – often written by a 'fortuitous' answer
ISW	Ignore Subsequent Working
MR	Misread
PA	Premature Approximation (resulting in basically correct work that is insufficiently accurate)
SOS	See Other Solution (the candidate makes a better attempt at the same question)
SR	Special Ruling (detailing the mark to be given for a specific wrong solution, or a case where some standard marking practice is to be varied in the light of a particular circumstance)

Penalties

MR –1	A penalty of MR –1 is deducted from A or B marks when the data of a question or part question are genuinely misread and the object and difficulty of the question remain unaltered. In this case all A and B marks then become "follow through ✓" marks. MR is not applied when the candidate misreads his own figures – this is regarded as an error in accuracy. An MR–2 penalty may be applied in particular cases if agreed at the coordination meeting.
PA –1	This is deducted from A or B marks in the case of premature approximation. The PA –1 penalty is usually discussed at the meeting.

Page 4	Mark Scheme	Syllabus	Paper
	GCE AS LEVEL – October/November 2012	9709	21

- 1 EITHER** State or imply non-modular inequality $(x-2)^2 \geq (x+5)^2$, or corresponding equation or pair of linear equations M1
 Obtain critical value $-\frac{3}{2}$ A1
 State correct answer $x \leq -\frac{3}{2}$ A1
- OR** State a correct linear equation for the critical value, e.g. $x-2 = -x-5$, or corresponding correct linear inequality, e.g. $x-2 \geq -x-5$ M1
 Obtain critical value $-\frac{3}{2}$ A1
 State correct answer $x \leq -\frac{3}{2}$ A1 [3]
- 2** Use law for the logarithm of a product, a quotient or a power M1*
 Obtain $x \log 5 = (2x-1) \log 3$ or equivalent A1
 Solve for x M1(dep*)
 Obtain answer $x = 1.87$ A1 [4]
- 3** Make relevant use of the $\cos 2\theta$ formula M1
 Obtain a correct quadratic in $\cos \theta$ A1
 Solve a quadratic in $\cos \theta$ M1
 Obtain answer $\theta = 60$ and no others in the range A1 [4]
 (Ignore answers outside the given range)
- 4 (i)** State $\frac{dx}{dt} = \frac{-2}{1-2t}$ or $\frac{dy}{dt} = -2t^{-2}$ B1
 Use $\frac{dy}{dx} = \frac{dy}{dt} \div \frac{dx}{dt}$ M1
 Obtain given answer correctly A1 [3]
- (ii)** Equate derivative to 3 and solve for t M1
 State or imply that $t = -1$ c.w.o. A1
 Obtain coordinates $(\ln 3, -2)$ A1 [3]

Page 5	Mark Scheme	Syllabus	Paper
	GCE AS LEVEL – October/November 2012	9709	21

- 5 (i) Attempt to integrate and use limits θ and π
Obtain $1 - \sin \theta$ M1
A1 [2]
- (ii) State that area of rectangle = $\theta \cos \theta$, equate area of rectangle to area of R
and rearrange to given equation B1 [1]
- (iii) Use the iterative formula correctly at least once M1
Obtain final answer 0.56 A1
Show sufficient iterations to justify its accuracy to 2 d.p. or show there is a
sign change in the interval (0.555, 0.565) B1 [3]
- 6 (a) State or imply correct ordinates 0.125, 0.08743..., 0.21511... B1
Use correct formula, or equivalent, correctly with $h = 0.5$ and three ordinates M1
Obtain answer 0.11 with no errors seen A1 [3]
- (b) Attempt to expand brackets and divide by e^{2x} M1
Integrate a term of form ke^{-x} or ke^{-2x} correctly A1[√]
Obtain 2 correct terms A1
Fully correct integral $x + 4e^{-x} - 2e^{-2x} + c$ A1 [4]
- 7 (i) Substitute $x = -1$, equate to zero and obtain a correct equation in any form B1
Substitute $x = 3$ and equate to 12 M1
Obtain a correct equation in any form A1
Solve a relevant pair of equations for a or for b M1
Obtain $a = -4$ and $b = 6$ A1 [5]
- (ii) Attempt division by $x^2 - 2$ and reach a partial quotient of $2x - k$ M1
Obtain quotient $2x - 4$ A1
Obtain remainder -2 A1 [3]
- 8 (i) Differentiate using chain or quotient rule M1
Obtain derivative in any correct form A1
Obtain given answer correctly A1 [3]
- (ii) Differentiate using product rule M1
State derivative of $\tan \theta = \sec^2 \theta$ B1
Use trig identity $1 + \tan^2 \theta = \sec^2 \theta$ correctly M1
Obtain $2\sec^3 \theta - \sec \theta$ A1 [4]
- (iii) Use $\tan^2 x = \sec^2 \theta - 1$ to integrate $\tan^2 x$ M1
Obtain $3\sec \theta$ from integration of $3\sec \theta \tan \theta$ B1
Obtain $\tan \theta - 3\sec \theta$ A1
Attempt to substitute limits, using exact values M1
Obtain answer $4 - 3\sqrt{2}$ A1 [5]