



Cambridge International AS Level

MATHEMATICS

9709/22

Paper 2 Pure Mathematics 2

October/November 2022

MARK SCHEME

Maximum Mark: 50

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the October/November 2022 series for most Cambridge IGCSE™, Cambridge International A and AS Level components and some Cambridge O Level components.

This document consists of **12** printed pages.

PUBLISHED**Generic Marking Principles**

These general marking principles must be applied by all examiners when marking candidate answers. They should be applied alongside the specific content of the mark scheme or generic level descriptors for a question. Each question paper and mark scheme will also comply with these marking principles.

GENERIC MARKING PRINCIPLE 1:

Marks must be awarded in line with:

- the specific content of the mark scheme or the generic level descriptors for the question
- the specific skills defined in the mark scheme or in the generic level descriptors for the question
- the standard of response required by a candidate as exemplified by the standardisation scripts.

GENERIC MARKING PRINCIPLE 2:

Marks awarded are always **whole marks** (not half marks, or other fractions).

GENERIC MARKING PRINCIPLE 3:

Marks must be awarded **positively**:

- marks are awarded for correct/valid answers, as defined in the mark scheme. However, credit is given for valid answers which go beyond the scope of the syllabus and mark scheme, referring to your Team Leader as appropriate
- marks are awarded when candidates clearly demonstrate what they know and can do
- marks are not deducted for errors
- marks are not deducted for omissions
- answers should only be judged on the quality of spelling, punctuation and grammar when these features are specifically assessed by the question as indicated by the mark scheme. The meaning, however, should be unambiguous.

GENERIC MARKING PRINCIPLE 4:

Rules must be applied consistently, e.g. in situations where candidates have not followed instructions or in the application of generic level descriptors.

GENERIC MARKING PRINCIPLE 5:

Marks should be awarded using the full range of marks defined in the mark scheme for the question (however; the use of the full mark range may be limited according to the quality of the candidate responses seen).

GENERIC MARKING PRINCIPLE 6:

Marks awarded are based solely on the requirements as defined in the mark scheme. Marks should not be awarded with grade thresholds or grade descriptors in mind.

Mathematics Specific Marking Principles	
1	Unless a particular method has been specified in the question, full marks may be awarded for any correct method. However, if a calculation is required then no marks will be awarded for a scale drawing.
2	Unless specified in the question, answers may be given as fractions, decimals or in standard form. Ignore superfluous zeros, provided that the degree of accuracy is not affected.
3	Allow alternative conventions for notation if used consistently throughout the paper, e.g. commas being used as decimal points.
4	Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored (isw).
5	Where a candidate has misread a number in the question and used that value consistently throughout, provided that number does not alter the difficulty or the method required, award all marks earned and deduct just 1 mark for the misread.
6	Recovery within working is allowed, e.g. a notation error in the working where the following line of working makes the candidate's intent clear.

Mark Scheme Notes

The following notes are intended to aid interpretation of mark schemes in general, but individual mark schemes may include marks awarded for specific reasons outside the scope of these notes.

Types of mark

- M** Method mark, awarded for a valid method applied to the problem. Method marks are not lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. Correct application of a formula without the formula being quoted obviously earns the M mark and in some cases an M mark can be implied from a correct answer.
- A** Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated method mark is earned (or implied).
- B** Mark for a correct result or statement independent of method marks.
- DM or DB** When a part of a question has two or more ‘method’ steps, the M marks are generally independent unless the scheme specifically says otherwise; and similarly, when there are several B marks allocated. The notation DM or DB is used to indicate that a particular M or B mark is dependent on an earlier M or B (asterisked) mark in the scheme. When two or more steps are run together by the candidate, the earlier marks are implied and full credit is given.
- FT** Implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A or B marks are given for correct work only.
- A or B marks are given for correct work only (not for results obtained from incorrect working) unless follow through is allowed (see abbreviation FT above).
 - For a numerical answer, allow the A or B mark if the answer is correct to 3 significant figures or would be correct to 3 significant figures if rounded (1 decimal place for angles in degrees).
 - The total number of marks available for each question is shown at the bottom of the Marks column.
 - Wrong or missing units in an answer should not result in loss of marks unless the guidance indicates otherwise.
 - Square brackets [] around text or numbers show extra information not needed for the mark to be awarded.

Abbreviations

AEF/OE	Any Equivalent Form (of answer is equally acceptable) / Or Equivalent
AG	Answer Given on the question paper (so extra checking is needed to ensure that the detailed working leading to the result is valid)
CAO	Correct Answer Only (emphasising that no ‘follow through’ from a previous error is allowed)
CWO	Correct Working Only
ISW	Ignore Subsequent Working
SOI	Seen Or Implied
SC	Special Case (detailing the mark to be given for a specific wrong solution, or a case where some standard marking practice is to be varied in the light of a particular circumstance)
WWW	Without Wrong Working
AWRT	Answer Which Rounds To

PUBLISHED

Question	Answer	Marks	Guidance
1	Use $\sec \theta = \frac{1}{\cos \theta}$ and $\operatorname{cosec} \theta = \frac{1}{\sin \theta}$ or other appropriate identities	B1	Must be using $\sec^2 \theta = 1 + \tan^2 \theta$ and $\operatorname{cosec}^2 \theta = 1 + \cot^2 \theta$.
	Obtain $\tan \theta = k$ using correct identities	M1	OE For any non-zero constant k , if using other identities, must come from a 3-term quadratic equation.
	Obtain $\tan \theta = 5$ and hence 78.7°	A1	AWRT
	Obtain 258.7° and no other solutions in the range	A1	AWRT
		4	

Question	Answer	Marks	Guidance	
2	Solve $4x - 1 = x + 3$ to obtain $x = \frac{4}{3}$	B1		
	Attempt solution of linear equation where signs of $4x$ and x are different	M1		
	Obtain final value $x = -\frac{2}{5}$	A1		
	Substitute numerical values and apply modulus signs correctly to obtain $ \frac{-12}{5} - \frac{1}{3} $ or equivalent, retaining exactness and with no subsequent squaring	M1	Allow <i>their</i> p and q , $p < q$.	
	Obtain $\frac{31}{15}$	A1	or exact equivalent.	
	Alternative method for Question 2			
	State or imply non-modulus equation $(4x - 1)^2 = (x + 3)^2$	B1		
	Attempt solution of 3-term quadratic equation	M1		
Obtain final values $-\frac{2}{5}$ and $\frac{4}{3}$	A1			

PUBLISHED

Question	Answer	Marks	Guidance
2	Substitute numerical values and apply modulus signs correctly to obtain $ \frac{-12}{5} - \frac{1}{3} $ or equivalent, retaining exactness and with no subsequent squaring	M1	Allow <i>their</i> $p < q$.
	Obtain $\frac{31}{15}$	A1	or exact equivalent.
		5	

Question	Answer	Marks	Guidance
3	State or imply equation is $\ln y = \ln A + k \ln x$	B1	
	Equate k to gradient of line	M1	or eliminate $\ln A$ from simultaneous equations using appropriate values.
	Obtain $k = 2.4$	A1	
	Substitute appropriate values to find $\ln A$	M1	
	Obtain $\ln A = 1.526$ and hence $A = 4.6$	A1	AWRT
	Alternative method for Question 3		
	State or imply $e^{3.47} = Ae^{0.81k}$ and $e^{2.87} = Ae^{0.56k}$	B1	
	Eliminate A	M1	
	Obtain $k = 2.4$	A1	
	Substitute appropriate values to find A	M1	
	Obtain $A = e^{1.526}$ and hence $A = 4.6$	A1	
		5	

PUBLISHED

Question	Answer	Marks	Guidance
4(a)	Substitute $x = -\frac{1}{2}$, equate to zero and attempt solution	M1	
	Obtain $a = 6$	A1	
	Divide by $2x + 1$ at least as far as the x term	M1	or use of identity or by inspection.
	Obtain $3x^2 + 10x - 8$	A1	
	Conclude $(2x + 1)(3x - 2)(x + 4)$	A1	
		5	
4(b)	Apply logarithms and use power law for $e^{4y} = k$ where $k > 0$	M1	Using <i>their</i> answer to (a).
	Obtain $y = -0.101$ and no other answers	A1	or greater accuracy.
		2	

PUBLISHED

Question	Answer	Marks	Guidance
5(a)	Attempt use of product rule to differentiate $x \ln(4x+1)$	*M1	
	Obtain $\ln(4x+1) + \frac{4x}{4x+1} - 3$	A1	
	Equate first derivative to zero and attempt correct rearrangement to obtain the form $4x+1 = \frac{ax+b}{\ln(4x+1)}$	DM1	OE
	Confirm $x = \frac{2x+0.75}{\ln(4x+1)} - 0.25$	A1	AG – necessary detail needed. Allow verification.
		4	
5(b)	Consider sign of $x - \frac{2x+0.75}{\ln(4x+1)} + 0.25$ or equivalent for 1.8 and 1.9	M1	
	Obtain $-0.017...$ and $0.035...$ or equivalents and justify conclusion	A1	AG – necessary detail needed.
		2	
5(c)	Use iteration process correctly at least once	M1	
	Obtain final answer 1.83	A1	answer required to exactly 3 s.f.
	Show sufficient iterations to 5 sf to justify answer or show sign change in the interval [1.825, 1.835]	A1	
		3	

PUBLISHED

Question	Answer	Marks	Guidance
6	Integrate $\frac{6}{3x+2}$ to obtain form $k_1 \ln(3x+2)$	*M1	for any constant k_1 .
	Obtain correct $2\ln(3x+2)$	A1	
	Apply limits correctly	DM1	
	Obtain $2\ln 14 - 2\ln 2$ and hence $\ln 49$	A1	at this stage or later.
	Integrate $3e^{-x} - 3$ to obtain form $k_2e^{-x} + k_3x$	M1	for any non-zero constants k_2, k_3 .
	Obtain correct $-3e^{-x} - 3x$	A1	
	Apply limits to obtain $-3e^{-4} - 12 + 3$	A1	OE; implied if $\int (y_1 - y_2) dx$ approach used.
	Use correct procedure to find exact total area	M1	
	Obtain $\ln 49 + 9 + 3e^{-4}$	A1	
		9	

Question	Answer	Marks	Guidance
7(a)	State $\frac{dx}{d\theta} = -6\sin 2\theta$ and $\frac{dy}{d\theta} = 4\cos \theta$	B1	
	Use $\frac{dy}{dx} = \frac{dy}{d\theta} / \frac{dx}{d\theta}$ and equate to 2	M1	
	Use $\sin 2\theta = 2\sin \theta \cos \theta$ and attempt value of $\sin \theta$	M1	
	Obtain $\sin \theta = -\frac{1}{6}$	A1	

Question	Answer	Marks	Guidance
7(a)	Obtain $\theta = 3.31$ only	A1	AWRT; and no second answer.
	Alternative method for Question 7(a)		
	Using $x = 3(2\cos^2\theta - 1)$, $x = 3(1 - 2\sin^2\theta)$ or $x = 3(\cos^2\theta - \sin^2\theta)$ to obtain $\frac{dx}{d\theta} = a \sin\theta \cos\theta$	M1	
	Obtain $\left(\frac{dy}{dx} = \right) \frac{4\cos\theta}{-12\sin\theta\cos\theta} = 2$	A1	
	Attempt value of $\sin\theta$	M1	
	Obtain $\sin\theta = -\frac{1}{6}$	A1	
	Obtain $\theta = 3.31$ only in the given range	A1	
		5	
7(b)	State or imply $9\cos 2\theta + 4\sin\theta = 0$ and use identity to obtain quadratic in $\sin\theta$	M1	
	Obtain $18\sin^2\theta - 4\sin\theta - 9 = 0$	A1	OE
	Attempt solution to find negative value of $\sin\theta$	DM1	
	Obtain $\sin\theta = -0.604\dots$	A1	Or $\frac{2 - \sqrt{166}}{18}$, $\theta = 3.79\dots$
	Substitute value of $\sin\theta$ (or <i>their</i> θ between π and $\frac{3}{2}\pi$) in expression for first derivative	M1	
	Obtain 0.551	A1	AWRT

PUBLISHED

Question	Answer	Marks	Guidance
7(b)	Alternative method for question 7(b)		
	Cartesian equation of curve $1 - \frac{y^2}{8} = \frac{x}{3}$ oe	M1	Must be a complete method, allow unsimplified.
	Intersection of line and curve $27x^2 + 8x - 24 = 0$ oe	M1	
	$x = 0.8062\dots$	A1	
	$\theta = 3.791\dots$	A1	
	Substitute value of θ (or <i>their</i> θ between π and $\frac{3}{2}\pi$) in expression for first derivative	M1	
	Obtain 0.551	A1	AWRT
		6	