

Mark Scheme (Results)



January 2009

GCE


GCE Mathematics (6677/01)




**January 2009
6677 Mechanics M1
Mark Scheme**

Question Number	Scheme	Marks
1	$-6\mathbf{i} + \mathbf{j} = \mathbf{u} + 3(2\mathbf{i} - 5\mathbf{j})$ $\Rightarrow \mathbf{u} = -12\mathbf{i} + 16\mathbf{j}$ $\Rightarrow u = \sqrt{(-12)^2 + 16^2} = 20$	M1 A1 A1 cso M1 A1 [5]
2	(a)  or  shape values (b) $19.6 = \frac{1}{2} \times 2 \times u$ $u = 19.6$	B1 B1 (2) M1 A1 A1 (3) [5]
3	(a) $2u \rightarrow \leftarrow 4u$ $km \quad m$ $u \leftarrow \rightarrow v$ $km2u - 4mu = -kmu + mv$ $u(3k - 4) = v$ (b) $k > 2 \Rightarrow v > 0 \Rightarrow \text{dir}^n \text{ of motion reversed}$ (c) For B, $m(u(3k - 4) - -4u)$ $= 7mu$	M1 A1 A1 (3) M1A1A1 cso (3) M1 A1 f.t. A1 (3) [9]

Question Number	Scheme	Marks
4 (a) $C + D = 120g$ $M(Q), 80g \cdot 0.8 - 40g \cdot 0.4 = D \cdot 1.6$ solving $C = 90g; D = 30g$ (b) $2F + F = 40g + 20g + 60g$ $M(Q), 60gx + 20g \cdot 0.8 = 40g \cdot 0.4 + F \cdot 1.6$ solving $QX = x = \frac{16}{15} \text{ m} = 1.07\text{m}$		M1 A1 M1 A1 M1 A1 A1 (7) M1 A1 M1 A1 M1 A1 (6) [13]

Question Number	Scheme	Marks
5 (a)		B2 -1 e.e.o.o. (labels not needed) (2)
(b)	$F = \frac{1}{2} R$ $(\uparrow), R \cos \alpha + F \sin \alpha = mg$ $R = \frac{1.1g}{(\cos \alpha + \frac{1}{2} \sin \alpha)} = 9.8 \text{ N}$ $(\rightarrow), P + \frac{1}{2} R \cos \alpha = R \sin \alpha$ $P = R(\sin \alpha - \frac{1}{2} \cos \alpha)$ $= 1.96$	B1 M1 A2 M1 A1 (6) M1 A2 M1 A1 (5) [13]

Question Number	Scheme	Marks
6 (a) (b) (c)	$\tan \theta = \frac{2}{1} \Rightarrow \theta = 63.4^\circ$ angle is 153.4° $(4 + p)\mathbf{i} + (q - 5)\mathbf{j}$ $(q - 5) = -2(4 + p)$ $2p + q + 3 = 0 *$ $q = 1 \Rightarrow p = -2$ $\Rightarrow \mathbf{R} = 2\mathbf{i} - 4\mathbf{j}$ $\Rightarrow \mathbf{R} = \sqrt{2^2 + (-4)^2} = \sqrt{20}$ $\sqrt{20} = m8\sqrt{5}$ $\Rightarrow m = \frac{1}{4}$	M1 A1 A1 (3) B1 M1 A1 A1 (4) B1 M1 M1 A1 f.t. M1 A1 f.t. A1 cao (7) [14]

Question Number	Scheme	Marks
7	(a) $T - 5g \sin \alpha = 5a$ $15g - T = 15a$ solving for a $a = 0.6g$ solving for T $T = 6g$	M1 A1 M1 A1 M1 A1 M1 A1 (8)
	(b) For Q : $\begin{array}{l} 5g - N = 5a \\ N = 2g \end{array}$	M1 A1 A1 f.t. (3)
	(c)  $F = 2T \cos\left(\frac{90^\circ - \alpha}{2}\right)$ $= 12g \cos 26.56^\circ$ $= 105 \text{ N}$	M1 A2 A1 f.t. A1 (5) [16]