

Cambridge
International
A Level

Cambridge International Examinations
Cambridge International Advanced Level

CANDIDATE
NAME

CENTRE
NUMBER

--	--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--

BIOLOGY

9700/42

Paper 4 A2 Structured Questions

October/November 2015

2 hours

Candidates answer on the Question Paper.

Additional Materials: Answer paper available on request.

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name in the spaces provided at the top of this page.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

DO **NOT** WRITE IN ANY BARCODES.

Section A

Answer **all** questions.

Section B

Answer **one** question.

Circle the number of the Section B question you have answered in the grid below.

You may lose marks if you do not show your working or if you do not use appropriate units.

Electronic calculators may be used.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.

For Examiner's Use	
Section A	
1	
2	
3	
4	
5	
6	
7	
8	
Section B	
9 or 10	
Total	

This document consists of **21** printed pages, **1** blank page and **2** lined pages.

Section A

Answer **all** the questions.

1 (a) The molecules listed below are all associated with respiration.

- | | | | |
|---------------------|-----------------|----------------|---------------|
| ATP synthase | glucose | ATP | NAD |
| oxaloacetate | pyruvate | citrate | oxygen |

From these molecules identify:

a phosphorylated nucleotide

a 3-carbon compound

a coenzyme

an enzyme [4]

(b) A sample of tree sap, rich in sugars, was found to be contaminated with yeast. This sample was tested for the concentration of ethanol at regular intervals.

The results are shown in Fig. 1.1.




Fig. 1.1

- (i) Calculate the percentage increase in ethanol concentration between 15 and 45 hours.
Show your working.

answer % [2]

- (ii) Suggest why the concentration of ethanol decreased after 45 hours.

.....
.....
.....[1]

[Total: 7]

BLANK PAGE

- 2 Gold ions (Au^{3+}) are toxic to most microorganisms. However, the bacterium *Delftia acidovorans* is frequently found in sticky layers, called biofilms, that form on the surface of gold deposits.

D. acidovorans produces a peptide synthase that catalyses the synthesis of a small peptide called delftibactin. When isolated, delftibactin can precipitate Au^{3+} ions as small particles of metallic gold. Delftibactin is a secondary metabolite.

- (a) Name another example of a *secondary metabolite* and explain what is meant by the term.

example

explanation

.....
.....
.....[3]

- (b) A mutant strain of *D. acidovorans* has been identified in which the gene coding for peptide synthase is inactive.

The wild-type (normal) and mutant *D. acidovorans* were grown on agar plates and then flooded with gold chloride solution, which contains Au^{3+} ions. The appearance of such a plate after this treatment is shown in Fig. 2.1.




Fig. 2.1

With reference to Fig. 2.1, suggest how delftibactin protects *D. acidovorans* from toxic Au^{3+} ions.

.....
.....
.....
.....
.....
.....
.....[3]

The endosperm contains starch stores. There are also small quantities of sucrose stored in the aleurone layer.

Water uptake stimulates the production of a plant growth regulator in the seed, which in turn activates the synthesis of enzymes in the aleurone layer. These enzymes hydrolyse starch to maltose and glucose.

Name the plant growth regulator involved in the activation of the synthesis of the enzymes.

.....[1]

(c) An investigation was carried out into the role of a gene, *TaSUT1*, which codes for a sucrose transporter protein, in the germination of wheat seeds.

- Wheat seeds were germinated and left to grow for 3, 7 or 10 days.
- Samples of tissues from the roots, seeds and shoots of the seedlings were tested for the presence of mRNA transcribed from *TaSUT1*.
- The extracted mRNA was mixed with a probe, and then placed on agarose gel across which a voltage was applied.

The results are shown in Fig. 4.2.




Fig. 4.2

(i) Suggest why the researchers looked for mRNA transcribed from the *TaSUT1* gene, rather than for the gene itself.

.....
.....
.....
.....[2]

(ii) Explain what the results in Fig. 4.2 indicate about the sequence of activity of *TaSUT1*, from day 3 to day 10, in the root, seed and shoot of a seedling.

.....
.....
.....
.....[2]

(d) *TaSUT1* codes for the sucrose transporter protein, SUT. This protein transports only sucrose.

To investigate where this protein was present in a germinating wheat seedling, a fluorescent antibody for SUT was added to sections of tissues from the seedling.

(i) Suggest how this enabled the researchers to determine the areas where SUT was located.

.....
.....
.....
.....
.....[2]

(ii) Immediately after germination began, SUT was found in the membranes of cells in the aleurone layer. It was also determined that the most common sugar in the endosperm in the first hours after germination was sucrose.

Explain how these results support the hypothesis that the **first** source of sugar for the embryo during germination is sucrose from the aleurone layer and **not** sugars produced by the hydrolysis of starch.

.....
.....
.....
.....
.....[2]

5 (a) Explain the meaning of the term biodiversity.

.....
.....
.....
.....
.....[2]

(b) The Javan gibbon, *Hylobates moloch*, is an endangered species.

Fig. 5.1 shows a female Javan gibbon with an infant.



Fig. 5.1

Javan gibbons live in fragmented patches of undisturbed forest in western Java, Indonesia. Habitat loss has reduced the population of wild gibbons to around 4500 individuals.

(i) Suggest why the separation of their habitat into small fragments, rather than a single large area, poses a threat to the long-term survival of this species.

.....
.....
.....
.....
.....
.....
.....
.....[3]

- 6 Scorpions are predatory arthropods. They have a pair of grasping claws at the front of their bodies and a tail with a stinger. The stinger is used to inject venom into their prey to cause paralysis.

Fig. 6.1 shows a scorpion.



Fig. 6.1

(a) Scorpion venom contains two active components:

- a toxin that affects ion channels at synapses of the nervous system of their prey
- an inhibitor of an enzyme found at these synapses.

For **each** component of the venom, suggest and explain **one** way in which it may stop the correct functioning of the synapse.

toxin

.....

.....

.....

.....

inhibitor

.....

.....

.....

.....

.....

[4]

(b) Scorpions stand very still on the sand. Moving prey will disturb grains of sand, and scorpions are able to detect this movement using sensory organs, known as slit hairs, at the tips of their legs. Some of the cells of a slit hair act as sensory receptors.

(i) State the role of a sensory receptor.

.....
.....
.....[1]

(ii) When a slit hair is bent by the movement of the sand the potential difference across the cell surface membranes of the slit hair cells becomes more positive inside compared to the outside.

State the name given to the initial change in potential difference that may lead to an action potential.

.....[1]

(iii) Action potentials may then be sent by the cells in the slit hairs to the central nervous system (CNS) of the scorpion.

Explain how the scorpion is able to distinguish between a small and a large movement of sand.

.....
.....
.....
.....
.....
.....
.....
.....[2]

[Total: 8]

- 7 (a) One way to estimate the rate of photosynthesis is to measure the rate of uptake of carbon dioxide.

Fig. 7.1 shows the relationship between light intensity and relative carbon dioxide uptake and production in a dicotyledonous plant.




Fig. 7.1

- (i) State **one physical factor** that may limit the rate of photosynthesis at **Y**.
[1]
- (ii) State **two** features of a dicotyledonous leaf that can affect the rate of photosynthesis.

[2]
- (iii) Explain the shape of the curve as light intensity increases from **0** to **X**.

[2]

(b) The uptake of radioactively-labelled carbon dioxide in chloroplasts was investigated. Three tubes, each containing different components of chloroplasts, were exposed to light. The results of the investigation are shown in Table 7.1.

Table 7.1

tube	contents	uptake of radioactively-labelled carbon dioxide / counts per minute
A	stroma and grana	96 000
B	stroma, ATP and reduced NADP	97 000
C	stroma	4 000

- (i) Name the substance that combines with carbon dioxide in a chloroplast.
[1]
- (ii) Explain why the results in tube **B** are similar to those in tube **A**.

[2]
- (iii) Explain why the uptake in tube **C** was less than the uptake in tube **B**.

[2]

- (c) Complete the following paragraph by using the most suitable words to fill in the gaps.

In a photosystem, several hundred accessory pigment molecules surround a primary pigment molecule, called, in the membrane. The position of the primary pigment is also called the Light energy is absorbed by the accessory pigments and passed on to the primary pigment. Electrons are excited to a higher energy level. They are emitted from the primary pigment and are captured by electron acceptors and eventually pass along the, producing ATP. [4]

[Total: 14]

