

Mark Scheme (Results)

Summer 2015

Pearson Edexcel International GCSE
Mathematics A (4MA0)
Paper 4H

Pearson Edexcel Level1/Level 2 Certificate
Mathematics A (KMA0)
Paper 4H

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

Summer 2015

Publications Code UG042089

All the material in this publication is copyright

© Pearson Education Ltd 2015

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.
- **Types of mark**
 - M marks: method marks
 - A marks: accuracy marks
 - B marks: unconditional accuracy marks (independent of M marks)
- **Abbreviations**
 - cao – correct answer only
 - ft – follow through
 - isw – ignore subsequent working
 - SC - special case
 - oe – or equivalent (and appropriate)
 - dep – dependent
 - indep – independent
 - eeo – each error or omission
 - awrt – answer which rounds to

- **No working**

If no working is shown then correct answers normally score full marks

If no working is shown then incorrect (even though nearly correct) answers score no marks.

- **With working**

If there is a wrong answer indicated on the answer line always check the working in the body of the script (and on any diagrams), and award any marks appropriate from the mark scheme.

If it is clear from the working that the "correct" answer has been obtained from incorrect working, award 0 marks.

Any case of suspected misread loses A (and B) marks on that part, but can gain the M marks.

If working is crossed out and still legible, then it should be given any appropriate marks, as long as it has not been replaced by alternative work.

If there is a choice of methods shown, then no marks should be awarded, unless the answer on the answer line makes clear the method that has been used.

If there is no answer on the answer line then check the working for an obvious answer.

- **Ignoring subsequent work**

It is appropriate to ignore subsequent work when the additional work does not change the answer in a way that is inappropriate for the question: eg. Incorrect cancelling of a fraction that would otherwise be correct.

It is not appropriate to ignore subsequent work when the additional work essentially makes the answer incorrect eg algebra.

Transcription errors occur when candidates present a correct answer in working, and write it incorrectly on the answer line; mark the correct answer.

- **Parts of questions**

Unless allowed by the mark scheme, the marks allocated to one part of the question CANNOT be awarded in another.

Apart from Questions 7c, 13, 19a, 20b, 21b and 23, where the mark scheme states otherwise, the correct answer, unless clearly obtained by an incorrect method, should be taken to imply a correct method.

Question	Working	Answer	Marks	Comments
1	$1 - \frac{2}{5}$ or $\frac{3}{5}$ oe		3	M1 or for $\frac{2}{5} \times 20$ oe or 8 or $\frac{8}{20}$
	$\frac{3}{5} \times 20$ oe	12		M1 (dep) for 20 - '8' A1 NB: $\frac{12}{20}$ gains M2, A0
				Total 3 marks

2	$15 \div (6 - 4) (=7.5)$ $"7.5" \times 3 (=22.5)$	22.5(0)	3	M1 or use of cancelled ratios (eg $3 : 6 : 4 = 0.75 : 1.5 : 1$, M1dep $15 \div 0.5 (=30)$ or cancelled ratios, (eg 30×0.75) or M2 for $15 \div \frac{2}{3}$ oe A1 oe
	Alternative			
	$\frac{6n}{13} - \frac{4n}{13} = 15$ or $\frac{2n}{13} = 15$ oe or $15 \times \frac{13}{2}$ oe or $n = 97.5$		3	M1
	$\frac{3}{3+6+4} \times 97.5$ oe			M1dep
		22.5(0)		A1 oe
	Alternative			
	$(3 : 6 : 4) \dots 21 : 42 : 28$ and $24 : 48 : 32$ $22.5 : 45 : 30$	22.5(0)	3	M1 For using ratios and seeing $21 : 42 : 28$ and $24 : 48 : 32$ M1dep Correct line or $(21+24) \div 2$ A1 oe
				Total 3 marks

3	(a)(i)		90	1	B1
	(a)(ii)		25	1	B1
	(b)		line from (13 45, 45) to (15 15, 45) and line from (15 15, 45) to (16 30, 0)	2	B2 B1 for line from ((13 45, 45) to (15 15, 45) or for a line from (x, 45) to (16 30, 0) where x is a time before 1630
					Total 4 marks

4	(a)	$(25 + 1) \div 2$ or 13 or 12.5		2	M1 or listing scores and clear attempt to find middle value A1
	(b)	$1 \times 9 + 2 \times 6 + 3 \times 3 + 4 \times 2 + 5 \times 1 + 4 \times 6$ oe (=67) "67" $\div 25$ or $\frac{9+12+9+8+5+24}{25}$ oe (allow one error in a product)		3	M1 sight of at least 4 products and intention to add M1dep for division of sum of products by 25 (can be their 25 if evidence of adding frequencies) A1 accept 2.7 or 3 if preceded by $\frac{67}{25}$
					Total 5 marks

5	(a) $y = -x$ drawn		2	M1 or a congruent shape with the correct orientation in the 1 st quadrant or a correct reflection in $y = x$ A1
	(b)	Rotation about $(0, -1)$ 90° clockwise	3	B1 Rotation B1 (centre) $(0, -1)$ B1 90° clockwise or -90° or 270° anti-clockwise or $+270^\circ$ NB. If more than one transformation given then no marks should be awarded
				Total 5 marks


6 (a)	angle MRQ (or RMQ) = x or $\frac{180-y}{2}$	$180 - 2x$	2	M1 could be marked on diagram or for a correct equation in x and y A1 oe eg $2(90 - x)$, $2(180 - x) - 180$ etc
(b)	(6 - 2) × 180 oe (=720) "720" - (90 + 115 + 144 + 87), [720 - 436] or 284 "284" ÷ 2	142	4	M1 or (180 - 360 ÷ 6) × 6 M1dep M1dep A1
	Alternative 180 - 90(=90), 180 - 115(=65), 180 - 144(=36), 180 - 87(=93) 360 - ("90" + "65" + "36" + "93"), [360 - 284] (= 76) 180 - ("76" ÷ 2)	142	4	M1 A correct method to find each of the exterior angles at A,C,D & E - angles could be seen on diagram. M1dep A correct method to find the total of the remaining exterior angles M1dep A correct method to find k A1
				Total 6 marks

7	(a)		$24 - 18y$	1	B1
	(b)		$e(e + 4)$	1	B1 Allow $(e + 0)(e + 4)$ ignore missing bracket at end
	(c)	Eg. $7x - 2x = -3 - 8$		3	M1 for correct rearrangement with x terms on one side and numbers on the other or the correct simplification of either x terms or numbers on one side in a correct equation eg. $5x + 8 = -3$; $7x = 2x - 11$
		$5x = -11$			M1 Award also for $-5x = 11$
			-2.2 oe		A1 -2.2 oe dependent on at least M1 awarded; if no correct algebraic working then award no marks
	(d)	$y^2 + 10y - 2y - 20$		2	M1 for 3 correct terms out of 4 or for 4 correct terms ignoring signs or for $y^2 + 8y + c$ for any non-zero value of c or for $\dots + 8y - 20$
			$y^2 + 8y - 20$		A1 cao
	(e)		$4e^2f(5e^3f - 4)$	2	B2 B1 for a correct but incomplete factorised answer with a minimum of 2 out of 4, e^2 or f outside the bracket, ie $4e^2(5e^3f^2 - 4f)$, $4f(5e^5f - 4e^2)$, $e^2f(20e^3f - 16)$, $4ef(5e^4f - 4e)$, $2e^2f(10e^3f - 8)$ or $4e^2f$ (a two term algebraic expression)
					Total 9 marks

8	$\cos 39 = \frac{11.3}{x}$ oe		3	M1	or $\frac{x}{\sin 90} = \frac{11.3}{\sin(180-90-39)}$
	$(x =) \frac{11.3}{\cos 39}$			M1	or $x = \frac{11.3}{\sin(180-90-39)} \times \sin 90$
		14.54		A1	awrt 14.54
	Alternative				
	$\tan 39 = \frac{y}{11.3}$; $y = 9.15\dots$ "9.15" ² + 11.3 ² = x ² oe			M1	Must get to correct Pythagoras statement
	$(x =) \sqrt{9.15^2 + 11.3^2}$ oe			M1	
		14.54		A1	awrt 14.54 (NB: 14.5 with no working gains M0A0)
					Total 3 marks

9	(a)	$-5 - 4 < x \leq 3 - 4$		2	M1	subtraction of 4 from either side in an inequality or one side of inequality correct (eg $x \leq -1$) or for $-5-4(=-9)$ and $3-4(=-1)$
			$-9 < x \leq -1$		A1	Accept $x > -9$, $x \leq -1$
	(b)		$-3, -2, -1, 0, 1$	2	B2	B1 for one omission or addition
						Total 4 marks

10 (a)	$131 - 111$	20	2	M1 For 60 & 20 or 60.75 & 20.25 seen in working or in diagram. A1 Allow answers in the range 20-21 from correct readings.
(b)	$80 - 50 (=30)$ $\frac{"30"}{80} \times 100$	37.5	3	M1 or for $\frac{50}{80} \times 100 (=62.5)$ M1 or for $100 - "62.5"$ A1 Accept 38% from correct working
				Total 5 marks

11	<p>(12 =) $2 \times 2 \times 3$ or (120 =) $2 \times 2 \times 2 \times 3 \times 5$ (condone 2,2,3 or 2,2,2,3,5) [factors could be seen at the end of a 'factor tree' or in a 'factor ladder'] or Venn diagram with the middle and one other region correct:</p>  <p>Where 10 may be 2,5 and 4 may be 2,2</p>	40	2	M1 or for a <u>list</u> of at least 5 consecutive multiples of 4 or a <u>list</u> of at least 5 factors of 120 or for $12x = 120 \times 4$ oe (eg $\frac{120}{12} \times 4 (= x)$) or $12 \div 4 (= 3)$ and $120 \div "3"$ A1 accept $2 \times 2 \times 2 \times 5$ or $2^3 \times 5$
				Total 2 marks

12	eg $\frac{16}{100} \times 65000$ oe or 10400	65000 $\times 0.84^3$		3	M1 For $\frac{16}{100} \times 65000$ oe or 10400	(M2 for 65000×0.84^3) or (M1 for 65000×0.84 or 54600 or 65000×0.84^2 or 45864 or 65000×0.84^4 or 32361.63..)
	$\frac{16}{100} \times (65000 - "10400")$ $= 8736$ $\frac{16}{100} \times (65000 - "10400" - "8736")$ $= 7338.24$ $65000 - "10400" - "8736" - "7338.24"$				M1 For completing Method	
					Accept (1 – 0.16) as equivalent to 0.84 throughout	
					SC: If no other marks gained, award M1 for 65000×0.48 oe (=31200) or 65000×0.52 oe (=33800)	
			38525.76		A1 for 38525 – 38526	
					Total 3 marks	


<p>13</p>	$\frac{-2 \pm \sqrt{2^2 - 4 \times 3 \times -7}}{2 \times 3}$ $\frac{-2 \pm \sqrt{4 + 84}}{6} = \frac{-2 \pm \sqrt{88}}{6}$	<p>1.23, -1.90</p>	<p>3</p>	<p>M1 correct substitution into the quadratic formula, allow one sign error in numbers; some evaluation may be seen. M1 Indep for simplification of discriminant to $\sqrt{88}$ or $\sqrt{4 + 84}$ or $2\sqrt{22}$ A1 1.23 (or better), -1.90 (accept answers in range -1.90 to -1.89) provided at least M1 awarded</p>
Total 3 marks				

<p>14 (a)</p>	$\frac{20}{8} \times 3 \text{ oe}$	<p>7.5</p>	<p>2</p>	<p>M1 A1 oe</p>
<p>(b)</p>	$1875 \div \left(\frac{20}{8}\right)^3 \text{ oe}$	<p>120</p>	<p>2</p>	<p>M1 for $\left(\frac{20}{8}\right)^3$ or $\left(\frac{8}{20}\right)^3$ oe, accept ratios A1</p>
Alternative				
	$\frac{1875}{20} \times \left(\frac{8}{20}\right)^2 (=15) \text{ oe}$	<p>120</p>		<p>M1 A1</p>
Total 4 marks				

<p>15 (a)</p>	<p>Probabilities on branches correct.</p>	$\frac{6}{10}, \frac{4}{10}, \frac{5}{9}, \frac{4}{9}, \frac{6}{9}, \frac{3}{9}$	<p>3</p>	<p>B1 for $\frac{6}{10}, \frac{4}{10}$ oe on LH branches</p> <p>B1 for $\frac{5}{9}, \frac{4}{9}$ oe on top RH branches</p> <p>B1 for $\frac{6}{9}, \frac{3}{9}$ oe on bottom RH branches</p> <p>Decimals given on the 2nd set of branches to be to at least 2dp (truncated or rounded).</p>
<p>(b)</p>	<p>$\frac{6}{10} \times \frac{4}{9}$ or $\frac{4}{10} \times \frac{6}{9}$ or ft from their tree diagram</p> <p>$\frac{6}{10} \times \frac{4}{9} + \frac{4}{10} \times \frac{6}{9}$ or ft from their tree diagram</p>	$\frac{48}{90}$ oe	<p>3</p>	<p>M1 or $\frac{6}{10} \times \frac{5}{9} + \frac{4}{10} \times \frac{3}{9} \left(= \frac{42}{90} \right)$</p> <p>M1dep or $1 - \frac{42}{90}$</p> <p>A1 Allow 0.53(33...)</p> <p>Note: If all 4 probability products are seen at the ends of the branches on the tree diagram or in lists in the working space for (b), marks can only be awarded in (b) if it is clear which product(s) they are intending to use.</p>
<p>Total 6 marks</p>				

16	$\frac{x}{\sin 62} = \frac{14.6}{\sin 105}$		3	M1 or $\frac{\sin 62}{x} = \frac{\sin 105}{14.6}$
	$\frac{14.6}{\sin 105} \times \sin 62$			M1
		13.3		A1 Accept answers in the range 13.3 – 13.35
				Total 3 marks

17	$\begin{pmatrix} 5 \\ -1 \end{pmatrix} + \begin{pmatrix} 2 \\ -3 \end{pmatrix}$ or $\begin{pmatrix} 5 \\ -1 \end{pmatrix} - \begin{pmatrix} -2 \\ 3 \end{pmatrix}$		2	M1 Or for $\begin{pmatrix} 7 \\ a \end{pmatrix}$ or $\begin{pmatrix} b \\ -4 \end{pmatrix}$
		$\begin{pmatrix} 7 \\ -4 \end{pmatrix}$		A1 Must be written as a vector.
				Total 2 marks

<p>18 (a)</p>		<p>5, 10, 8, 13</p>	<p>3</p>	<p>B1 for 8 in intersection and 13 in correct position B1 for 5 in correct position B1 for 10 in correct position</p>
<p>(b)</p>		<p>31</p>	<p>1</p>	<p>B1 or ft from diagram</p>
<p>(c)</p>		<p>10</p>	<p>1</p>	<p>B1 or ft from diagram</p>
				<p>Total 5 marks</p>

19	(a)	$5 \times 7 + 5\sqrt{2} - 7\sqrt{8} - \sqrt{8}\sqrt{2}$ Or $35 + 5\sqrt{2} - 7\sqrt{8} - \sqrt{8}\sqrt{2}$ Or $35 + 5\sqrt{2} - 7\sqrt{8} - \sqrt{16}$ Or $35 + 5\sqrt{2} - 7(2\sqrt{2}) - \sqrt{16}$		3	M1	4 terms, allow one sign error Note 5×7 may be 35 $-7\sqrt{8}$ may be $-7 \times 2\sqrt{2}$ but not $-14\sqrt{2}$ $\sqrt{8}\sqrt{2}$ may be $\sqrt{16}$ or 4 or for $\sqrt{8} = 2\sqrt{2}$
		$35 + 5\sqrt{2} - 14\sqrt{2} - 4$ or $31 + 5\sqrt{2} - 14\sqrt{2}$			M1dep	
					A1	show from correct working
	(b)	$\frac{3c - \sqrt{c}}{\sqrt{c}} \times \frac{\sqrt{c}}{\sqrt{c}}$ or $\frac{3c\sqrt{c} - c}{c}$ or $\frac{\sqrt{c}(3\sqrt{c} - 1)}{\sqrt{c}}$		2	M1	
		A1				
					Total 5 marks	

20	(a)	Eg. $2n$ is always even so $2n + 1$ is odd	explanation	1	B1
	(b)	$(2n + 1) + (2n + 3) + (2n + 5) + (2n + 7)$ oe $8n + 16$ $8(n + 2)$	show	3	M1 M1dep A1 or a complete explanation from correct algebraic working
		Alternative			
		let x be an even number $x + 1 + x + 3 + x + 5 + x + 7$ oe $4x + 16$ $4(x + 4)$		3	M1 For defining x (at beginning or end) and summing 4 consecutive odd numbers M1dep A1 For a complete explanation from correct algebraic working., eg $x + 4$ must be even and 4 times an even number = $4 \times 2n$ which is a multiple of 8
		Alternative			
		Let y be an odd number $y + y + 2 + y + 4 + y + 6$ oe $4y + 12$ $4(y + 3)$		3	M1 For defining y (at beginning or end) and summing 4 consecutive odd numbers M1dep A1 For a complete explanation from correct algebraic working. eg $(y + 3)$ is (odd + odd) which is even and therefore a multiple of 2, and $4 \times 2n$ is a multiple of 8
					Total 4 marks

21	(a)		$3x^2 + 12x$	2	B2 B1 for $3x^2$ or $12x$
	(b)	" $3x^2 + 12x = 0$ $3x(x + 4) = 0$ or $x(x + 4) = 0$ or $x(3x + 12) = 0$ or correct use of the formula (all values correctly substituted) or completing the square (as far as $(3)((x+2)^2-4)=0$) $x = 0$ and $x = -4$		4	M1 ft M1 ft as long as in the form of a quadratic equation. A1 dep on M1 A1 Could be $x=0,y=5;x=-4,y=37$
			(0, 5) and (-4, 37)		
					Total 6 marks

22		$\frac{1}{2}ab\sin 150$ oe or $\frac{1}{2}(a+1)(b+2)$ oe (must be $\sin 150$, not $\sin C$)		5	M1 Or $\frac{1}{2}(a+1)(b+2)\sin 90$
		$3 \times \frac{1}{2}ab\sin 150 = \frac{1}{2}(a+1)(b+2)$ oe			M1 correct equation, eg may see $\frac{3}{4}ab = \dots$
		$3 \times \frac{1}{2}ab\sin 150 = \frac{1}{2}(ab + b + 2a + 2)$ oe			M1 expansion of brackets in a correct equation
		$3ab - 2ab - 4a = 2b + 4$ oe or $3ab\sin 150 - ab - 2a = b + 2$ oe			M1 isolation of terms in a in a correct equation (may be on either side of equation & can still have $\sin 150$)
			$\frac{2b+4}{b-4}$ oe		A1 eg $\frac{-2b-4}{4-b}$; $\frac{\frac{1}{2}b+1}{\frac{1}{4}b-1}$, $\frac{b+2}{3b\sin 150 - b - 2}$, etc
					Total 5 marks

23	$x^2 + (10 - 2x)^2 = 20$ oe			M1 or $\left(\frac{10-y}{2}\right)^2 + y^2 = 20$ oe
	$100 - 20x - 20x + 4x^2$ or $100 - 40x + 4x^2$			M1 (indep) for correct expansion of $(10 - 2x)^2$ or $\left(\frac{10-y}{2}\right)^2$ even if unsimplified
	$x^2 + 100 - 20x - 20x + 4x^2 = 20$ or $x^2 + 100 - 40x + 4x^2 = 20$			$\left(\frac{100-10y-10y+y^2}{4}\right) + y^2 = 20$ or $\left(\frac{100-20y+y^2}{4}\right) + y^2 = 20$
	$5x^2 - 40x + 80 (= 0)$ or $x^2 - 8x + 16 (= 0)$		5	A1 $5y^2 - 20y + 20 (= 0)$ or $y^2 - 4y + 4 (= 0)$
	$5(x-4)(x-4) (= 0)$ or $(x-4)(x-4) (= 0)$ or $(5x-20)(x-4) (= 0)$ or $\frac{-8 \pm \sqrt{(-8)^2 - 4(1)(16)}}{2(1)}$ oe (may be partially evaluated; condone lack of brackets around negative numbers)			M1 $5(y-2)(y-2) (= 0)$ or $(y-2)(y-2) (= 0)$ or $(5y-10)(y-2) (= 0)$ or $\frac{-4 \pm \sqrt{(-4)^2 - 4(1)(4)}}{2(1)}$ oe (may be partially evaluated; condone lack of brackets around negative numbers)
	$x = 4$ $y = 2$	$x = 4 ; y = 2$		A1 dep on all preceding method marks No marks for $x = 4 ; y = 2$ with no working. Accept (4, 2)
				Total 5 marks

