

Mark Scheme (Results)

January 2012

International GCSE Chemistry (4CH0)
Paper 1C
Science Double Award (4SC0) Paper
1C

## **Edexcel and BTEC Qualifications**

Edexcel and BTEC qualifications come from Pearson, the world's leading learning company. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information, please call our GCE line on 0844 576 0025, our GCSE team on 0844 576 0027, or visit our qualifications website at <a href="https://www.edexcel.com">www.edexcel.com</a>. For information about our BTEC qualifications, please call 0844 576 0026, or visit our website at <a href="https://www.btec.co.uk">www.btec.co.uk</a>.

If you have any subject specific questions about this specification that require the help of a subject specialist, you may find our Ask The Expert email service helpful.

Ask The Expert can be accessed online at the following link:

http://www.edexcel.com/Aboutus/contact-us/

Alternatively, you can speak directly to a subject specialist at Pearson about Edexcel qualifications on our dedicated Science telephone line: 0844 576 0037

## Pearson: helping people progress, everywhere

Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for raising achievement through innovation in education. Find out more about how we can help you and your students at: <a href="https://www.pearson.com/uk">www.pearson.com/uk</a>

January 2012
Publications Code UG030278
All the material in this publication is copyright
© Pearson Education Ltd 2012

## INTERNATIONAL GCSE CHEMISTRY 4CHO 4SCO /1C - JANUARY 2012

|   | Question number |    |    | Answer         | Notes   | Marks |
|---|-----------------|----|----|----------------|---|-------|
| 1 | а               |    | M1 | beaker         | Accept phonetic spellings   | 1     |
|   |                 |    | M2 | water          |   | 1     |
|   |                 |    | М3 | glass rod      |   | 1     |
|   | M4              |    | M4 | funnel         |   | 1     |
|   | M5              |    | M5 | conical flask  |   | 1     |
|   |                 |    | M6 | water          |   | 1     |
|   |                 |    |    |                |   |       |
|   | b               | i  | M1 | (filter) paper | Accept phonetic spellings   | 1     |
|   |                 |    |    |                | Ignore alternatives to filter, such as kitchen / chromatography - |       |
|   |                 |    |    |                | the essential word is paper                                       |       |
|   |                 |    |    |                |   |       |
|   |                 | ii | M1 | sand           | Accept phonetic spellings   | 1     |
|   |                 |    |    |                |   |       |
|   | С               |    | M1 | cross in box 4 |   | 1     |
|   |                 |    | M2 | cross in box 5 |   | 1     |
|   |                 |    |    |                |   |       |

Total 10 marks

|   | iesti<br>umb | _  |          | Answer   | Notes  | Marks |
|---|--------------|----|----------|--|--|-------|
| 2 | а            | İ  | M1       | (hydrated) iron(III) oxide / Fe <sub>2</sub> O <sub>3</sub>  | Allow (hydrated) iron oxide and Fe <sub>2</sub> O <sub>3</sub> .xH <sub>2</sub> O  | 1     |
|   |              | ii | M1       | oxygen / O <sub>2</sub>  | Allow air Do not accept O  | 1     |
|   |              |    | M2       | water / H <sub>2</sub> O   | Accept poorly written formulae such as H2O and O <sup>2</sup> Accept phonetic spellings Answers can be in either order Reject salt / acid - eg salt water does not score | 1     |
|   | b            |    | M1       | cross in box 4 (oxidation)   |  | 1     |
|   | С            | i  | M1       | Zinc / Zn  | Accept phonetic spellings  | 1     |
|   |              | ii | M1       | cross in box 2 (Bucket)  |  | 1     |
|   | d            |    | M1<br>M2 | oiling / greasing / painting / covering with plastic / coating with tin or named metal (aluminium or below) in reactivity series / attaching magnesium or zinc blocks (to ships) | Any two for 1 each Ignore sacrificial protection and galvanising and alloying  | 2     |

**Total 8 marks** 

|   | Questio<br>n<br>numbe<br>r |    |          | Answer   | Notes   | Marks |
|---|----------------------------|----|----------|--|---|-------|
| 3 | 3 a M1                     |    | M1       | NH <sub>4</sub> <sup>+</sup>   | Award 1 if wrong way around   | 1     |
|   |                            |    | M2       | CI <sup>-</sup>  | Penalise missing charges both times   | 1     |
|   | b                          | i  | M1       | (add) sodium hydroxide/NaOH (solution) (and warm)  test (gas / ammonia) with (damp red) litmus (paper) | Accept any identified Group 1 or Group 2 hydroxide  If no reagent added, max 1 mark for correct test AND result even if dipped into solution  If just hydroxide or OH <sup>-</sup> ions, do not award M1 but continue marking  If any other incorrect reagent added, then 0/3 Accept use of universal indicator  Accept holding litmus above tube etc | 1     |
|   |                            |    | М3       | OR test with hydrogen chloride / conc HCI  (litmus paper) turns blue OR white smoke/solid/powder       | Reject blue litmus for M2 and M3 Do not penalise ammonium instead of ammonia in M2 Do not allow (dilute) hydrochloric acid Do not award M3 if litmus dipped into solution (even if only implied)  | 1     |
|   |                            | ii | M1<br>M2 | (add) silver nitrate/AgNO <sub>3</sub> (solution) (dilute) nitric acid                                 | If missing or incorrect reagent, 0/3 Do not accept any other acid or just acidified If acid missing or wrong, M3 can still be awarded   | 1 1   |
|   |                            |    | M3       | white precipitate / solid / suspension   | If bleaching litmus paper mentioned, only M1 can be awarded   | 1     |

| n | Questio<br>n<br>numbe<br>r |    |          | Answer   | Notes   | Marks  |
|---|----------------------------|----|----------|--|---|--------|
| 3 | С                          |    | M1       | reversible / goes both ways  | Ignore equilibrium  | 1      |
|   | d                          | i  | M1       | ammonium chloride / NH <sub>4</sub> Cl   | Do not accept ammonia chloride If name and formula given, both must be correct  | 1      |
|   |                            | ii | M1       | ammonia / NH <sub>3</sub> / molecules / they / it are / move / diffuse /travel faster / quicker  | Ignore descriptions such as lighter / smaller / denser  Accept phonetic spellings including amonia / ammonium  Do not accept hydrogen chloride / hydrochloric acid / HCI / ammonium chloride / NH <sub>4</sub> CI in place of ammonia  Accept all other words with same meaning as faster - eg speedier  Do not accept react faster or travel further  Accept reverse statements such as hydrogen chloride slower | 1      |
|   | е                          |    | M1<br>M2 | Corrosive / burns / damages skin or eyes<br>Wear eye protection eg goggles or mask<br>/ gloves / place bung in the end of the<br>tube / use of fume cupboard | Ignore harmful / irritant / toxic / poisonous Allow tongs / tweezers if reference to cotton wool Ignore lab coats M1 and M2 are independent   | 1<br>1 |

Total 11 marks

|   |   | stion<br>nber |          | Answer  | Notes   | Marks |
|---|---|---------------|----------|---|---|-------|
| 4 | а | i             | M1       | bubbles / fizzing / effervescence<br>OR<br>solid/magnesium disappears/dissolves<br>OR | Allow just gas (given off) Ignore wrongly named gas   | 1     |
|   |   |               |          | flask gets warm   | Allow temperature increases but not heat produced   |       |
|   |   | ii            | M1       | magnesium chloride / MgCl <sub>2</sub>  | Accept phonetic spellings<br>Accept poorly written formulae such as<br>MGCl <sub>2</sub> and MgCL <sub>2</sub>  | 1     |
|   | b | i             | M1<br>M2 | $2H_2 + O_2 \rightarrow 2H_2O$  | correct formulae = 1 balancing = 1 Ignore heat anywhere Ignore state symbols  | 1     |
|   |   | ii            | M1       | condensation  | Accept phonetic spellings   | 1     |
|   | С | i             | M1       | blue  | Do not accept any other colours even in combination with blue, eg blue-green  Accept phonetic spellings Ignore qualifiers such as pale / dark / light Ignore mention of solution / liquid / solid | 1     |

| Question number |          | Answer   | Notes   | Marks |
|-----------------|----------|--|---|-------|
| 4 C ii          | M1<br>M2 | measure boiling point / melting/freezing point OR distil / boil / freeze 100 °C / 0 °C | Value must match property Accept ° or C in place of °C Do not award M2 if only value given Ignore evaporates M2 dependent on M1 | 1     |

**Total 8 marks** 

|   |   | stion<br>nber |    | Answer | Notes  | Marks |
|---|---|---------------|----|--------|--|-------|
| 5 | а | i             | M1 | S      | Accept diagram:  | 1     |
|   |   |               |    |        | H<br> <br>H —C — Br<br> <br>H  |       |
|   |   | ii            | M1 | T/U    | Accept diagrams:   | 1     |
|   |   |               |    |        | $\begin{array}{cccccccccccccccccccccccccccccccccccc$   |       |
|   |   | iii           | M1 | T/U    | Accept diagrams:   | 1     |
|   |   |               |    |        | H H H C=C H  Do not penalise if both T and U are given Do not award the mark if either or both of T or U is given and any other letter is included |       |

|   |   | stion<br>nber |          | Answer  | Notes  | Marks |
|---|---|---------------|----------|---|--|-------|
| 5 | b |               | M1       | (add) bromine (water)   | If bromide, then 0/2 Do not allow bromine in UV light, but M2 can be awarded   | 1     |
|   |   |               | M2       | decolourised / goes<br>colourless   | Ignore starting colour of bromine Ignore clear / discolours Reject bleached  | 1     |
|   | С |               | M1       | displayed formula of but-1-<br>ene, but-2-ene or<br>methylpropene   | All atoms and bonds must be shown<br>Allow dienes  | 1     |
|   | d | i             | M1       | $C_nH_{2n+2}$   | Accept x and other letters in place of n<br>Accept answers like C <sub>n</sub> H <sub>2n</sub> +2<br>Ignore brackets | 1     |
|   |   | ii            | M1       | same/similar chemical properties / reactions / behaviour / characteristics  | Ignore specific example such as react with oxygen Ignore similar (type of) reactivity                                |       |
|   |   |               | M2<br>M3 | gradation /gradual change /<br>trend / increase / decrease<br>of physical properties<br>(neighbouring members)<br>differ by CH <sub>2</sub> | Accept reference to specific property, eg boiling point Reject same / similar physical properties                    | 2     |
|   |   |               | M4       | same functional group   | Any two for 1 each Accept two answers on one answer line   |       |

| Question number |    | Answer  | Notes  | Marks |
|-----------------|----|---|--|-------|
| 5 e             | M1 | (compounds / molecules with) same molecular formula / same number of each type of atom different structures / structural formulae / atoms arranged differently / different displayed formulae | Ignore same chemical formula Ignore hydrocarbons If atoms or elements instead of compounds or molecules, max 1 for Q | 1     |

Total 11 marks

|   | Question number |    |    | Answer   | Notes   | Marks |
|---|-----------------|----|----|--|---|-------|
| 6 | а               | į  | M1 | H—O—H with both bonds represented by 2 shared electrons                        | Accept 2 dots, 2 crosses or 1 of each Atoms do not have to be labelled with H or O If wrongly labelled, only M1 can be awarded  | 1     |
|   |                 |    | M2 | 8 electrons in outer shell of O<br>AND<br>2 electrons in outer shell of both H | Ignore inner shell of O<br>Reject if H has 2 shells<br>M2 dependent on M1   | 1     |
|   |                 | ii | M1 | (strong electrostatic) attraction between bonding/shared pair of electrons     | Must refer to pair or two electrons   | 1     |
|   |                 |    | M2 | and nuclei (of hydrogen and oxygen)  | Accept word nucleus instead of nuclei if clear reference to 2 atoms 0/2 if any mention of ions / electron transfer M2 dependent on mention of both attraction and electrons in M1 | 1     |

|   |   | tion<br>ber |    | Answer  | Notes  | Marks |
|---|---|-------------|----|---|--|-------|
| 6 | b |             | M1 | idea of electron transfer / loss and gain of electrons                            |  | 1     |
|   |   |             | M2 | direction of transfer, eg sodium to oxygen / sodium loses and oxygen gains        |  | 1     |
|   |   |             | M3 | correct number of electrons involved, eg (each) sodium loses 1 and oxygen gains 2 | Ignore charges on ions  Ignore covalent 0/3 if any mention of electron sharing All marks may be scored on diagrams or by reference to electronic configurations Max 2 if molecules mentioned | 1     |
|   |   | ii          | M1 | (sodium) loses electron(s)  | Ignore oxygen gains electrons  | 1     |

| Question number |  |        | Answer   | Notes  | Marks |
|-----------------|--|--------|--|--|-------|
| 6 c             |  | M<br>1 | attractions between water molecules are weak(er) / easily overcome / need little energy to break                   | Allow (named) intermolecular forces in place of attractions  | 1     |
|                 |  | M<br>2 | attractions between (sodium and oxide) ions are strong(er) / ionic bonds are strong /need a lot of energy to break | Do not award M2 if any mention of intermolecular forces / metallic bonding Any implication of <u>breaking</u> covalent bonds = 0/2 | 1     |

|   | Question number |    |          | Answer                                       | Notes   | Marks  |
|---|-----------------|----|----------|--|---|--------|
| ( | ó d             | i  | M1       | S  |   |        |
|   |                 |    | M2<br>M3 | aq   |   | 2      |
|   |                 |    |          |  | All three correct = 2 marks Two correct = 1 mark One/none correct = 0 marks Do not award M1 for g or if not possible to be sure that it is s and not g Do not award marks for abbreviations such as sol / liq |        |
|   |                 | ii | M1<br>M2 | blue / purple<br>OH <sup>-</sup> / hydroxide | Allow indigo or violet<br>M1 and M2 independent   | 1<br>1 |

Total 14 marks

|   | Question number |    |        | Answer                        | Notes   | Marks |
|---|-----------------|----|--------|-------------------------------|---|-------|
| 7 | а               | i  | M<br>1 | Chlorine / /Cl <sub>2</sub>   | Allow CI Accept phonetic spellings Do not penalise poorly written formulae such as CL / cl / cL | 1     |
|   |                 |    | M<br>2 | Iodine / I <sub>2</sub>       | Allow I<br>Accept phonetic spellings  | 1     |
|   |                 | ii | M<br>1 | Astatine / At <sub>2</sub>    | Allow At Accept phonetic spellings Do not penalise poorly written formulae such as AT / at / aT | 1     |
|   | b               |    | M<br>1 | $H_2 + CI_2 \rightarrow 2HCI$ | correct formulae = 1<br>balancing = 1   | 1     |
|   |                 |    | M<br>2 |                               | Max 1 for symbol or formula error, eg HcL, Cl <sup>2</sup>                                      | 1     |

|   | Question number |    |          | Answer  | Notes   | Marks |
|---|-----------------|----|----------|---|---|-------|
| 7 | O               |    | M1<br>M2 | red<br>(hydrochloric) acid / hydrogen ions / H <sup>+</sup><br>(formed) | Ignore acidic and references to pH  | 1     |
|   |                 | ii | M1       | blue  | Allow no colour change Do not accept changes (from red) to blue   | 1     |
|   |                 |    | M2       | no reaction/acid/hydrogen ions/H <sup>+</sup> (formed)                  | Reject any reference to alkaline<br>Ignore not acidic and references to<br>pH<br>Ignore reference to not dissolving | 1     |

Total 9 marks

|   |   | tion<br>ber |    | Answer                            | Notes  | Marks |
|---|---|-------------|----|-----------------------------------|--|-------|
| 8 | а |             | M1 | exothermic                        | Accept phonetic spellings Do not accept endothermic or any spelling that could be taken as endothermic or a hybrid such as exdothermic | 1     |
|   | b | i           | M1 | volume of solution                | Allow amount of solution   | 1     |
|   |   |             | M2 | concentration (of solution)       |  | 1     |
|   |   |             | М3 | amount / mass of metal            | Allow quantity of metal  | 1     |
|   |   |             | M4 | same surface area of metal        | Allow same size pieces / same state of subdivision   | 1     |
|   |   |             | M5 | same (rate/time of) stirring      |  | 1     |
|   |   |             | M6 | same starting/initial temperature | Ignore references to room temperature Any two for 1 each   | 1     |
|   |   | ii          | M1 | 18.7(0)                           |  | 1     |
|   |   |             | M2 | 26.8(0)                           |  | 1 1   |
|   |   |             | МЗ | 8.1(0)                            | Conseq on M1 and M2  | 1     |
|   |   | iii         | M1 | Zn / zinc                         | Accept phonetic spellings  | 1     |
|   |   |             | M2 | x                                 |  | 1     |
|   |   |             |    |                                   |  |       |

|   | Question number |  |    | Answer  | Notes   | Marks |
|---|-----------------|--|----|---|---|-------|
| 8 | С               |  | M1 | $Zn + XSO_4 \rightarrow ZnSO_4 + X$                               | Ignore state symbols  | 1     |
|   | d               |  | M1 | would react with water OR forms insoluble calcium sulfate/product | Allow too reactive/very reactive/too high in the reactivity series Do not allow more reactive than other metals (in experiment) | 1     |

**Total 10 marks** 

|   | Question number |     |          | Answer   | Notes  | Marks  |
|---|-----------------|-----|----------|--|--|--------|
| 9 | а               | i   | M1       | air / atmosphere   |  | 1      |
|   |                 |     | M2       | water / natural gas / hydrocarbons                         | Allow methane  | 1      |
|   |                 | ii  | M1       | iron / Fe  | Ignore iron oxide Accept phonetic spellings Do not penalise other included numbers - eg Fe(II) / Fe(III) / Fe <sup>2+</sup> / Fe <sup>3+</sup> | 1      |
|   |                 | iii | M1       | 450 °C   | Accept temperature of 350°C to 550°C or temperatures in K If range given, both values must be within acceptable range                          | 1      |
|   |                 |     | M2       | 200 atm(ospheres)  | Accept pressure of 150 atm to 250 atm or pressures in Pa Unit needed for mark If two conditions given, both must be correct                    |        |
|   |                 | iv  | M1<br>M2 | cooled / temperature lowered ammonia liquefies / condenses | M1 and M2 are independent Do not award M2 if implication that other gases condense   | 1<br>1 |

|   |   | stion<br>nber |       | Answer  | Notes  | Marks |
|---|---|---------------|-------|---|--|-------|
| 9 | b |               | M1    | $n(N_2) = (56 \times 10^6) \div 28 / 2 \times 10^6$ | No penalty for missing or incorrect power of 10                | 1     |
|   |   |               | M2    | $n(NH_3) = M1 \times 2 / 4 \times 10^6$             | Conseq on M1   | 1     |
|   |   |               | МЗ    | $m(NH_3) = M2 \times 17 / 68 t (onnes)$             | Conseq on M2   | 1     |
|   |   |               |       |   | Correct final answer with units scores 3                       |       |
|   |   |               |       |   | Accept answers in grams and kilograms                          |       |
|   |   |               |       |   | 34 t scores 2 marks  |       |
|   |   |               |       |   | Final answer of 68 with missing or incorrect units scores 2    |       |
|   |   |               |       | OR  | M1 for 28 and 34 (need not be in this                          |       |
|   |   |               |       | 34 × 56   | expression)  |       |
|   |   |               |       | 28  | M2 is for expression shown                                     |       |
|   |   |               |       | = 68 t(onnes)                                       | M3 is for answer with units                                    |       |
|   |   |               |       | , ,   |  |       |
|   |   | (1)           | N 4 4 |   |  | 1     |
|   | С | (i)           | M1    | increased   | Allow long grows with American                                 | 1     |
|   |   |               | M2    | shift to left                                       | Allow less ammonia / products Allow moves in reverse direction | I     |
|   |   |               |       |   | Ignore reference to favouring                                  |       |
|   |   | (ii)          | M1    | shift to right                                      | Allow more ammonia / products                                  | 1     |
|   |   | (11)          | ''' ' | Sime to right                                       | Allow moves in forward direction                               | •     |
|   |   |               |       |   | Ignore reference to favouring                                  |       |
|   |   |               | M2    | fewer moles/molecules (of gas) on the right         | Allow more moles/molecules on the left                         | 1     |
|   |   |               |       |   | Do not penalise incorrect numbers, eg                          |       |
|   |   |               |       |   | 3 moles on the left and 2 moles on the                         |       |
|   |   |               |       |   | right  |       |
|   |   |               |       |   | Ignore references to rate                                      |       |
|   |   |               |       |   | M2 dependent on M1   |       |

|   | Question number |     |    | Answer   | Notes   | Marks |
|---|-----------------|-----|----|--|---|-------|
| 9 | d               | i   | M1 | 60   |   | 1     |
|   |                 | ii  | M1 | setting out correct division of each % by $A_{\rm r}$ OR 2.5, 5 and 3.75 | Award 0 for whole question if division by atomic numbers / wrong way up / multiplication used If molecular masses used for all three elements, no M1, but can award M2 and M3 | 1     |
|   |                 |     | M2 | division by smallest (gives 1 : 2 : 1.5)                                 | No penalty for subsequently rounding 1.5 to 2 if clear they have divided by smallest  | 1     |
|   |                 |     | M3 | $N_2H_4O_3$  | Accept elements in any order Allow NH <sub>4</sub> NO <sub>3</sub> If % O wrong or missing, only M1 and M2 can score  | 1     |
|   |                 | iii | M1 | ammonium nitrate   | Accept phonetic spellings Do not accept ammonia in place of ammonium Do not accept nitrite or nitride in place of nitrate Ignore all formulae                                 | 1     |

|    | Question number |    |          | Answer  | Notes  | Marks |
|----|-----------------|----|----------|---|--|-------|
| 10 | а               | i  | M1       | layers / sheets / planes / rows of (positive) ions  | Allow atoms/ particles in place of positive ions Reject molecules / protons / electrons  | 1     |
|    |                 |    | M2       | slide (over each other)   | Allow slip / flow / move in place of slide Accept explanation in terms of non-directional bonding Do not award M2 if protons / electrons Do not award M2 if no mention of layers or equivalent | 1     |
|    |                 | ii | M1<br>M2 | delocalised electrons / sea of electrons<br>move / flow (through structure) / mobile<br>(when voltage/potential difference applied) | Ignore free electrons M2 needs mention of electrons Any mention of ions moving = 0/2   | 1 1   |

|    | uesti<br>umb |    |          | Answer   | Notes   | Marks  |
|----|--------------|----|----------|--|---|--------|
| 10 | g            | į  | M1       | brown precipitate                                | Accept solid / suspension Ignore qualifiers such as pale / light / dark / muddy / dirty Ignore grey Ignore references to turning brown Reject bubbles or equivalent Do not penalise wrong identity of precipitate Accept solid / suspension Accept orange / orange-brown / red- brown Ignore qualifiers such as pale / light / dark Reject bubbles or equivalent Do not penalise wrong identity of precipitate Award 1 for both colours correct but precipitate missing | 1      |
|    |              | ii | M1<br>M2 | $FeSO_4 + 2NaOH \rightarrow Fe(OH)_2 + Na_2SO_4$ | Correct formulae = 1<br>Balancing = 1   | 1<br>1 |

Total 8 marks

|    | stio<br>nber |        |    | Answer  | Notes  | Marks |
|----|--------------|--------|----|---|--|-------|
| 11 | а            | M      | 11 | (total) volume different/not constant / not 50 / is 55                  | Allow too much water / sodium thiosulfate added / reference to numbers eg should be 10 instead of 15 or 35 instead of 40                     | 1     |
|    | b            | M<br>M | 12 | All six points plotted correctly to nearest gridline  curve of best fit | Deduct 1 mark for each error If plotting cannot be seen judge accuracy from the line. Do not award mark for joining dots                     | 2     |
|    |              | IVI    | 13 | <u>curve</u> or best fit  | or multiple lines or if all of the data points are completely misplotted   | '     |
|    | С            | M      |    | 1000 ÷ 26.6<br>37.6   | Ignore units M2 can be awarded for use of another student's result Award 2 marks for correct final answer Award 1 mark for 38 / 37.59 / 37.5 | 1 1   |

|    | iesti<br>umb |    |                | Answer   | Notes   | Marks       |
|----|--------------|----|----------------|--|---|-------------|
| 11 | d            | i  | M1<br>M2       | rate (directly) proportional to concentration                                    | Accept concentration (directly) proportional to rate Accept specific quantitative expression, eg rate doubles as concentration doubles Allow 1 mark for qualitative expression, rate increases as concentration increases | 2           |
| 11 | d            | ii | M1<br>M2<br>M3 | more particles / ions (in a given volume) collide (successfully) more frequently | Reject atoms / molecules  Reject with more energy Ignore greater chance of collision Must be reference to frequency or number of collisions per unit time Allow "increased frequency of collisions" for M2 and M3         | 1<br>1<br>1 |

Total 11 marks

www.dynamicpapers.com

Further copies of this publication are available from Edexcel Publications, Adamsway, Mansfield, Notts, NG18 4FN

Telephone 01623 467467 Fax 01623 450481 Email <u>publication.orders@edexcel.com</u> Order Code xxxxxxxxx January 2012

For more information on Edexcel qualifications, please visit <a href="https://www.edexcel.com/quals">www.edexcel.com/quals</a>

Pearson Education Limited. Registered company number 872828 with its registered office at Edinburgh Gate, Harlow, Essex CM20 2JE





