Write your name here Surname		Other names	
Pearson Edexcel International Advanced Level	Centre Number		Candidate Number
Chemistry Advanced Subsidiar Unit 2: Application of	ry	nciples	of Chemistry
Wednesday 17 January 201 Time: 1 hour 30 minutes	18 – Morning		Paper Reference WCH02/01
Candidates must have: Scient	ific calculator.		Total Marks

### Instructions

- Use **black** ink or **black** ball-point pen.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions.
- Answer the questions in the spaces provided
  - there may be more space than you need.

## Information

- The total mark for this paper is 80.
- The marks for **each** question are shown in brackets
  - use this as a guide as to how much time to spend on each question.
- Questions labelled with an asterisk (\*) are ones where the quality of your written communication will be assessed
  - you should take particular care with your spelling, punctuation and grammar, as well as the clarity of expression, on these questions.
- A Periodic Table is printed on the back cover of this paper.

## **Advice**

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.
- Show all your working in calculations and units where appropriate.

Turn over ▶







DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

#### **SECTION A**

Answer ALL the questions in this section. You should aim to spend no more than 20 minutes on this section. For each question, select one answer from A to D and put a cross in the box  $\boxtimes$ . If you change your mind, put a line through the box 🔀 and then mark your new answer with a cross  $\boxtimes$ .

- 1 Which is a polar molecule?
  - ☑ A BeCl₂
  - B BCI₃

  - ☑ D NCl,

(Total for Question 1 = 1 mark)

- 2 Which bond angles are present in a molecule of methanol?
  - A 90° and 104.5°
  - **■ B** 104.5° and 109.5°

  - **D** 90° and 180°

(Total for Question 2 = 1 mark)

- This question is about the hydrides of carbon, nitrogen, oxygen and fluorine.
  - (a) The hydride with the highest boiling temperature is

(1)

- A CH<sub>4</sub>
- B NH<sub>3</sub>
- ☑ D HF
- (b) The hydride which has the strongest hydrogen bond in the pure liquid is

(1)

- A CH<sub>4</sub>
- B NH,
- D HF

(Total for Question 3 = 2 marks)



X

X

X

**NOT WRITE IN THIS** 

DO NOT

4 On descending Group 2, from magnesium to barium, what are the trends in the first ionisation energy of the elements, and in the solubility of the sulfates?

	First ionisation energy	Solubility of sulfate
Α	increases	increases
В	increases	decreases
C	decreases	increases
D	decreases	decreases

(Total for Question 4 = 1 mark)

5 Flame tests are carried out on the chlorides of four Group 2 metals.
Select the metal chlorides that give these flame colours.

	Flame colour				
	Colourless Crimson Pale green			Yellow-red	
⊠ A	magnesium	calcium	strontium	barium	
⊠ B	barium	calcium	magnesium	strontium	
⊠ C	barium	strontium	magnesium	calcium	
⊠ D	magnesium	strontium	barium	calcium	

(Total for Question 5 = 1 mark)

- 6 The s-block metal nitrate that decomposes on heating to form a nitrite is
  - **A** lithium nitrate.
  - **B** sodium nitrate.
  - **C** magnesium nitrate.
  - **D** calcium nitrate.

(Total for Question 6 = 1 mark)

Use this space for any rough working. Anything you write in this space will gain no credit.

DO NOT WRITE IN THIS AREA

			(Total for Question 10 = 1 mark
	X	D	H <sub>2</sub> O
	×	C	NO <sub>2</sub>
	×	В	CO <sub>2</sub>
	×	A	СО
10	Th	e gr	eenhouse gas with the highest mean concentration in the atmosphere is
		_	(Total for Question 9 = 1 mark
			sulfur trioxide.
		С	sulfur. sulfur dioxide.
	X	В	hydrogen sulfide. sulfur.
9		duce	concentrated sulfuric acid reacts with solid potassium bromide, sulfuric acid is ed to
			(Total for Question 8 = 1 mark
	X	D	sulfuric acid.
	×	C	phosphoric(V) acid.
	×	В	nitric acid.
	X	A	hydrochloric acid.
8	Th	e be	est way to prepare hydrogen iodide from potassium iodide is to add concentrated
			(Total for Question 7 = 1 mark
	×	D	lodine is a grey solid that dissolves in hexane to form a brown solution.
	X	c	lodine is a brown liquid that dissolves in hexane to form a pink solution.
	×	В	Chlorine is a pale green gas that dissolves in hexane to form a pale green solution
	$\times$	Α	Chlorine is a pale green gas that dissolves in hexane to form a brown solution.


DO NOT WRITE IN THIS AREA

⊠ A	ion-dipole.		
ВВ	ion-ion.		
⊠ C	dipole-dipole.		
⊠ D	hydrogen bonding.		
		(Total	for Question 11 = 1 mark)
<b>2</b> What a	re the properties of the	liquid 2-chlorobutane?	
	Solubility in water	Effect of a charged rod on a stream of the liquid	
⊠ A	insoluble	stream diverted	
<b>⊠</b> B	insoluble	stream unaffected	
⊠ C	soluble	stream diverted	
	Soldbic		
⊠ D	soluble	stream unaffected	
			for Question 12 = 1 mark)
3 How m concen	soluble  nany organic elimination ntrated solution of potas  1 2 3	(Total products form when 2-bromob sium hydroxide in ethanol?	outane is heated with a
<ul><li>☑ D</li><li>3 How m concen</li><li>☑ A</li><li>☑ B</li><li>☑ C</li></ul>	soluble  nany organic elimination ntrated solution of potas  1 2 3	(Total products form when 2-bromob sium hydroxide in ethanol?	
■ D  How m concen ■ A ■ B ■ C ■ D	soluble  nany organic elimination ntrated solution of potas  1 2 3 4	(Total products form when 2-bromobsium hydroxide in ethanol? (Total with the formula $C_4H_9OH$ , would	for Question 13 = 1 mark)
3 How m concen  A B C D  Which is peak do	soluble  nany organic elimination ntrated solution of potas  1 2 3 4  two isomeric alcohols, w	(Total products form when 2-bromobsium hydroxide in ethanol? (Total with the formula $C_4H_9OH$ , would ass spectra?	for Question 13 = 1 mark)
■ D  3 How m concen ■ A ■ B ■ C ■ D  4 Which to peak do ■ A	soluble  nany organic elimination ntrated solution of potas  1  2  3  4  two isomeric alcohols, we ue to CH <sub>2</sub> OH <sup>+</sup> in their materials	(Total products form when 2-bromobsium hydroxide in ethanol?  (Total vith the formula $C_4H_9OH$ , would ass spectra?	for Question 13 = 1 mark)
■ D  3 How m concen ■ A ■ B ■ C ■ D  4 Which to peak do ■ A ■ B	soluble  nany organic elimination ntrated solution of potas  1  2  3  4  two isomeric alcohols, we ue to CH <sub>2</sub> OH <sup>+</sup> in their many butan-1-ol and 2-methy	(Total products form when 2-bromobsium hydroxide in ethanol?  (Total vith the formula $C_4H_9OH$ , would ass spectra?	for Question 13 = 1 mark)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

**15** Part of the infrared (IR) spectrum of a compound is shown.



Bond	Wavenumber range / cm <sup>-1</sup>
O—H (alcohol)	3750–3200
O—H (carboxylic acid)	3300–2500
C—H (alkane)	2962–2853
C—H (aldehyde)	2900–2820 and 2775– 2700
C=O (aldehyde or ketone)	1740–1680

The compound could be

- A propan-1-ol.
  - **B** propanoic acid.
- C propanal.
- **D** propanone.

(Total for Question 15 = 1 mark)

DO NOT WRITE IN THIS AREA

<b>16</b> Two a	cohols are oxidised under mild conditions.
	alcohols each form a compound that gives a red precipitate on heating with Benedict's solution or Fehling's solution.
These	alcohols could be
⊠ A	propan-1-ol and propan-2-ol.
	propan-1-ol and butan-1-ol.
	propan-2-ol and butan-2-ol.
⊠ D	butan-1-ol and butan-2-ol.
	(Total for Question 16 = 1 mark)
	ic compounds which react with sodium but are <b>not</b> oxidised by ed potassium dichromate(VI) are
⊠ A	primary alcohols.
⊠ B	secondary alcohols.
<b>⊠</b> C	tertiary alcohols.
⊠ D	ketones.
	(Total for Question 17 = 1 mark)
<b>18</b> Which	statement about the carbon footprint of fuels is true?
⊠ A	Hydrogen has a zero carbon footprint as it does not produce carbon dioxide.
⊠ B	Methane has a zero carbon footprint as it occurs naturally.
⊠ C	Biodiesel has a zero carbon footprint as it absorbs as much carbon dioxide in production as it produces in combustion.
⊠ D	No fuel has been discovered with a zero carbon footprint.



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

19 Dinitrogen tetroxide and nitrogen dioxide form an equilibrium mixture in a gas syringe.

 $N_2O_4(g) \rightleftharpoons 2NO_2(g)$ Pale brown Dark brown

The pressure is rapidly doubled and then the mixture allowed to stand.

The colour would

- ☑ A go darker then go paler.
- **B** go darker and remain darker.
- **C** go paler and remain paler.
- **D** go paler then go darker.

(Total for Question 19 = 1 mark)

**TOTAL FOR SECTION A = 20 MARKS** 



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

**BLANK PAGE** 



DO NOT WRITE IN THIS AREA

#### **SECTION B**

# Answer ALL the questions. Write your answers in the spaces provided.

**20** This question is about the preparation and properties of 1-iodobutane.

- (a) 1-iodobutane is prepared by warming a mixture of damp red phosphorus with iodine to produce phosphorus(III) iodide,  $PI_3$ . This reacts with butan-1-ol to form 1-iodobutane,  $C_4H_9I$ .
  - \*(i) Draw a diagram to show the shape of phosphorus(III) iodide. Predict the I—P—I bond angle.

Explain why the molecule has this shape and bond angle.

(4)

Diagram

Bond angle .....

Explanation	 	 	

(ii) Complete the balanced equation for the formation of 1-iodobutane. State symbols are not required.

(1)

.....C<sub>4</sub>H<sub>9</sub>OH + PI<sub>3</sub> 
$$\rightarrow$$

DO NOT WRITE IN THIS AREA

	(iii) Draw <b>skeletal</b> formulae of the four structure.	ctural isomers of $C_4H_9I$ . (2)
_		
b)	<ul> <li>1-iodobutane, dissolved in ethanol, reacts value a yellow precipitate. The reaction involves to the first step, 1-iodobutane forms but Identify the attacking reagent, and state</li> </ul>	two steps.
		(2)
:ki	ng reagent	(2)
	ng reagentnd mechanism of this reaction	
	nd mechanism of this reaction(ii) Write the <b>ionic</b> equation for the formati	
	nd mechanism of this reaction	
	nd mechanism of this reaction(ii) Write the <b>ionic</b> equation for the formati	on of the yellow precipitate.
e ai	nd mechanism of this reaction(ii) Write the <b>ionic</b> equation for the formati Include state symbols.	on of the yellow precipitate.
aı	nd mechanism of this reaction(ii) Write the <b>ionic</b> equation for the formati Include state symbols.	on of the yellow precipitate.
e ai	nd mechanism of this reaction(ii) Write the <b>ionic</b> equation for the formati Include state symbols.	on of the yellow precipitate. (1)



21 This question is about nitrogen monoxide, NO.	
(a) Nitrogen monoxide is formed in internal combustion engines.	
$N_2(g) + O_2(g) \rightleftharpoons 2NO(g)$ $\Delta H_{298}^{\oplus} = +180 \text{ kJ mol}^{-1}$	
Explain how, if at all, an increase in temperature and an increase in pressure affect	
this equilibrium. Justify your answers.	(3)
(b) In industry, nitrogen monoxide is produced by the oxidation of ammonia at high temperature, with a platinum catalyst.	
$4NH_3(g) + 5O_2(g) \rightarrow 4NO(g) + 6H_2O(g)$	

First element

..... to .....

Second element \_\_\_\_\_ from \_\_\_\_ to \_\_\_\_

reaction. State the relevant oxidation numbers.

(ii) Use the Maxwell-Boltzmann distribution to explain why increasing the temperature will result in a higher rate for this reaction. A diagram is not required.

(i) Identify the two elements which change their oxidation number in this

(1)

(2)

DO NOT WRITE IN THIS AREA

	(iii) Use the Maxwell-Boltzmann distribution to explain why the platinum catalyst produces a higher rate for this reaction. A diagram is not required.	(1)
(c)	Nitrogen monoxide is a major pollutant. High in the atmosphere, it is a greenhouse gas and it depletes the ozone layer.  (i) Explain why nitrogen monoxide is a greenhouse gas and how the presence of nitrogen monoxide in the atmosphere leads to global warming.	(3)
	(ii) Write <b>two</b> equations to show how the free radical, nitrogen monoxide, depletes the ozone layer. Indicate free radicals in the usual way. Hence write the equation which shows the overall change taking place.	
	State symbols are not required.	(3)
	(Total for Question 21 = 13 ma	rks)



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

- **22** Potassium iodate(V), KIO<sub>3</sub>, is made by adding iodine to boiling concentrated potassium hydroxide solution.
  - (a) (i) Balance the equation for the reaction.

(2)

(ii) State the type of redox reaction between iodine and concentrated potassium hydroxide.

(1)

(b) What would you **see** when a slight excess of iodine has been added?

(1)

(c) Potassium iodate(V) crystallises as the solution cools.

Suggest why potassium iodate(V), rather than potassium iodide, crystallises out.

(1)

DO NOT WRITE IN THIS AREA

(d)	The purity of the potassium iodate(V) formed is determined using the method	
	outlined below.	
	$0.100\mathrm{g}$ of the potassium iodate(V) sample is dissolved in distilled water and the volume made up to $100\mathrm{cm}^3$ .	
	A 10.0 cm <sup>3</sup> portion is taken and added to an excess of a mixture of potassium iodide in dilute sulfuric acid.	
	The iodine formed is titrated with 0.0100 mol dm <sup>-3</sup> sodium thiosulfate solution.	
	The titration is repeated and the mean titre is 27.45 cm <sup>3</sup> .	
	(i) Name the indicator that should be used for the titration and state when it should be added to the reaction mixture.	(2)
		(2)
	(ii) Give the colour change for the indicator at the end-point.	
	(ii) Give the colour change for the indicator at the end-point.	(1)
	From to	
	(iii) Calculate the number of moles of thiosulfate ions used in the titration.	
	(III) Calculate the number of moles of thiosulfate ions used in the titration.	(1)
	(iv) Calculate the number of moles of potassium iodate(V) in the 10.0 cm <sup>3</sup> portion,	
	given that 6 mol of thiosulfate ions is equivalent to 1 mol of iodate(V) ions.	(1)



DO NOT WRITE IN THIS AREA

(v) Calculate the mass of potassium iodate(V) in the original sample.

(3)

(vi) Calculate the percentage purity by mass of potassium iodate(V) in the original sample. Give your answer to **two** significant figures.

(2)

(vii) Suggest why the potassium iodate(V) obtained is not 100% pure.

(1)

(Total for Question 22 = 16 marks)

**TOTAL FOR SECTION B = 41 MARKS** 



AREA

**NOT WRITE IN THIS** 

NOT WRITE IN THIS AREA

#### **SECTION C**

## Answer ALL the questions. Write your answers in the spaces provided.

**23** Glucose occurs naturally in many fruits. It is a white powder at room temperature and is extremely soluble in water. Glucose may be represented by the structure below.

Glucose

The fermentation of glucose is fundamental to brewing and baking. Glucose breaks down to form carbon dioxide and ethanol.

Drinks with a high alcohol content are obtained by distillation from a fermentation mixture.

For many years, the alcohol content of such drinks was measured as degrees proof. Originally this was defined by the gunpowder test. A pellet of gunpowder was soaked in the drink. If the gunpowder would still ignite, the alcohol drink was at least 100° proof. The reason for introducing this measure was that, from the sixteenth century, the tax on alcoholic drinks was related to their alcohol content.

Nowadays, most countries have adopted alcohol percentage by volume (ABV), which is the volume of ethanol, in cm<sup>3</sup>, present in 100 cm<sup>3</sup> of the drink.

Today, most ethanol for chemical use is produced by an addition reaction of ethene.

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

*(a) (i)	Name all the intermolecular forces between glucose molecules. For each type of force, indicate the atoms in the molecule involved.								
	A detailed explanation of how these forces arise is <b>not</b> required.	(6)							
(ii) Explain why glucose is very soluble in water.									

(b) Complete the equation for the fermentation of glucose. State symbols are not required.

$$C_6H_{12}O_6 \rightarrow$$

DO NOT WRITE IN THIS AREA

(c) Suggest <b>two</b> advantages for the taxation of alcoholic drinks.	(2)
(d) The ABV in a 100° proof drink is found to be 57.15%.	
(i) Calculate the degrees proof of pure ethanol.	(1)
(ii) Calculate the concentration of ethanol, in $mol  dm^{-3}$ , in a solution when the ABV is 57.15%. [Density of ethanol = $0.789  g  cm^{-3}$ ]	(3)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(e) Potassium nitrate is the main ingredient of gunpowder. Suggest how the gunpowder test for measuring the degrees proof of alcohol drinks works.

(1)

(f) Balance this simplified equation for the decomposition of gunpowder.

(1)

......KNO<sub>3</sub>(s) + S(s) + .....C(s) 
$$\rightarrow$$
 K<sub>2</sub>S(s) + N<sub>2</sub>(g) + .....CO<sub>2</sub>(g)

(g) Write the equation, including state symbols, for the formation of ethanol from ethene and suggest conditions for the industrial preparation.

(2)

(Total for Question 23 = 19 marks)

TOTAL FOR SECTION C = 19 MARKS TOTAL FOR PAPER = 80 MARKS

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

# **BLANK PAGE**



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

# **BLANK PAGE**



	0 (8)	4.0 <b>He</b> helium 2	20.2	Ne	neon 10	39.9	Ar	argon 18	83.8	고	krypton 36	131.3	Xe	xenon 54	[222]	R	radon 86		_					
	7	h (77)	19.0	ıL	fluorine 9		ַ		6.62		bromine k	126.9		fodine 53	[210]		astatine 85		Etenients with atomic numbers 112-110 have been reported but not fully authenticated	175		lutetium 71	[257]	<b>Lr</b> lawrencium 103
	9	(16)	16.0	0	oxygen 8	32.1		16	79.0		Ε	127.6	Б	tellurium 52	[509]	8	polonium 84	46 1000	ticated	173		ytterbium 70	[254]	No nobelium la 102
	2	(15)	14.0	z	nitrogen 7	31.0	Δ -	phosphorus 15	74.9	As	arsenic 33	121.8	Sb	antimony 51	209.0	Bi	bismuth 83	4	but not fully authenticated	169		thulium 69	[526]	Md mendelevium 101
	4	(14)	12.0	U	carbon 6	28.1		14	72.6	ge	germanium 32	118.7	Sn	50 ti	207.2	Ъ	lead 82		but not fi	167	ם	erbium 68	[253]	Fm fermium 100
	m	(13)	10.8	В	boron 5	27.0	IV.	atuminium 13	2.69	Ga	gallium 31	114.8	Г	indium 49	204.4	F	thallium 81	44	ieirs with	165		holmium 67	[254]	<b>Es</b> einsteinium 99
ents								(12)	65.4	Zn	zinc 30	112.4	Ъ	cadmium 48	200.6	Hg	mercury 80			163	Dy	dysprosium 66	-	Cf Es californium einsteinium 98 99
The Periodic Table of Elements						(11)			63.5	J	copper 29	107.9	Ag	silver 47	197.0	Αn	plog 79	[272]	roentgenium 111	159		terbium 65	[245]	<b>BK</b> berkelium 97
le of					(10)				58.7	ź	nickel 28	106.4	Pd	palladium 46	195.1	£	platinum 78	[271]	meitnerium damstadtium roentgenium	157		gadolinium 64	[247]	Cm curium 96
c Tab						(6)			58.9	ပိ	cobalt 27	102.9		rhodium 45	192.2	ŀ	iridium 77	[268]	meitnerium 109	152	Eu	samarium europium 62 63	[243]	Am americium 95
riodi		1.0 <b>H</b> hydrogen						(8)	55.8	Fe	0	101.1	Ru	molybdenum technetium ruthenium 42 44	190.2	o	osmium 76	[277]	hassium 108	150	Sm	samarium 62	[242]	Np Pu Am neptunium plutonium americium 93 94 95
he Pe							(2)		54.9	Wn	chromium manganese 24 25	[86]	2	technetium 43	186.2	Re	rhenium 75	_		[147]	Pm	praseodymium promethium 59 60 61	[237]	Np neptunium 93
<b>—</b>			: mass	atomic symbol	name atomic (proton) number			(9)	52.0		chromium 24	95.9	Wo	motybdenum 42	183.8	≯	tungsten 74	[566]	seaborgium 106	144	PN	neodymium 60	238	U uranium 92
		Key	relative atomic mass					(2)	50.9	>	vanadium 23	92.9	Q.	niobium 41	180.9	Ta	tantalum 73	[292]	dubnium 105	141	P	ргазеодутіцт 59	[231]	Pa protactinium 91
			relat	ato	atomic			(4)	47.9	ï	ţţ	91.2	Zr	zirconium 40	178.5	Ξ	hafnium 72	[261]	nutherfordium 104	140	e O	cerium 58	232	thorium 90
			_					(3)	45.0	Sc	scandium 21	88.9		yttrium 39	138.9	La*	lanthanum 57	[227]	actinium 89		es			
	2	(2)	0.6	Be	beryllium 4	24.3	Wg	magnesium 12	40.1	Ca	ŭ	97.8		strontium 38	137.3	Ba	barium 56	[326]	radium 88		* Lanthanide series	* Actinide series		
	- 8		6.9 Li		lithium 3	23.0	Na sodium		39.1	¥	potassium 19	85.5	В	rubidium 37	132.9	ర	caesium 55	[223]	francium 87		* Lant	* Actir		

