

Cambridge IGCSE

Cambridge International Examinations

Cambridge International General Certificate of Secondary Education (9–1)

CANDIDATE NAME									
CENTRE NUMBER						DIDATE 1BER	Ξ		

CHEMISTRY

0971/04

Paper 4 Theory (Extended)

For Examination from 2018

SPECIMEN PAPER

1 hour 15 minutes

Candidates answer on the Question Paper.

No Additional Materials are required.

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams, graphs or rough working.

Do not use staples, paper clips, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

Answer all questions.

Electronic calculators may be used.

You may lose marks if you do not show your working or if you do not use appropriate units.

A copy of the Periodic Table is printed on page 20.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.



1 The following table gives information about six substances.

substance	melting point / °C	boiling point / °C	electrical conductivity as a solid	electrical conductivity as a liquid
Α	839	1484	good	good
В	-188	-42	poor	poor
С	776	1497	poor	good
D	-117	78	poor	poor
E	1607	2227	poor	poor
F	– 5	102	poor	good

(a)	which substance could be a metal?	[1]
(b)	State all the substances that are liquid at room temperature.	- 4 -
(c)	Which substance could have a macromolecular structure similar to that of silicon(IV) oxide	[1] ə?
		[1]
(d)	Which substance could be propane?	F41
		[1]
(e)	Which substance could be sodium chloride?	
		[1]
	[Total	l: 5]

[Total: 6]

2 The table gives the composition of three particles.

particle	number of protons	number of electrons	number of neutrons
Α	15	15	16
В	15	18	16
С	15	15	17

(a)	Wh	at is the evidence in the table for each of the following?	
	(i)	Particle A is an atom.	
	(ii)	A, B and C are all particles of the same element.	[1]
			[1]
	(iii)	Particles A and C are isotopes of the same element.	
			[2]
(b)	(i)	What is the electronic structure of particle A ?	[1]
	(ii)	Is element A , a metal or a non-metal? Give a reason for your choice.	
			[1]

[2]


3	Kinetic theory	explains t	he properties	of	matter	in	terms	of	the	arrangement	and	movement	of
	particles.												

(a)	Nitrogen	is a	gas	at	room	temperature.	Nitrogen	molecules,	N_2 ,	are	spread	far	apart	and
	move in a	a ran	dom	ma	nner a	at high speed.								

(i)	Draw the electronic structure of a nitrogen molecule
	Show only the outer electron shells.

	(ii)	Compare the movement and arrangement of the molecules in solid nitrogen to those in nitrogen gas.
		[3]
(b)	the Use	ealed container contains nitrogen gas. The pressure of the gas is due to the molecules of gas hitting the walls of the container. The the kinetic theory to explain why the pressure inside the container increases when the aperature is increased.
		[2]

The following apparatus can be used to measure the rate of diffusion of a gas.



The following results were obtained.

gas	temperature /°C	rate of diffusion in cm³/min
nitrogen	25	1.00
chlorine	25	0.63
nitrogen	50	1.05

(c) (i) Explain why nitrogen gas diffuses faster than chlorine gas.

		[2]
(ii)	Explain why the nitrogen gas diffuses faster at the higher temperature.	
		[1]
	[Total:	10]


Chror	nium is a transition element.
	i) State two differences in the physical properties of chromium and sodium.
	[2
(i	
	[2
(b) C	Chromium is used to electroplate steel objects. The diagram shows how this could be done.
	lead anode object to be plated chromium(III) sulfate(aq)
(i) Give two reasons why steel objects are plated with chromium.
	[2
(i	i) The formula of the chromium(III) ion is Cr^{3+} and of the sulfate ion is SO_4^{2-} . Give the formula of chromium(III) sulfate.
	[1
(ii	i) Write the ionic half-equation for the reaction at the negative electrode (cathode).
	[2
(iv	 A colourless gas, which relights a glowing splint, is formed at the positive electrode (anode).
	State the name of this gas.

www.dynamicpapers.com

7

(v)	During electroplating, it is necessary to add more chromium(III) sulfate but during copper plating using a copper anode, it is not necessary to add more copper(II) sulfate.
	Explain this difference.
	[2]
	[Total: 12]

5 Iron is extracted from its ore, hematite, in the blast furnace.



Describe the reactions involved in this extraction.

Include one equation for a redox reaction and one for an acid/base reaction.
[5]

[Total: 5]

9

BLANK PAGE

6	Soluble salts	can be	made	using a	base	and an	acid

(a)	Complete this method of preparing dry crystals of the soluble salt
	cobalt(II) chloride-6-water from the insoluble base $cobalt(II)$ carbonate

step 1 Add an excess of cobalt(II) carbonate to hot dilute hydrochloric acid.	
step 2	
step 3	
	••••
step 4	
	[4]

[Total: 10]

(b) (i) 5.95 g of cobalt(II) carbonate were added to 40 cm³ of hydrochloric acid, concentration 2.0 mol/dm³.

Calculate the maximum yield of cobalt(II) chloride-6-water and show that the cobalt(II) carbonate was in excess.

$$CoCO_3 + 2HCl \rightarrow CoCl_2 + CO_2 + H_2O$$

 $CoCl_2 + 6H_2O \rightarrow CoCl_2.6H_2O$

maximum yield:

	number of moles of HC1 used =
	number of moles of CoCl ₂ formed =
	number of moles of $CoCl_2.6H_2O$ formed =
	mass of one mole of $CoCl_2.6H_2O = 238g$
	maximum yield of CoCl ₂ .6H ₂ O =g
	to show that cobalt(II) carbonate is in excess:
	number of moles of HCl used = (use your value from above)
	mass of one mole of $CoCO_3 = 119g$
	number of moles of CoCO ₃ in 5.95g of cobalt(II) carbonate =[5]
(ii)	Explain how these calculations show that cobalt(II) carbonate is in excess.
	[1]

7 Iodine reacts with chlorine to form dark brown iodine monochloride.

$$I_2 + Cl_2 \rightarrow 2ICl$$

This reacts with more chlorine to give yellow iodine trichloride. An equilibrium forms between these iodine chlorides.

$$ICl(I) + Cl_2(g) \rightleftharpoons ICl_3(s)$$

dark brown yellow

(a)	Wha	at do you understand by the term equilibrium?	
			[2]
(b)		en the equilibrium mixture is heated, it becomes a darker brown colour. Igest if the reverse reaction is endothermic or exothermic. Give a reason for your choice	
		l	[1]
(c)	The	pressure on the equilibrium mixture is decreased.	
	(i)	How would this affect the position of equilibrium? Give a reason for your choice.	
		It would move to the	
		reason	
			[1]
	(ii)	Describe what you would observe.	

(d) Calculate the overall energy change for the reaction between iodine and chlorine using the bond energy values shown.

$$I_2 + Cl_2 \rightarrow 2ICl$$

Bond	Energy / kJ per mol
I–I	151
C <i>l</i> –C <i>l</i>	242
I–C <i>l</i>	208

Show your working.

[3]

(e) Draw a labelled energy level diagram for the reaction between iodine and chlorine using the information in (d).

[2]


[Total: 10]

The alcohols form an homologous series.
(a) Give three characteristics of an homologous series.
[3]
(b) The following two alcohols are members of an homologous series and they are isomers.
$CH_3 - CH_2 - CH_2 - CH_2 - OH$ and $(CH_3)_2CH - CH_2 - OH$
(i) Explain why they are isomers.
[2]
(ii) Deduce the structural formula of another alcohol which is also an isomer of these alcohols.

[1]

[1]

(c) Copper(II) oxide can oxidise butanol to liquid \mathbf{X} , whose pH is 4.



(i) Give the name of another reagent which can oxidise butanol.


		[1]
(ii)	Which homologous series does liquid X belong to?	
		[1]
iii)	State the formula of liquid X .	

[Total: 14]

(d) The alcohol ethanol can be made by fermentation. Yeast is added to aqueous glucose.

$$C_6 H_{12} O_6(aq) \ \to \ 2 C_2 H_5 OH(aq) \ + \ 2 CO_2(g)$$

Carbon dioxide is given off and the mixture becomes warm, as the reaction is exothermic. The graph shows how the rate of reaction varies over several days.



(i)	Suggest a	method (of measi	uring the	rate	of this	reaction.

		[2]
(ii)	Why does the rate initially increase?	
		[1]
(iii)	Suggest two reasons why the rate eventually decreases.	
		[2]

17

BLANK PAGE

rne	ere are two types of polymerisation, addition and condensation.	
(a)	Explain the difference between these two types of polymerisation.	
		[2]
		ر کا
(b)	Some plastics, formed by polymerisation, are non-biodegradable.	
	Describe two pollution problems that are caused by non-biodegradable plastics.	
		••••••
		[2]

(c) The polymer known as PVA is used in paints and adhesives. Its structural formula is shown below.

Deduce the structural formula of its monomer.

[1]

(d) A condensation polymer can be made from the following monomers.

Draw the structural formula of this polymer.

[3]

[Total: 8]

								Group	gn								
-	=											≡	2	>	5	5	=
							,										2
							I										He
				Kon			нуфлодил										halium
65	,	_	de	atomic number	[_	-					4	e	7	æ	đ	5
	- a		40	onio onno	7 7							0	, (- 2		ь	No.
3	90		also	atornic symbol	500							0	٠.	Ζ,		- 3	2
7	the party		relan	ramar margine adomir mass	2255							11	12	14	16	10	8
=	12											13	4	16	9	17	18
Na	M											Ą	S	۵	S	õ	Ā
multus 99	magnetalem 2.4											ahminium	mpgs.	phosphorus	aufter 9.5	dilonne	ucture
65	20 2	21	22	23	24	25	88	27	28	59	98	3 5	32	33	3	35	36
¥	Ö	S	F	>		M	Đ.	ပိ	ž	3	Zu	Ga	Ge	As	Se	Ä	Ż
potassium		seandam	Standard	wanadum	5	напдалеве	inori	cobst	rigge	saddeo	Sinc	gallum	permanium	amenic	selectum	bromine	hryphon
39	40	45	48	51	52	55	8	20	69	99	99	02	73	75	7.9	80	88
37	38	33	40	41	Г	43	4	45	46	47	48	49	20	15	252	53	\$
&	ഗ്	>	Zr	g	Mo	J _C	Z	문	P	Ag	S	Ę	Sn	Sb	ъ Д	П	Xe
ndsidum	stronbum	ytrum	zircznium	risbium	Ę	technolom	nzhankm	тороф	mulpolloq	shur	cadmium	inclum	Ę	artimony	Salbutum	indini	XOREGE
92	88	88	9	88	╗	1	101	103	106	108	112	116	119	122	128	127	131
99	99	57-71	72	23		75	92	77	7.8	78	90	18	82	8	8	85	98
S	Ba	lanthanoids	Ξ	_a ⊣a		Se.	ő	'n	ă	Au	Ď	F	ЬЬ	ē	Po	¥	瀀
Catchum	berium		hafrium	lartahm	6	manim	mrama	moja	platinum	pod	metouty	frallum	bead	biamuth	polonium	astalne	recon
133	137		1/8	181	T	186	38	182	195	187	201	504	707	502	1	1	1
87	88	89-103	9	8		107	8	8	110	111	112		114		116		
Ľ.	Sa.	401roids	抷	8		늄	£	ž	S	Bg	5		FZ		^		
frandum	radkırı		ntherfordum	abrium	seaborgam	pohilum	hassum	megneunu	ametachun	noentgerium	офетель		Jerovium		Peemorium		
		57	58	29	99	19	62	63	64	65	99	29	89	69	70	7	
lanthanoids	spice	Гa	පී	Ā		Pm	Sm	ū	В	2	ò	운	ш	Тщ	Ϋ́	3	
		lanthanum 430	mread	prancipaline 141	neotymium 144	promethium	samanum 150	europium	gadolinium 457	150	dysprosium 4.8.9	holmium	eftkum 197	tollum 180	yterbium 479	Mothum 475	
		88	06	2	П	88	3	36	96	97	86	8	100	101	102	183	
actinoids	nt.	Ac	£	Pa		Š	P	Am	S	ă	Ö	es E	E	PM	°	_	
		actrium	mount	protectinum	E	neptunism	phronium	americkum	ornum	perform	callfornium	einsteinum	Semium	тегдевунт	nobelum	lawrendum	
		(232	231	238	1	1	1	1	1	ť	ť	1	ť	1	ı	
The vol.	The volume of one mole of any gas is 24 dm3 at room t	e mole of	any gas	is 24 dm ³	at room	emperatu	a pue aur	emperature and pressure (r.t.p.)	r.t.p.)								

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.